

The Influence of Spinous Process Union on Clinical Outcomes After Spinous Process Osteotomy for Lumbar Spinal Stenosis After 2 Years: A Secondary Analysis From the NORDSTEN-Study

Sondre Hagerup, Jens Ivar Brox, Hasan Banitalebi, Kari Indrekvam, Tor Åge Myklebust and Erland Hermansen

Int J Spine Surg 2024, 18 (1) 47-53

doi: <https://doi.org/10.14444/8576>

<https://www.ijssurgery.com/content/18/1/47>

This information is current as of May 4, 2025.

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://ijssurgery.com/alerts>

The Influence of Spinous Process Union on Clinical Outcomes After Spinous Process Osteotomy for Lumbar Spinal Stenosis After 2 Years: A Secondary Analysis From the NORDSTEN-Study

SONDRE HAGERUP, MD¹; JENS IVAR BROX, MD, PhD^{2,3}; HASAN BANITALEBI, MD^{2,4}; KARI INDREKVAM, MD, PhD^{5,6}; TOR ÅGE MYKLEBUST, PhD^{7,8}; AND ERLAND HERMANSEN, MD, PhD^{1,5,6}

¹Department of Orthopedic Surgery, Ålesund Hospital, Møre and Romsdal Hospital Trust, Norway, Europe; ²Institute of Clinical Medicine, University of Oslo, Oslo, Norway, Europe; ³Department of Physical Medicine and Rehabilitation, Oslo University Hospital, Norway, Europe; ⁴Department of Diagnostic Imaging, Akershus University Hospital, Norway, Europe; ⁵Kysthospitalet i Hagevik, Orthopedic Clinic, Haukeland University Hospital, Bergen, Norway, Europe; ⁶Department of Clinical Medicine, University of Bergen, Bergen, Norway, Europe; ⁷Department of Research and Innovation, Møre and Romsdal Hospital Trust, Ålesund, Norway, Europe; ⁸Department of Registration, Cancer Registry of Norway, Oslo, Norway, Europe

ABSTRACT

Background: Lumbar spinal stenosis is a prevalent and increasingly important cause of low back pain, leg pain, and walking impairment. Minimally invasive decompressive techniques such as spinous process (SP) osteotomy have become more common in recent years. The main aim of this study was to investigate the proportion of complete SP union and whether complete radiological healing after the osteotomy is associated with superior clinical outcome after 2 years.

Methods: In this retrospective cohort study, 149 patients were included from the Spinal Stenosis Trial, a part of the Norwegian Degenerative spondylolisthesis and spinal STENosis study. Computed tomography imaging was performed 2 years postoperatively. The number of osteotomies and the number of SP unions were recorded. Patients were divided into groups based on the degree of union: nonunion, partial union, and complete union. Rate of success (>30% improvement in Oswestry Disability Index [ODI]) and mean change in ODI were the primary outcome measures. We compared the differences between baseline and follow-up between the Degree of Union groups.

Results: The study included 102 of 149 eligible patients. Ten patients (9.8%) were classified as having nonunion, 15 (14.7%) as having partial union, and 77 (75.5%) as having complete union. Of the 155 osteotomies, there were 122 classified as union (77%). The success rate was 74%, with no influence of SP union. The mean change in the ODI was –20.1 (95% CI –37.0, 14.2) with no influence of SP union.

Conclusions: We found no influence of SP union, classified by computed tomography, on clinical outcome 2 years after SP osteotomy in patients with lumbar spinal stenosis.

Clinical Relevance: Supplying useful information about SPO to assist surgeons in the choice of decompressive technique.

Level of Evidence: 2.

Lumbar Spine

Keywords: spinous process osteotomy, lumbar spinal stenosis, laminotomy, posterior decompression

INTRODUCTION

Degenerative lumbar spinal stenosis (LSS) is a clinical syndrome where degenerative changes in and around the spinal canal compress the neural and vascular contents of the spinal canal. This causes a pattern of pain and weakness in the lower extremities and possibly in the lower back, often accompanied by reduced walking distance. LSS is a prevalent and increasingly significant cause of low back pain and impaired walking among older adults. Among Americans older than 65 years, LSS is the most common indication for spine surgery.¹ Two large substudies of the Wakayama

Spine Study found that 9.8% of the general population had symptomatic LSS, and only 17.5% of those with severe radiological LSS had symptoms.^{2,3} The most recent Cochrane review concluded that despite many studies, the evidence in favor of decompressive surgery is sparse; however, the number of surgeries performed suggests that surgery is perceived as an effective treatment option.

Laminectomy was once considered the reference method of surgical treatment, but in recent decades, minimally invasive midline retaining procedures, such as bilateral laminotomy, unilateral laminotomy with

crossover, and to a lesser extent spinous process osteotomy (SPO), have become more popular. These minimally invasive techniques have been shown to yield similar clinical results as total laminectomies and are considered less invasive.⁴

When performing multilevel decompression, it has been debated whether to perform lumbar fusion as well. Studies have suggested no added value of fusion in the clinical outcome of multi- and single-level decompression.^{5,6}

The SPO procedure provides good visualization of the spinal canal and the lateral recesses.⁷ There are 2 main approaches to SPO for posterior decompression, one with the osteotomy being performed at the base of the spinous process (SP) and the other where the SP is split longitudinally from the dorsal surface. However, concern has been raised about whether nonunion of the osteotomy can cause pain.

Several studies have examined the union rates following SP osteotomies and their correlation to clinical outcomes. In a 2014 study involving 55 patients who underwent SP base osteotomy, a union rate of 44% was observed after 2 years. Patients with SP union in all levels showed significantly better scores on the Oswestry Disability Index (ODI) and EQ5D after 2 years; however, no significant difference was found in pain scores.⁸ In a 2019 study involving 73 patients who underwent SP splitting osteotomy, a union rate of 52% was observed between the SP and lamina after 6 to 18 months, with no impact on clinical outcome.⁹ In 2 other studies, one with 53 patients reported a union rate of 55% after 1 year, while the other with 50 patients reported a 100% union rate after 9 months.^{10,11}

The main aim of the present study was to investigate the proportion of complete SP union and examine whether radiological healing of the SP after the osteotomy (osteotomy at the base of SP) is associated with superior clinical outcome after 2 years.

MATERIALS AND METHODS

Study Population and Data Collection

The data for this retrospective cohort study was collected from the Spinal Stenosis Trial (SST), a randomized controlled trial part of the NORwegian Degenerative spondylolisthesis and spinal STENosis study (NORDSTEN-study).

The SST included 437 Norwegian patients aged 18 to 80 years who underwent surgery between April 2014 and October 2018, with claudication or radiating pain and radiological findings indicating LSS without

Table 1. Eligibility criteria for the Spinal Stenosis Trial of the NORDSTEN-study.

Inclusion Criteria
1. Clinical symptoms of lumbar spinal stenosis, neurogenic claudication, or bilateral radiating pain.
2. Not responding to at least 3 mo of nonsurgical treatment.
3. Radiological findings corresponding to the clinical symptoms: central stenosis or lateral recess stenosis.
4. Able to give informed consent and to answer the questionnaires.
5. Age >18 y.
6. Able to understand the Norwegian language, both oral and written.
Exclusion Criteria
1. Degenerative lumbar spondylolisthesis with a slip ≥ 3 mm verified on standing plain x-rays in lateral view.
2. Not willing to participate in the trial.
3. Former surgery in the level of stenosis.
4. Fracture or former fusion in the thoracolumbar region.
5. Age >80 y.
6. ASA classification of 4 or 5.
7. Cauda equina syndrome (bowel or bladder dysfunction) or fixed complete motor deficit.
8. Lumbosacral scoliosis $>20^\circ$, verified on anteroposterior view.
9. Stenosis in >3 levels.
10. Distinct symptoms in one or both of their legs due to other diseases, such as polyneuropathy, vascular claudication, or osteoarthritis.

Abbreviation: ASA, American Society of Anesthesiologists.

spondylolisthesis.¹² The patients were randomized into 3 surgical intervention groups, and the 149 patients allocated to the SPO arm were eligible for inclusion in this study. See Table 1 for inclusion and exclusion criteria.

This study was approved by the Norwegian Committees for Medical and Health Research Ethics (2011/2034).

Surgical Intervention

Access to the spinal canal was obtained by an osteotomy at the base of the SP above (and sometimes under) the affected level. The SP was then retracted to 1 side with intact supraspinous and intraspinal ligaments, giving midline access to the spinal canal (Figure 1). A laminotomy of the lower part of the superior lamina and the upper part of the inferior lamina was then performed, followed by a medial facetectomy. Both nerve roots were visualized, and the lateral recesses were decompressed. After the decompression was performed, the retractors were released, and the osteotomized SP fell back to its original position. Additional details have been published in the SST protocol.¹²

Radiological Evaluation

Computed tomography (CT) was performed 2 years postoperatively. The lumbar spine was visualized with 3-mm thick sagittal, coronal, and axial reformatted CT images. The CT examinations were evaluated by a

SPINOUS PROCESS OSTEOTOMY




Figure 1. Illustration of a spinous process osteotomy. The red zone illustrates the focus of decompression.

board-certified radiologist (H.B.) and a junior physician (S.H.). The former instructed the latter in the methodology prior to the examination. To improve inter-rater agreement, both examiners performed 20 evaluations together as a pilot. Data from the pilot study were not used in the final analysis. Afterward, all patients were examined by the 2 examiners independently. Each SP was examined individually and classified as union if there was observable bridging callus formation between the distal and proximal fragment of the SP. If no bridging callus was present, the SP was defined as nonunion. Examples are shown in Figure 2.

The patients were categorized into 3 groups, based on degree of union (DOU). If none of the patient's osteotomized SPs were classified as union, the patient was placed in the nonunion group. If some but not all SPs were classified as union, the patient was placed in the partial union group, and if all SPs were classified as union, the patient was placed in the complete union group. Inconsistencies between the 2 examiners were reviewed independently by an orthopedic surgeon (E.H.) to obtain consensus.




Figure 2. Transverse and sagittal computed tomography images of the lumbar spine. A nonunion is shown in the top 2 images and a union in the bottom 2 images.

Primary Outcome Measures

Mean change in ODI score from baseline to 2-year follow-up was chosen as the primary outcome. The ODI is the most used outcome measure in spinal surgery; we used the Norwegian version 2.0.^{12–14} The ODI is expressed as an index ranging from no disability (0) to the worst imaginable disability (100).

The proportion of patients classified as achieving successful outcomes was chosen as the second primary outcome. Success was defined as $\geq 30\%$ improvement in ODI score from baseline to follow-up. This criterion has been used in the other publications from the NORDSTEN-study and is considered a relevant outcome measure.¹⁵

Secondary Outcome Measures

Mean change in patient-reported outcome measures (PROMs) comprised the secondary outcome measures. Mean change in symptom severity and functional outcome was measured by change scores on the Zürich claudication questionnaire, a standardized spinal stenosis-specific self-administered questionnaire consisting of a symptom severity part (ranging from 1–5), a daily function part (ranging from 1–4), and a surgery satisfaction part.¹⁶ Mean change in low back and leg pain was quantified using a numeric rating scale (NRS,

Table 2. Demographics and baseline pain and function scores in the 3 degree of union groups.

Variable	Total (n = 102)	Nonunion (n = 10)	Partial Union (n = 15)	Complete Union (n = 77)	P
Age, y, mean (SD)	66.2 (8.4) (n = 102)	66.5 (7.2) (n = 10)	65.1 (11.6) (n = 15)	66.3 (8.0) (n = 77)	0.87
Smokers, %	18.6% (n = 19)	20% (n = 2)	40% (n = 6)	14.3% (n = 11)	0.51
Men, %	64.7% (n = 66)	40% (n = 4)	80% (n = 12)	65% (n = 50)	0.12
BMI, mean (SD)	27.4 (4.3) (n = 97)	27.4 (3.8) (n = 10)	28.1 (3.6) (n = 14)	27.3 (4.5) (n = 73)	0.82
ODI, mean (SD)	35.7 (14.5) (n = 101)	41.0 (17.8) (n = 10)	38.6 (17.5) (n = 15)	34.4 (13.4) (n = 76)	0.28
ZCQ-S, mean (SD)	3.3 (0.5) (n = 98)	3.5 (0.3) (n = 10)	3.3 (0.5) (n = 15)	3.2 (0.5) (n = 73)	0.20
ZCQ-F, mean (SD)	2.5 (0.54) (n = 99)	2.6 (0.4) (n = 10)	2.5 (0.5) (n = 15)	2.4 (0.6) (n = 74)	0.51
LP NRS, mean (SD)	6.3 (1.9) (n = 97)	6.6 (1.8) (n = 10)	6.4 (2.5) (n = 14)	6.3 (1.9) (n = 73)	0.86
LBP NRS, mean (SD)	5.8 (2.2) (n = 98)	7.1 (1.9) (n = 10)	4.9 (2.2) (n = 14)	5.8 (2.2) (n = 74)	0.04
EQ5D, mean (SD)	0.41 (0.30) (n = 100)	0.29 (0.30) (n = 10)	0.43 (0.29) (n = 15)	0.43 (0.30) (n = 75)	0.37

Abbreviations: EQ5D, EuroQol health related quality of life index; LBP, low back pain; LP, leg pain; NRS, numeric rating scale; ODI, Oswestry Disability Index; ZCQ-S/F, Zürich Claudication Questionnaire Symptoms/Function.

ranging from 1–10). Mean change in health-related quality of life was measured by EQ-5D, a prominent health-related quality of life tool that evaluates health status across 5 dimensions (mobility, self-care, usual activities, pain/discomfort, and anxiety/depression) and a visual analog scale for self-rated health assessment (indexed scores ranging from –0.59 to 1).¹⁷ Radiological outcomes included the number of osteotomies defined as union and the number of patients in each DOU group.

Statistical Analysis

All analyses were performed using IBM SPSS statistics version 28.0.0.0 (190). The baseline data and the change from baseline to follow-up in PROMs were examined for DOU-group differences. For the continuous variables (age, body mass index, and PROMs), analysis of variance with Tukey's honestly significant difference for post hoc testing to control for making type-I error was used. For the categorical variables (sex,

smoking, and success), the χ^2 test was used. In the cases with an expected count of less than 5, the Fisher's exact test was used. Due to the skewed DOU-group sizes, another analysis was performed on the change from baseline PROM data, with the nonunion and partial union treated as 1 group and compared with the complete union group using an independent 2-sample *t* test. A weighted Cohen's kappa analysis was performed to assess interrater agreement between the 2 examiners.

RESULTS

Of 149 patients eligible for inclusion, 102 were included in this study (Figure 3). Ten (9.8%) patients were classified as nonunion, 15 (14.7%) as a partial union, and 77 (75.5%) as a complete union. The number of patients in each DOU group who answered all follow-up PROMs varied (detailed with secondary outcomes later in this section). The total number of osteotomies, counting every cut SP on every patient, was 155, of which 122 (77.7%) were classified as union at 2-year follow-up. Two examiners agreed on all but 2 cases, resulting in an observed agreement rate of 98% and a weighted Cohen's kappa of 0.95 (95% CI 0.87–1.03).

Baseline Data

The baseline characteristics of the patients are shown in Table 2. There were no significant differences in the measured demographic data between the 3 DOU groups. The only baseline PROM with significant DOU-group differences was low back pain measured with NRS, with the nonunion group mean being 2.4 points higher than that of the partial union group ($P = 0.04$).

Primary and Secondary Outcomes

The mean change in ODI was –20.1 (95% CI –37.0, –14.2) from baseline to follow-up in the total cohort.


**Figure 3.** Flowchart of the patients in the current study.

Table 3. Primary outcome of change from baseline to follow-up in the 3 degree of union groups

Outcome Measure	Total (n = 101)	Nonunion (n = 10)	Partial Union (n = 15)	Complete Union (n = 76)	P
ODI, mean (SD)	-20.1 (17.0) (n = 100)	-25.6 (15.9) (n = 10)	-17.3 (20.7) (n = 14)	-19.9 (16.5) (n = 76)	0.49
Success	74% (n = 100)	90% (n = 10)	64.3% (n = 14)	73.7% (n = 76)	0.39

Abbreviation: ODI, Oswestry Disability Index.

Note: Success $\geq 30\%$ improvement in ODI.

The mean change from baseline to follow-up was -25.6 (95% CI $-14.2, -37.0$) in the nonunion group and -19.9 (95% CI $-16.7, -23.7$) in the complete union group. The differences between the DOU groups were not statistically significant ($P = 0.49$; Table 3).

There were no statistically significant differences in the mean change from baseline to follow-up between the union groups for any secondary outcome measures (Table 4). The number of patients in each DOU group who answered all follow-up PROMs varied, as noted throughout Table 4.

When t tests were performed with nonunion and partial union patients combined into 1 group (mean ODI change -20.8) vs the complete union group (mean ODI change -19.9), no significant differences were found ($P = 0.83$; Table 5).

DISCUSSION

The main finding of this study is that there was no influence of SP union on 2-year patient-reported outcomes. There was a high proportion of patients classified as complete union (75.5%), a high success rate (74%), and a high number of osteotomies classified as union (77.7%).

The mean improvement in ODI is in line with results after other decompressive techniques and above current estimates of minimal clinically important difference.^{7,8,18–20} The clinical results after different minimal invasive decompression techniques are similar, as shown in a recent randomized controlled trial.²¹

The previously mentioned studies examining union between the SP and lamina found union rates between 44% and 100% after 6 to 24 months in groups of 50 to 73 patients. Hermansen et al⁸ found patients with union

in all levels to have better clinical outcomes after 2 years, while Wi et al found no relationship between union and clinical outcome after 6 to 18 months. We found a high rate of union (77%) among our 102 patients, with no effect of union on clinical outcome. Two of the benefits of our study compared with the aforementioned studies are the larger sample size and the use of 2 independent examiners using CT imaging to evaluate union.

A limitation of this study is the unequal number of patients in the DOU groups, with only 10 patients classified as nonunion according to our classification. Another limitation is that the lack of an existing standardized method for evaluation of SP union makes our qualitative assessments harder to precisely reproduce. Our follow-up interval of 2 years may also be too short to determine the long-term effects of SP union. In a 2015 study investigating the effects of SP union following SP splitting osteotomy during open-door laminoplasty, the authors reported no significant impact of union on clinical outcome after 2 to 4 years. However, they observed that nonunion was associated with worse clinical outcomes after 10 to 12 years.²² Further studies with extended follow-up intervals of 5 and 10 years are required to observe the long-term effects of SP union on clinical outcomes after the SP base osteotomy procedure. Additional studies should also include magnetic resonance imaging scans to visualize potential noncalcified fibrotic tissue formation between the proximal and distal SP fragments. Another limitation is that we, in our analysis of the CT images, did not assess other readily available parameters relevant for LSS in general, such as changes in slip progression and facet joint degeneration in the index level and adjacent levels.

Table 4. Secondary outcome measures change from baseline to follow-up, mean (SD).

Outcome Measure	Total (n = 101)	Nonunion (n = 10)	Partial Union (n = 15)	Complete Union (n = 76)	P
ZCQ-S	-1.1 (0.9) (n = 98)	-1.4 (0.7) (n = 10)	-1.0 (0.8) (n = 15)	-1.0 (0.9) (n = 73)	0.45
ZCQ-F	-0.9 (0.8) (n = 98)	-0.9 (0.7) (n = 9)	-0.7 (0.7) (n = 15)	-0.9 (0.7) (n = 74)	0.55
LP NRS	-3.6 (3.1) (n = 95)	-3.6 (3.1) (n = 10)	-3.2 (3.3) (n = 13)	-3.6 (3.0) (n = 72)	0.93
LBP NRS	-2.8 (2.5) (n = 97)	-3.6 (2.6) (n = 10)	-1.7 (2.3) (n = 14)	-2.9 (2.5) (n = 73)	0.15
EQ5D	0.34 (0.33) (n = 93)	0.4 (0.3) (n = 10)	0.23 (0.38) (n = 14)	0.34 (0.32) (n = 69)	0.29

Abbreviations: EQ5D, EuroQol health related quality of life index; LBP, low back pain; LP, leg pain; NRS, numeric rating scale; ZCQ-S/F, Zürich Claudication Questionnaire Symptoms/Function.

Table 5. Primary and secondary outcome measures: change from baseline to follow-up, comparing the complete union group and not complete union group made from merging the nonunion group and partial union group.

Outcome Measure	Not Complete Union (n = 24)	Complete Union (n = 76)	P
ODI	-20.8 (19.0) (n = 24)	-19.9 (16.5) (n = 76)	0.83
Success, % ^a	75% (n = 18)	74% (n = 56)	0.90
ZCQ-S	-1.2 (0.8) (n = 25)	-1.0 (0.9) (n = 73)	0.55
ZCQ-F	-0.8 (0.7) (n = 24)	-0.9 (0.7) (n = 74)	0.46
LP NRS	-3.4 (3.1) (n = 23)	-3.6 (3.0) (n = 72)	0.79
LBP NRS	-2.5 (2.6) (n = 24)	-2.9 (2.5) (n = 73)	0.51
EQ5D	0.32 (0.35) (n = 24)	0.34 (0.32) (n = 69)	0.79

Abbreviations: EQ5D, EuroQol health related quality of life index; LBP, Lower Back Pain; LP, leg pain; ODI, Oswestry Disability Index; ZCQ-S/F, Zürich Claudication Questionnaire Symptoms/Function.

Note: Data presented as mean (SD) unless otherwise indicated.

^aSuccess ≥30% improvement in ODI.

Further research is needed on the long term clinical importance of SP union and if one should try to facilitate SP union, for example, through osteosutures. Our findings suggest that nonunion does not worsen clinical outcomes and that trying to facilitate union thus seems unnecessary. SPO is a viable primary decompressive approach for LSS or as a rescue technique when performing unilateral laminotomy with crossover. The surgeon can thereby choose a technique according to his or her preference and experience. SPO is, by some surgeons, considered to be less technically demanding and offers good visualization, especially when decompressing the lateral recess.

CONCLUSION

Most of the SPOs (77%) were classified as union. The success rate was high in the total sample (74%). We found no influence of SP union, classified by CT, on clinical outcome 2 years after SPO in patients with LSS.

REFERENCES

- Deasy J. Acquired lumbar spinal stenosis. *JAAPA*. 2015;28(4):19–23. doi:10.1097/01.JAA.0000462052.47882.f
- Ishimoto Y, Yoshimura N, Muraki S, et al. Prevalence of symptomatic lumbar spinal stenosis and its association with physical performance in a population-based cohort in Japan: the wakayama spine study. *Osteoarthritis Cartilage*. 2012;20(10):1103–1108. doi:10.1016/j.joca.2012.06.018
- Ishimoto Y, Yoshimura N, Muraki S, et al. Associations between radiographic lumbar spinal stenosis and clinical symptoms in the general population: the wakayama spine study. *Osteoarthritis Cartilage*. 2013;21(6):783–788. doi:10.1016/j.joca.2013.02.656
- Nerland US, Jakola AS, Solheim O, et al. Minimally invasive decompression versus open laminectomy for central stenosis of the lumbar spine: pragmatic comparative effectiveness study. *BMJ*. 2015;350(apr01 1):h1603. doi:10.1136/bmj.h1603
- Nolte MT, Louie PK, Basques BA, et al. Patients undergoing 3-level-or-greater decompression-only surgery for lumbar spinal stenosis have similar outcomes to those undergoing single-level surgery at 2 years. *Int J Spine Surg*. 2021;15(5):945–952. doi:10.14444/8124
- Försth P, Ólafsson G, Carlsson T, et al. A randomized, controlled trial of fusion surgery for lumbar spinal stenosis. *N Engl J Med*. 2016;374(15):1413–1423. doi:10.1056/NEJMoa1513721
- Hermansen E, Romild UK, Austevoll IM, et al. Does surgical technique influence clinical outcome after lumbar spinal stenosis decompression? a comparative effectiveness study from the norwegian registry for spine surgery. *Eur Spine J*. 2017;26(2):420–427. doi:10.1007/s00586-016-4643-9
- Hermansen E, Moen G, Fenstad AM, Birketvedt R, Indrekvam K. Spinous process osteotomy to facilitate the access to the spinal canal when decompressing the spinal canal in patients with lumbar spinal stenosis. *Asian Spine J*. 2014;8(2):138–144. doi:10.4184/asj.2014.8.2.138
- Wi SM, Lee HJ, Chang SY, et al. Restoration of the spinous process following muscle-preserving posterior lumbar decompression via sagittal splitting of the spinous process. *Clin Orthop Surg*. 2019;11(1):95–102. doi:10.4055/cios.2019.11.1.95
- Lee GW, Ahn M-W. Comparative study of two spinous process (SP) osteotomy techniques for posterior decompression surgery in lumbar spinal stenosis: sp base versus splitting osteotomy. *Eur Spine J*. 2018;27(7):1644–1652. doi:10.1007/s00586-018-5526-z
- Takaso M, Nakazawa T, Imura T, et al. Less invasive and less technically demanding decompressive procedure for lumbar spinal stenosis—appropriate for general orthopaedic surgeons. *Int Orthop*. 2011;35(1):67–73. doi:10.1007/s00264-010-0986-8
- Hermansen E, Austevoll IM, Romild UK, et al. Study-protocol for a randomized controlled trial comparing clinical and radiological results after three different posterior decompression techniques for lumbar spinal stenosis: the spinal stenosis trial (SST). *BMC Musculoskelet Disord*. 2017;18(1):121. doi:10.1186/s12891-017-1491-7
- [ICHOM]. The ICHOM Standard Set for Low Back Pain. 2020. <https://www.ichom.org/portfolio/low-back-pain>.
- Fairbank JCT, Pynsent PB. The Oswestry disability index. *Spine*. 2000;25(22):2940–2952. doi:10.1097/00007632-200011150-00017
- Austevoll IM, Gjestad R, Grotle M, et al. Follow-up score, change score or percentage change score for determining clinical important outcome following surgery? An observational study from the norwegian registry for spine surgery evaluating patient reported outcome measures in lumbar spinal stenosis and lumbar degenerative spondylolisthesis. *BMC Musculoskelet Disord*. 2019;20(1):31. doi:10.1186/s12891-018-2386-y
- Stucki G, Daltroy L, Liang MH, Lipson SJ, Fossel AH, Katz JN. Measurement properties of a self-administered outcome measure in lumbar spinal stenosis. *Spine*. 1996;21(7):796–803. doi:10.1097/00007632-199604010-00004
- Rabin R, de Charro F. EQ-SD: a measure of health status from the euroqol group. *Ann Med*. 2001;33(5):337–343. doi:10.3109/07853890109002087
- Zaina F, Tomkins-Lane C, Carragee E, Negrini S. Surgical versus non-surgical treatment for lumbar spinal stenosis. *Cochrane Database Syst Rev*. 2016;2016(1):CD010264. doi:10.1002/14651858.CD010264.pub2

19. Hung M, Saltzman CL, Kendall R, et al. What are the MCIDs for PROMIS, NDI, and ODI instruments among patients with spinal conditions. *Clin Orthop Relat Res*. 2018;476(10):2027–2036. doi:10.1097/CORR.0000000000000419
20. Maughan EF, Lewis JS. Outcome measures in chronic low back pain. *Eur Spine J*. 2010;19(9):1484–1494. doi:10.1007/s00586-010-1353-6
21. Hermansen E, Austevoll IM, Hellum C, et al. Comparison of 3 different minimally invasive surgical techniques for lumbar spinal stenosis. *JAMA Netw Open*. 2022;5(3):e224291. doi:10.1001/jamanetworkopen.2022.4291
22. Kakiuchi M, Fukushima W. Impact of Spinous process integrity on ten to twelve-year outcomes after posterior decompression for lumbar spinal stenosis: study of open-door laminoplasty using a spinous process-splitting approach. *J Bone Joint Surg Am*. 2015;97(20):1667–1677. doi:10.2106/JBJS.N.01370

Funding: This trial received funding from Central Norway Regional Health Authority (RHA) and Western Norway Regional Health Authority (RHA).

Disclosures: The authors report no disclosures or conflicts of interest.

Corresponding Author: Sondre Hagerup; sohare97@gmail.com

Published 27 February 2024

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2024 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.