

Increasing Nonconcurrent Overlapping Surgery Is Not Associated With Outcome Changes in Lumbar Fusion

Ali S. Farooqi, Austin J. Borja, Donald K.E.D. Detchou, Gregory Glauser, Krista Strouz, Scott D. McClintock and Neil R. Malhotra

Int J Spine Surg published online 25 May 2022
<https://www.ijssurgery.com/content/early/2022/06/14/8305>

This information is current as of May 17, 2025.

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://ijssurgery.com/alerts>

Increasing Nonconcurrent Overlapping Surgery Is Not Associated With Outcome Changes in Lumbar Fusion

ALI S. FAROOQI, BA¹; AUSTIN J. BORJA, BA¹; DONALD K.E.D. DETCHOU, BA¹; GREGORY GLAUSER, MD, MBA, MA¹; KRISTA STROUZ, BS^{2,3}; SCOTT D. MCCLINTOCK, PhD³; AND NEIL R. MALHOTRA, MD¹

¹Department of Neurosurgery, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, USA; ²McKenna EpiLog Fellowship in Population Health at the University of Pennsylvania, Philadelphia, USA; ³The West Chester Statistical Institute and Department of Mathematics, West Chester University, West Chester, USA

ABSTRACT

Background: There remains a paucity of literature on the impact of overlap on neurosurgical patient outcomes. The purpose of the present study was to correlate increasing duration of surgical overlap with short-term patient outcomes following lumbar fusion.

Methods: The present study retrospectively analyzed 1302 adult patients undergoing overlapping, single-level, posterior-only lumbar fusion within a single, multicenter, academic health system. Recorded outcomes included 30-day emergency department visits, readmission, reoperation, mortality, overall morbidity, and overall morbidity/surgical complications. The amount of overlap was calculated as a percentage of total overlap time. Comparison was made between patients with the most (top 10%) and least (bottom 40%) amount of overlap. Patients were then exact matched on key demographic factors but not by the attending surgeons. Subsequently, patients were exact matched by both demographic data and the attending surgeons. Univariate analysis was first carried out prior to matching and then on both the demographic-matched and surgeon-matched cohorts. Significance for all analyses was set at a P value of <0.05 .

Results: Within the whole population, increasing duration of overlap was not correlated with any short-term outcome ($P = 0.41$ – 0.91). After exact matching, patients with the most and least durations of overlap did not have significant differences with respect to any short-term outcomes ($P = 0.34$ – 1.00).

Conclusion: Increased amount of overlap is not associated with adverse short-term outcomes for single-level, posterior-only lumbar fusions.

Clinical Relevance: The present results suggest that increasing the duration of overlap during lumbar fusion surgery does not lead to inferior outcomes.

Level of Evidence: 3.

Lumbar Spine

Keywords: concurrent surgery, lumbar spine, overlapping surgery, patient safety, spinal fusion

INTRODUCTION

Surgeons often perform cases that can overlap in operating time, a practice known as overlapping surgery. Overlapping surgery came into public attention following a 2015 *Boston Globe* investigation, leading to increased scrutiny and demands for research evaluating the safety of overlapping surgery.¹ In response, the American College of Surgeons updated their guidelines regarding overlapping surgery.² They stated that concurrent surgery—in which a surgeon simultaneously performs the critical elements of multiple cases—is always inappropriate. In contrast, overlapping surgery—in which a surgeon alternates operating rooms after completing the critical elements of a case, allowing another practitioner to complete the noncritical elements of the case—is acceptable.

Given the recent public concern about patient safety, additional research is warranted to examine the impact of overlap on surgical outcomes.³ Previous literature across multiple surgical services has not demonstrated an increase in morbidity, mortality, or complication rate, despite occasionally finding an increased duration of surgery.^{4–7} There are similar results for overlap within neurosurgery, with previous studies reporting noninferior outcomes following all neurosurgical procedures,^{8,9} elective neurosurgical procedures,¹⁰ pituitary adenoma resection,¹¹ microvascular decompression,¹² and spine surgery.¹³ Although these reports suggest that surgical overlap is safe, few studies have evaluated whether an increasing amount of overlap during surgery confers additional risk to patients. Previous studies in orthopedic surgery

found no association with complication rate and increasing duration of overlap,^{5,14} but there are a paucity of studies that have evaluated the impact of varying degrees of neurosurgical overlap on short-term outcomes.

Here, we assess increasing duration of surgical overlap on 30-day outcomes in the setting of posterior-only, single-level lumbar fusion. Lumbar fusion is a common neurosurgical procedure that can have complication rates near 13% to 16% in the short-term postoperative period, making it particularly relevant to determine the risk factors for adverse outcomes.^{15,16} This study utilized coarsened exact matching (CEM) to compare outcomes among patients matched on an increasing number

of variables, including demographics and attending surgeon.

METHODS

Sample Selection

This Institutional Review Board–approved study was considered to pose a minimal risk to patients and was granted a waiver of informed consent. A total of 3799 consecutive patients undergoing single-level, posterior-only lumbar fusion at a single, multihospital, academic health system over a 6-year period (from 07 June 2013 to 29 April 2019) were enrolled retrospectively (Figure 1). Exclusion criteria included patients with missing health information, significantly elevated body




Figure 1. Flowchart of lumbar fusion cases included within the present study.

mass index (BMI), nonelective/nonroutine operations, nongeneral anesthesia, and unclean wound closure. Key patient characteristic data were acquired via EpiLog—a nonproprietary data acquisition system created by the senior author (N.R.M.) and built within the existing electronic health record for quality improvement initiatives.¹⁷

The whole population for analysis included only patients with overlap during operative time ($n = 1302$). The amount of surgical overlap for each patient was defined as a percentage of total operative duration. Univariate logistic regression was employed to correlate increasing amount of overlap with short-term patient outcomes, including readmission, reoperation, emergency department visits, mortality, morbidity, and surgical complications. Furthermore, exact-matched patients with the least amount (bottom 40%) and most amount (top 10%) of overlap time were compared.

Data Collection and Matching Process

CEM was performed to account for potentially confounding variables when comparing patients with the least and most amounts of overlap. CEM employs 1:1 matching via the original covariates.¹⁸ This contrasts propensity score matching, which converts the covariates into a composite value for matching, otherwise ignoring the original values. Under the present CEM protocol, an exact match was defined as a match on every prespecified covariate.

Patients with the least and most amounts of overlap were exact matched on 27 unique patient characteristics including gender, race, insurance status, median household income, level of education, tobacco use history, BMI, Charlson Comorbidity Index (CCI) score and its underlying components, American Society of Anesthesiologists (ASA) grade, and surgical history. Binary matching was employed for race (white or nonwhite) and median household income (above or below the median value for the dataset). Ternary matching was performed for BMI, CCI, and ASA grade, with patients binned into either a low-, medium-, or high-value cohort. Exact matching was performed for the remaining matching covariates. Unmatched patients were removed from the matched analysis. After matching on demographic data, matches were limited to those with the same operating surgeon.

Statistical Methods

Demographic, clinical, and outcome data were extracted from the electronic health record via

EpiLog and placed into defined spreadsheets. Univariate logistic regression was performed using SAS version 9.4 (SAS Institute Inc., Cary, North Carolina) to analyze the duration of overlap as a continuous variable (ie, 1% increase in amount of overlap). For CEM, binning of the matching variables and removal of missing values were performed using SAS version 9.4, whereas matching was completed with the MatchIt programming package in R Statistics (R Core Team, Vienna, Austria, 2017). McNemar test was used to compare means of outcomes between all exact-matched groups. Statistical significance was defined as a P value < 0.05 for all analyses.

RESULTS

Patient Characteristics: Before Exact Matching

Prior to exact matching, patients with the most surgical overlap demonstrated greater CCI baseline score ($P = 0.031$), but a comparable distribution of comorbidities, and a similar ASA grade (Figure 2A). No significant difference in surgical history was observed. Patients with the most amount of overlap also had significantly shorter total length of surgery ($P = 0.017$), but a comparable length of multilayered wound closure. Patients with the most amount of overlap also had significantly shorter length of stay ($P = 0.006$) and duration of follow-up ($P = 0.018$), but no difference in the number of non-neurosurgical interventions within 30 days of the index operation (Table 1). Patients with increasing overlap had significantly lower operating room costs ($P < 0.001$) and differed by the insurance provider ($P = 0.017$).

Patient Characteristics: Demographic-Matched Cohort

Patients were exact matched on demographic variables but not limited to the same operating surgeon ($n = 158$). Matched cohorts demonstrated similar CCI score and ASA grade and a similar distribution of comorbidities, but patients with the most surgical overlap had significantly lower rates of nonmetastatic cancer ($P = 0.028$) (Figure 2B). No significant difference in surgical history was observed. Patients with the most amount of overlap had a shorter total length of surgery ($P = 0.018$), but not the length of multilayered wound closure. No difference was observed in duration of follow-up, but patients with the most overlap were observed to have significantly shorter lengths of stay ($P = 0.001$) (Table 2). There was no difference in the insurance provider, but patients with




Figure 2. Comparison of baseline health characteristics, distribution of comorbidities, and operative time among patients with the most and least amount of surgical overlap. *Significance at P value <0.05 . ASA, American Society of Anesthesiologists grade; CCI, Charlson Comorbidity Index score; CHF, congestive heart failure; CKD, chronic kidney disease; COPD, chronic obstructive pulmonary disease; CVA, cerebrovascular accident; DM, diabetes mellitus; HIV, human immunodeficiency virus; LD, liver disease; MI, myocardial infarction; PUD, peptic ulcer disease; PVD, peripheral vascular disease; RA, rheumatoid arthritis.

more overlap had a lower operating room cost ($P < 0.001$).

Patient Characteristics: Demographic-Matched Cohort Limited by Surgeon

After exact matching on demographics, matches were limited to those with the same operating surgeon ($n = 68$). No significant differences were observed in CCI score, distribution of comorbidities, ASA grade, and surgical history. Furthermore, there was no difference in length of surgery or length of multilayered wound closure (Figure 2C). Finally, no significant differences were observed in the duration of follow-up, length of stay, insurance type, or cost of surgery (Table 3).

Patient Outcomes: Whole Population

Within the whole population, increasing duration of overlap was not correlated with any morbidity or mortality outcome, including 30-day readmission, reoperation, emergency department visits, or mortality ($P = 0.41-0.91$). When evaluating overall morbidity or overall morbidity including surgical complications, there was also no significant correlation ($P = 0.70-0.78$) (Table 4).

Outcomes in Demographic-Matched Cohort

After exact matching patients with the most overlap to those with the least overlap after on demographics, but not by the surgeon, there was no difference in any

Table 1. Characteristics of patients whose surgeries had the least amount of overlap (bottom 40% of patients with overlapping surgeries) and the most amount of overlap (top 10% of patients with overlapping surgeries).

Variable	Least Overlap (N = 520)	Most Overlap (N = 131)	P Value
Age (y), mean (range)	60.30 (24–88)	63.90 (26–87)	0.001
Gender, n (%)			0.96
Male	225 (43.3%)	57 (43.5%)	
Female	295 (56.7%)	74 (56.5%)	
Race, n (%) Asian	6 (1.2%)	1 (0.8%)	0.55
Black	55 (10.6%)	12 (9.2%)	
White	441 (84.8%)	117 (89.3%)	
Hispanic	4 (0.8%)	0 (0%)	
Unknown	10 (1.9%)	0 (0%)	
Other	2 (0.4%)	1 (0.8%)	
East Indian	2 (0.4%)	0 (0%)	
Insurance, n (%)			0.017
Commercial	20 (3.8%)	2 (1.5%)	
Medicare	213 (41.0%)	73 (55.7%)	
Medicaid	26 (5.0%)	4 (3.1%)	
Managed care	192 (36.9%)	33 (25.2%)	
Blue cross	69 (13.3%)	19 (14.5%)	
Tobacco use within past 12 mo, n (%)			0.08
Yes	81 (15.6%)	11 (8.4%)	
No	437 (84.0%)	120 (91.6%)	
Unknown	2 (0.4%)	0 (0%)	
Fasting blood glucose (mg/dL), mean (range)	133.95 (60–303)	127.84 (79–209)	0.32
Surgeries before index operation (count), mean (range)			
Ever	0.58 (0–12)	0.61 (0–14)	0.39
90 d Prior	0.07 (0–5)	0.02 (0–1)	0.13
Direct cost of operative supplies and implants (\$), mean (range)	10,254 (116–50,894)	6696 (342–27,592)	<0.001
Length of hospital stay (h), mean (range)	104.64 (24–1070)	96.59 (26–563)	0.006
Length of follow-up (mo), mean (range)	43.62 (0.5–83.11)	38.24 (0.7–80.74)	0.018
Non-neurosurgical operations 30 d after index operation (count), mean (range)	0.07 (0–3)	0.09 (0–2)	0.31

Note: Statistically significant ($P < 0.05$) values are bolded.

Table 2. Characteristics of the subset of patients with the least amount of overlap (bottom 40% of patients with overlap) and the most amount of overlap (top 10% of patients with overlap), subsequently matched on demographic variables but not the operating surgeon.

Variable	Least Overlap (N = 79)	Most Overlap (N = 79)	P Value
Age (y), mean (range)	61.96 (24–79)	63.20 (26–79)	0.50
Gender, n (%)			>0.99
Male	32 (40.5%)	32 (40.5%)	
Female	47 (59.5%)	47 (59.5%)	
Race, n (%)			>0.99
Black	2 (2.5%)	2 (2.5%)	
White	77 (97.5%)	77 (97.5%)	
Insurance, n (%)			0.53
Commercial	3 (3.8%)	1 (1.3%)	
Medicare	39 (49.4%)	42 (53.2%)	
Medicaid	4 (5.1%)	1 (1.3%)	
Managed care	22 (27.8%)	21 (26.6%)	
Blue cross	11 (13.9%)	14 (17.7%)	
Tobacco use within past 12 mo, n (%)			>0.99
Yes	1 (1.3%)	1 (1.3%)	
No	78 (98.7%)	78 (98.7%)	
Fasting blood glucose (mg/dL), mean (range)	136.26 (92–222)	128.48 (79–207)	0.35
Surgeries before index operation (count), mean (range)			
Ever	0.19 (0–4)	0.14 (0–4)	0.98
90 d Prior	0.03 (0–2)	0.01 (0–1)	>0.99
Direct cost of operative supplies and implants (\$), mean (range)	10,606 (960–42,850)	6029 (342–20,689)	<0.001
Length of hospital stay (h), mean (range)	106.43 (48–400)	84.92 (26–367)	0.001
Length of follow-up (mo), mean (range)	43.36 (3.75–81.27)	40.16 (0.66–80.75)	0.41
Non-neurosurgical operations 30 d after index operation (count), mean (range)	0.04 (0–1)	0.08 (0–1)	0.31

Note: Statistically significant ($P < 0.05$) values are bolded.

Table 3. Characteristics of the subset of patients with the least amount of overlap (bottom 40% of patients with overlap) and the most amount of overlap (top 10% of patients with overlap), subsequently matched on both demographic variables and limited to the same operating surgeon.

Variable	Least Overlap (N = 34)	Most Overlap (N = 34)	P Value
Age (y), mean (range)	56.74 (24–76)	58.26 (26–76)	0.64
Gender, n (%)			>0.99
Male	19 (55.9%)	19 (55.9%)	
Female	15 (44.1%)	15 (44.1%)	
Race, n (%)			>0.99
Black	1 (2.9%)	1 (2.9%)	
White	33 (97.1%)	33 (97.1%)	
Insurance, n (%)			0.20
Commercial	4 (11.8%)	0 (0%)	
Medicare	12 (35.3%)	13 (38.2%)	
Medicaid	2 (5.9%)	1 (2.9%)	
Managed care	11 (32.4%)	10 (29.4%)	
Blue cross	5 (14.7%)	10 (29.4%)	
Tobacco use within past 12 mo, n (%)			>0.99
Yes	1 (2.9%)	1 (2.9%)	
No	33 (97.1%)	33 (97.1%)	
Fasting blood glucose (mg/dL), mean (range)	122.48 (92–174)	128.73 (79–207)	0.46
Surgeries before index operation (count), mean (range)			
Ever	0.44 (0–10)	0.24 (0–4)	0.98
90 d Prior	0.06 (0–2)	0.03 (0–1)	>0.99
Direct cost of operative supplies and implants (\$), mean (range)	7120 (116–26,035)	6400 (342–17,459)	0.75
Length of hospital stay (h), mean (range)	86.73 (25–242)	69.65 (31–151)	0.10
Length of follow-up (mo), mean (range)	34.84 (3.0–78.44)	41.42 (1.3–77.69)	0.21
Non-neurosurgical operations 30 d after index operation (count), mean (range)	0.12 (0–1)	0.09 (0–1)	0.70

30-day adverse outcome, overall morbidity, or overall morbidity/surgical complications ($P = 0.38$ – 1.00). There was no 30-day mortality recorded in this exact-matched cohort (Figure 3, Table 4).

Outcomes in Demographic-Matched Cohort Limited by Surgeon

After exact matching cohorts on both demographics and limiting to the same operating surgeon, there was no significant difference in any 30-day adverse outcome, overall morbidity, or overall morbidity/surgical complications ($P = 0.34$ – 0.69). No 30-day mortality events were recorded in this exact-matched cohort (Figure 3, Table 4).

DISCUSSION

Increasing duration of overlap, in the setting studied, does not lead to adverse short-term outcomes following single-level, posterior-only lumbar fusion. Duration of overlap was not correlated with any short-term patient outcome in the whole population. When comparing demographic exact-matched cohorts with the least and most amounts of overlap, there were no significant differences in any short-term morbidity outcome. Comparison of exact-matched cohorts yielded the same results, when matching was further limited by the surgeon performing the operation.

Table 4. Short-term outcome data analyzing the impact of increasing duration of overlap as a continuous variable. Exact-matched patients with the most amount of overlap (top 10%) and least amount of overlap (bottom 40%) were also compared.

30-d Outcomes	Increasing Duration of Overlap	After Exact Match Not Limited by Surgeon	After Exact Match Limited by Surgeon
Readmission	$P = 0.91$ 1.00 (0.99–1.01)	$P > 0.99$ 1.00 (0.17–5.82)	$P = 0.69$ 0.67 (0.08–4.48)
Reoperation	$P = 0.91$ 1.00 (0.99–1.01)	$P = 0.38$ 3.00 (0.32–78.99)	$P = 0.69$ 1.50 (0.22–12.61)
Emergency department visits	$P = 0.41$ 1.00 (0.99–1.01)	$P = 0.63$ 0.50 (0.02–6.57)	$P = 0.69$ 0.67 (0.08–4.48)
Mortality	$P = 0.90$ 1.00 (0.94–1.05)	N/A	N/A
Morbidity	$P = 0.70$ 1.00 (0.99–1.01)	$P = 0.51$ 0.60 (0.12–2.60)	$P = 0.51$ 0.60 (0.12–2.60)
Morbidity/surgical complication	$P = 0.78$ 1.00 (0.99–1.01)	$P = 0.39$ 0.57 (0.15–1.97)	$P = 0.34$ 0.50 (0.10–2.01)

Note: Data are presented as OR (95% CI).




Figure 3. Distribution of short-term outcomes following single-level, posterior lumbar fusion. No differences were observed in any morbidity or mortality outcomes between exact matched cohorts. ED, emergency department.

There is a paucity of research on the impact of increasing overlap duration on neurosurgical outcomes. Previous studies in orthopedic surgery have not demonstrated any association between duration of overlap and adverse 30- or 90-day patient outcomes.^{5,14} Our results supplement these findings by demonstrating that an increasing amount of overlap time during lumbar fusion does not lead to an increased risk of 30-day complications when controlling for patient demographic variables and surgeon variability. Future studies should examine increasing overlap time, longer-term outcomes, and outcomes following other procedures to thoroughly assess the safety of increasing overlap time in neurosurgery.

Previous studies have found that individuals selected for overlapping surgery are typically healthier at baseline and have lower CCI score, fewer comorbidities, and lower ASA grade.^{10,19} However, this study found that patients with the most overlap had similar CCI score, similar distribution of comorbidities, a higher ASA grade as compared with patients with the least overlap. This difference suggests that, although surgeons

typically select healthier patients as candidates for overlapping surgery, it is not always the case.

Previous studies within neurosurgery found that patients with overlap typically had a longer length of surgery as compared with patients without overlap, but no additional risk of adverse outcomes.⁸⁻¹³ Here, among the whole population and the cohort exact matched on key patient characteristics, patients with the most amount of overlap were observed to have shorter total length of surgery; these results disappeared after controlling for the attending surgeon, potentially suggesting that the supervising surgeon drives operative time, rather than the amount of overlap time for any given patient. This contrasts prior work by George et al,¹⁴ which found that patients with a greater surgical overlap had a longer length of surgery during total joint arthroplasty. This difference in results may be due to variability in risk and complexity between the procedures studied, which further highlight the importance of evaluating the impact of overlap duration in a procedure-specific manner.

Limitations

This study was retrospective and is therefore susceptible to sampling biases. Furthermore, all patient health care encounters were recorded within the university-wide electronic health record; therefore, patient encounters that occurred at outside health system may not have been captured, leading to potential under-reporting of morbidity outcomes. This limitation was mitigated by an extensive mean duration of follow-up (38.2 and 43.6 months, for patients with the most and least overlap, respectively). During each follow-up visit, all external health care encounters were asked and recorded. Furthermore, any potential under-reporting of adverse events would be expected to equally affect patients with the most and least amounts of overlap, preserving the internal validity of this study.

While certain demographic and clinical variables were excluded from our CEM protocol, the matching variables used herein were carefully selected from the surgical literature. Race,²⁰⁻²⁴ smoking status,²⁵ BMI,^{26,27} CCI,²⁸⁻³⁰ duration of surgery,^{31,32} and ASA score³³ have all been demonstrated to independently impact post-operative morbidity. By controlling these variables via CEM, we expect to ensure the overall validity of the study findings.

The present analysis used a decile-based matching protocol, which has previously been employed by the present authors.³⁴⁻³⁷ This approach allowed analysis of the extremes of overlap duration, while ensuring

an adequate population sample assessment via exact matching. CEM resulted in 158 patients included in the matched analysis, with a match rate 60.3% among the most overlap cohort. Relative to other matching protocols, CEM includes fewer patients for analysis, but offers more precise control of confounding variables, resulting in enhanced focus on the impact of surgical overlap in 1-level spinal fusion surgery.³⁸

This study examined only single-level, posterior lumbar fusions, enhancing the internal validity of the results at the expense of generalizability. That said, among lumbar fusions, single-level instrumented fusions are the most frequently performed procedure. Furthermore, the differing number of levels of disease presents in different manners, which has implications for results and complications and thus was thought to be too heterogeneous for the intense focus on overlap undertaken herein.³⁹ Additional studies are needed to corroborate these findings across other spinal procedures.

CONCLUSION

In the presence of overlapping surgery, increasing duration of overlap during lumbar fusion does not lead to adverse short-term patient outcomes. When comparing exact-matched patients with the least and most amounts of overlap time, no significant differences were found in any specific morbidity outcome or overall morbidity. Further studies should evaluate longer-term patient outcomes among patients with increased amounts of overlap during lumbar fusion and other neurosurgical procedures.

ACKNOWLEDGMENTS

The EpiLog Project.

REFERENCES

1. Abelson J, Saltzman J, Kowalczyk L. Clash in the name of care. 2015.
2. American College of Surgeon. *Statements on Principles*. 2016.
3. Mello MM, Livingston EH. The evolving story of overlapping surgery. *JAMA*. 2017;318(3):233–234. doi:10.1001/jama.2017.8061
4. Ponce BA, Wills BW, Hudson PW, et al. Outcomes with overlapping surgery at a large academic medical center. *Ann Surg*. 2019;269(3):465–470. doi:10.1097/SLA.0000000000002701
5. Zhang AL, Sing DC, Dang DY, et al. Overlapping surgery in the ambulatory orthopaedic setting. *J Bone Joint Surg Am*. 2016;98(22):1859–1867. doi:10.2106/JBJS.16.00248
6. Goldfarb CA, Rizzo MG, Rogalski BL, Bansal A, Dy CJ, Brophy RH. Complications following overlapping orthopaedic procedures at an ambulatory surgery center. *J Bone Joint Surg Am*. 2018;100(24):2118–2124. doi:10.2106/JBJS.18.00244
7. Glauser G, Osiemo B, Goodrich S, et al. Assessment of short-term patient outcomes following overlapping orthopaedic surgery at a large academic medical center. *J Bone Joint Surg Am*. 2020;102(8):654–663. doi:10.2106/JBJS.19.00554
8. Bohl MA, Mooney MA, Sheehy JP, et al. Overlapping surgeries are not associated with worse patient outcomes: retrospective multivariate analysis of 14 872 neurosurgical cases performed at a single institution. *Neurosurgery*. 2018;83(1):53–59. doi:10.1093/neuros/nyx472
9. Glauser G, Agarwal P, Ramayya AG, et al. Association of surgical overlap during wound closure with patient outcomes among neurological surgery patients at a large academic medical center. *Neurosurgery*. 2019;85(5):E882–E888. doi:10.1093/neuros/nyz142
10. Agarwal P, Ramayya AG, Osiemo B, et al. Association of overlapping neurosurgery with patient outcomes at a large academic medical center. *Neurosurgery*. 2019;85(6):E1050–E1058. doi:10.1093/neuros/nyz243
11. Karsy M, Bowers CA, Scoville J, et al. Evaluation of complications and costs during overlapping transsphenoidal surgery in the treatment of pituitary adenoma. *Neurosurgery*. 2019;84(5):1104–1111. doi:10.1093/neuros/nyy269
12. Min L, Liu M, Zhang W, et al. Outcomes and safety of overlapping surgery in patients undergoing microvascular decompression for hemifacial spasm and trigeminal neuralgia. *World Neurosurg*. 2019;130:e1084–e1090. doi:10.1016/j.wneu.2019.07.092
13. Zygourakis CC, Sizdahkhani S, Keefe M, et al. Comparison of patient outcomes and cost of overlapping versus nonoverlapping spine surgery. *World Neurosurg*. 2017;100:658–664. doi:10.1016/j.wneu.2017.01.064
14. George J, Newman JM, Faour M, et al. Overlapping lower extremity total joint arthroplasty does not increase the risk of 90-day complications. *Orthopedics*. 2018;41(5):e695–e700. doi:10.3928/01477447-20180806-01
15. Saleh A, Thirukumar C, Mesfin A, Molinari RW. Complications and readmission after lumbar spine surgery in elderly patients: an analysis of 2,320 patients. *Spine J*. 2017;17(8):1106–1112. doi:10.1016/j.spinee.2017.03.019
16. Feng C, Zhang Y, Chong F, et al. Establishment and implementation of an enhanced recovery after surgery (ERAS) pathway tailored for minimally invasive transforaminal lumbar interbody fusion surgery. *World Neurosurg*. 2019;129: . doi:10.1016/j.wneu.2019.05.139
17. Gawande A. Why doctors hate their computers. *The New Yorker*. 2018;12.
18. Iacus SM, King G, Porro G. Causal inference without balance checking: coarsened exact matching. *Polit Anal*. 2017;20(1):1–24. doi:10.1093/pan/mpr013
19. Zygourakis CC, Keefe M, Lee J, et al. Comparison of patient outcomes in 3725 overlapping vs 3633 nonoverlapping neurosurgical procedures using a single institution's clinical and administrative database. *Neurosurgery*. 2017;80(2):257–268. doi:10.1093/neuros/nyw067
20. Martsof GR, Barrett ML, Weiss AJ, et al. Impact of race/ethnicity and socioeconomic status on risk-adjusted hospital readmission rates following hip and knee arthroplasty. *J Bone Joint Surg Am*. 2016;98(16):1385–1391. doi:10.2106/JBJS.15.00884
21. LaPar DJ, Bhamidipati CM, Harris DA, et al. Gender, race, and socioeconomic status affects outcomes after lung

- cancer resections in the United States. *Ann Thorac Surg.* 2011;92(2):434–439. doi:10.1016/j.athoracsur.2011.04.048
22. Seicean A, Seicean S, Neuhauser D, Benzel EC, Weil RJ. The influence of race on short-term outcomes after laminectomy and/or fusion spine surgery. *Spine (Phila Pa 1976).* 2017;42(1):34–41. doi:10.1097/BRS.0000000000001657
23. Lemaire A, Cook C, Tackett S, Mendes DM, Shortell CK. The impact of race and insurance type on the outcome of endovascular abdominal aortic aneurysm (AAA) repair. *J Vasc Surg.* 2008;47(6):1172–1180. doi:10.1016/j.jvs.2008.01.033
24. Alish H, Riley III LH, Skolasky RL. Insurance status, geography, race, and ethnicity as predictors of anterior cervical spine surgery rates and in-hospital mortality: an examination of United States trends from 1992 to 2005. *Spine (Phila Pa 1976).* 2009;34(18):1956–1962. doi:10.1097/BRS.0b013e3181ab930e
25. Turan A, Mascha EJ, Roberman D, et al. Smoking and perioperative outcomes. *Anesthesiology.* 2011;114(4):837–846. doi:10.1097/ALN.0b013e318210f560
26. Smits A, Lopes A, Das N, et al. Surgical morbidity and clinical outcomes in ovarian cancer - the role of obesity. *BJOG.* 2016;123(2):300–308. doi:10.1111/1471-0528.13585
27. Wilson JR, Tetreault LA, Schroeder G, et al. Impact of elevated body mass index and obesity on long-term surgical outcomes for patients with degenerative cervical myelopathy: analysis of a combined prospective dataset. *Spine (Phila Pa 1976).* 2017;42(3):195–201. doi:10.1097/BRS.0000000000001859
28. Grossman R, Mukherjee D, Chang DC, et al. Preoperative Charlson comorbidity score predicts postoperative outcomes among older intracranial meningioma patients. *World Neurosurg.* 2011;75(2):279–285. doi:10.1016/j.wneu.2010.09.003
29. Genther DJ, Gourin CG. Effect of comorbidity on short-term outcomes and cost of care after head and neck cancer surgery in the elderly. *Head Neck.* 2015;37(5):685–693. doi:10.1002/hed.23651
30. Rueth NM, Parsons HM, Habermann EB, et al. Surgical treatment of lung cancer: predicting postoperative morbidity in the elderly population. *J Thorac Cardiovasc Surg.* 2012;143(6):1314–1323. doi:10.1016/j.jtcvs.2011.09.072
31. JYS K, Khavanin N, Rambachan A, et al. Surgical duration and risk of venous thromboembolism. *JAMA Surg.* 2015;150(2):110–117. doi:10.1001/jamasurg.2014.1841
32. Chu D, Bakaeen FG, Wang XL, LeMaire SA, Coselli JS, Huh J. Does the duration of surgery affect outcomes in patients undergoing coronary artery bypass grafting? *Am J Surg.* 2008;196(5):652–656. doi:10.1016/j.amjsurg.2008.07.003
33. Dial BL, Esposito VR, Danilkowicz R, et al. Factors associated with extended length of stay and 90-day readmission rates following ACDF. *Global Spine J.* 2020;10(3):252–260. doi:10.1177/2192568219843111
34. Farooqi AS, Borja AJ, Detchou DKE, et al. Duration of overlap during lumbar fusion does not predict outcomes. *Clin Neurol Neurosurg.* 2021;205:106610. doi:10.1016/j.clineuro.2021.106610
35. Borja AJ, Glauser G, Strouz K, et al. Use of the LACE+ index to predict readmissions after single-level lumbar fusion. *J Neurosurg Spine.* 2021;2021:1–9. doi:10.3171/2021.9.SPINE21705
36. Farooqi AS, Borja AJ, Detchou DKE, et al. Postoperative outcomes and the association with overlap before or after the critical step of lumbar fusion. *J Neurosurg Spine.* 2021;2021:1–10. doi:10.3171/2021.5.SPINE202105
37. Farooqi AS, Borja AJ, Detchou DKE, et al. Overlap before the critical step of lumbar fusion does not lead to increased short-term morbidity. *Neurosurgery.* 2021;89(6):1052–1061. doi:10.1093/neuros/nyab360
38. Guy D, Karp I, Wilk P, Chin J, Rodrigues G. Propensity score matching versus coarsened exact matching in observational comparative effectiveness research. *J Comp Eff Res.* 2021;10(11):939–951. doi:10.2217/cer-2021-0069
39. Harada GK, Khan JM, Vetter C, et al. Does the number of levels fused affect spinopelvic parameters and clinical outcomes following posterolateral lumbar fusion for low-grade spondylolisthesis? *Global Spine J.* 2021;11(1):116–121. doi:10.1177/2192568220901527

Funding: N.R.M. received support from the Kevin and Bernadette McKenna Family Research Fund.

Declaration of Conflicting Interests: The authors report no conflicts of interest in this work.

Ethics Committee Approval: This study was approved by the IRB at the University of Pennsylvania. IRB number for this study is: 831423. A waiver of informed consent was granted by the University of Pennsylvania IRB as this study was considered to be minimal risk to patients. All ethical guidelines and rules were followed to protect patient privacy

Contributorship Statement: Conception and design: N.R.M. and A.S.F. Acquisition of data: N.R.M. and A.S.F. Analysis and interpretation of data: A.S.F., K.S., S.D.M., and N.R.M. Drafting the article: A.S.F., D.K.E.D.D., G.G., A.J.B., and N.R.M. Critically revising the article: A.S.F., D.K.E.D.D., G.G., A.J.B., K.S., S.D.M., and N.R.M. Approved the final version of the manuscript on behalf of all authors: A.S.F., D.K.E.D.D., G.G., A.J.B., K.S., S.D.M., and N.R.M. Administrative/technical/material support: D.K.E.D.D., G.G., A.J.B., N.R.M. Study supervision: N.R.M.

Corresponding Author: Neil R. Malhotra, Department of Neurosurgery, University of Pennsylvania, 3rd Floor Silverstein Pavilion, 3400 Spruce St, PA 19104, USA; neil.malhotra@pennmedicine.upenn.edu

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2022 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.