

Sacral Dysmorphism and Lumbosacral Transitional Vertebrae (LSTV) Review

DAVID M. MATSON, MD, LAUREN M. MACCORMICK, MD, JONATHAN N. SEMBRANO, MD,
DAVID W. POLLY JR, MD

Department of Orthopaedic Surgery, University of Minnesota, Minneapolis, Minnesota

ABSTRACT

Background: Anatomic variation in the relationship between the lumbar spine and sacrum was first described in the literature nearly a century ago and continues to play an important role in spine deformity, low back pain (LBP), and pelvic trauma. This review will focus on the clinical and surgical implications of abnormal lumbosacral anatomy in the context of sacroiliac joint (SIJ) disease, spine deformity, and pelvic trauma.

Methods: A PubMed search using the keywords “lumbosacral transitional vertebrae,” “LSTV,” “transitional lumbosacral vertebrae,” “TLSTV,” and “sacral dysmorphism” was performed. The articles presented here were evaluated by the authors.

Clinical Significance: The prevalence of LSTV varies widely in the literature from 3.9% to 35.6% in the spine literature, and sacral dysmorphism is described in upwards of 50% of the population in the trauma literature. The relationship between LSTV and LBP is well established. While there is no agreed-on etiology, the source of pain is multifactorial and may be related to abnormal biomechanics and alignment, disc degeneration, and arthritic changes.

Surgical Implications: Understanding abnormal lumbosacral anatomy is crucial for preoperative planning of SIJ fusion, spine deformity, and pelvic trauma surgery. LSTV can alter spinopelvic parameters crucial in planning spine deformity correction. Traditional safe zones for sacroiliac screw placement do not apply in the first sacral segment in sacral dysmorphism and risk iatrogenic nerve injury.

Conclusions: LSTV and sacral dysmorphism are common anatomic variants found in the general population. Abnormal lumbosacral anatomy plays a significant role in clinical evaluation of LBP and surgical planning in SIJ fusion, spine deformity, and pelvic trauma. Further studies evaluating the influence of abnormal lumbosacral anatomy on LBP and surgical technique would help guide treatment for these patients.

Other and Special Categories

Keywords: lumbosacral transitional vertebrae, LSTV, sacral dysmorphism, transitional lumbosacral vertebrae, low back pain, Bertolotti syndrome, sacroiliac joint fusion

INTRODUCTION

Anatomic variation in the relationship between the lumbar spine and sacrum was first described in the literature almost a century ago and continues to play an important role in spine deformity, low back pain (LBP), and pelvic trauma.^{1,2} As the indications for posterior pelvic screw fixation have expanded, the recognition of sacral dysmorphism as an important consideration has grown. Additionally, the understanding of the importance of lumbosacral transitional vertebrae (LSTV) in spine mechanics and surgical fixation has evolved. LSTV are a congenital anomaly and common anatomic variant that occurs when there is partial or complete fusion between the final lumbar vertebra and first sacral segment. Lumbarization of the first sacral segment

occurs with separation, either partial or complete, of S1 from the rest of the sacrum, creating a sixth lumbar vertebra. Sacralization of the last lumbar vertebra occurs when the last lumbar vertebra is incorporated into the sacrum, resulting in 4 lumbar vertebrae.³ Sacralization of L5 (7.5%) is slightly more common in the general population compared with lumbarization of S1 (5.5%).⁴

Genetic factors may play a role in the development of LSTV. Hox genes (Hox-10, Hox-11) have been found to play a significant role in the process of vertebral body segmentation and development.⁵ The development of vertebral bodies and disc formation occurs during the fourth week of embryonic development; however, the process by which the vertebrae consolidate and fuse into adult lumbosacral anatomy occurs through the fourth




Figure 1. Castellvi-Chan classification system of lumbosacral transitional anatomy. Image reprinted with permission from Castellvi et al.⁶

decade of life.⁶ The relative number of lumbar and sacral segments is influenced by load transmission through the lumbosacral region during development and is associated with the bipedal evolution of humans.^{3,7}

CLASSIFICATION AND PREVALENCE

The spectrum of anatomic variation in this population was described by Castellvi et al⁸ in 1984 (Figure 1). Using radiographs, the authors classified 7 different patterns of transitional vertebrae into 4 types based on morphology (types I–IV)



Figure 2. Castellvi-Chan type IIB lumbosacral transitional vertebra identified on pelvis outlet view radiograph.

and 2 variations based on laterality (for types I–III, A = unilateral and B = bilateral). Type I, also known as a dysplastic transverse process, is a large transverse process triangular in shape with dimensions measuring at least 19 mm in rostrocaudal width. Type II, or incomplete lumbarization/sacralization, is a large transverse process that follows the contour and articulates with the sacrum but is not fused, creating a diarthroidal joint between the final lumbar vertebra and the first sacral segment (Figure 2). Type III, or complete lumbarization/sacralization, is a large transverse process with bony fusion to the sacrum (Figure 3). Type IV exhibits lumbarization/sacralization that is incomplete (type II) on one side and complete (type III) on the contralateral side.

O'Driscoll et al⁹ developed a classification system based on T1- and T2-weighted magnetic resonance imaging (MRI) and the disc morphology between the uppermost sacral segment and the remainder of the sacrum. Type 1 describes no disc material present. Type 2 has small residual disc that does not extend the whole diameter from anterior to posterior. Type 3 has a well-formed disc extending the entire anteroposterior (AP) diameter of the sacrum. Type 4 has a well-formed disc extending the entire AP diameter of the sacrum and an associated abnormal upper sacral segment. The authors found good association between type 4 on MRI and fused LSTV on radiographs.

Sacral dysmorphism has been described in the orthopedic trauma literature more recently. Miller and Routt⁶ described several important findings on pelvic radiographs of sacral dysmorphism. The 7 characteristics may be present to varying degrees




Figure 3. (A) Castellvi-Chan type IIIA and (B) Castellvi-Chan type IIIB lumbosacral transitional vertebra identified on pelvis outlet view radiographs.

and include (1) collinearity of the upper portion of the sacrum and iliac crests on the outlet radiograph, (2) presence of mammillary bodies at the sacral ala on outlet radiograph, (3) noncircular anterior sacral foramina on outlet radiograph, (4) residual upper sacral disks on lateral radiograph, (5) tongue-and-groove sacroiliac joint surface on axial computed tomographic (CT) scan, (6) cortical indentation of the ala on the inlet radiograph, and (7) acute alar slope on the lateral radiograph (Figures 4 and 5). The subtler findings may be better appreciated on CT of the pelvis.

The prevalence of LSTV in the general population varies significantly in the literature, ranging from 4% to 35.6% in the spine literature.^{2,4,8,10-16}



Figure 4. Mammillary bodies and abnormal sacral neural exit present in the uppermost sacral segment of a dysmorphic sacrum.

Nardo et al¹⁴ evaluated radiographs of 4636 participants and determined the overall prevalence of LSTV to be 18.1%, with types I and II comprising nearly 80% of LST. The authors also found a significantly higher rate in men compared to women (28.1% vs 11.1%). Apazidas et al¹¹ reported the highest rate of LSTV at 35.6% and determined the prevalence of each of the Castellvi-Chan classification groups; type IA was most common at 14.7%. The prevalence of sacral dymorphism described in the trauma literature is estimated to be upwards of 50% (Table 1).^{13,17}

CLINICAL SIGNIFICANCE

LSTV and sacral dymorphism result in distinct anatomic changes at the lumbosacral junction and in the surrounding structures. Mahato¹⁸ found that



Figure 5. Tongue-and-groove sacroiliac joint surface of sacral dymorphism identified on CT scan.

Table 1. Incidence of lumbosacral transitional vertebrae or sacral dysmorphism.

Study	Incidence of Lumbosacral Transitional Vertebrae or Sacral Dysmorphism, %
Castellvi et al ⁸	30
Luoma et al ¹⁶	30
Wu et al ¹³	16.7
Gardner et al ¹⁷	44
Mahato et al ³	3.9
Apazidas et al ¹¹	35.6
Nardo et al ¹⁴	18.1
	Type I: 41.72
	Type II: 41.4
	Type III: 11.5
	Type IV: 5.2
Tang et al ¹⁵	15.8
	Type I: 44.8
	Type II: 43.2
	Type III: 7.2
	Type IV: 4.8

LSTV articulations at L5–S1 were associated with increased lordotic curves, L5 vertebral heights, and pedicle and angular dimensions. Furthermore, L5–S1 fusions were associated with smaller disc heights, wider and shorter L5 pedicles, narrower and taller transverse processes, and straighter spines overall. According to Mahato¹⁹, sacralization reduces S1 pedicle height and sagittal angulation of the pedicles while increasing the downward slope. Lumbarization was associated with more obtuse pedicles in the sagittal plane and a smaller length between the facet and sacral promontory. The anatomic variations may affect spinal instrumentation at this level and must be taken into consideration with regard to preoperative planning.

Mechanical LBP is one of the most common musculoskeletal concerns. LBP has been associated with sacroiliac dysfunction as well as LSTV. The association between LSTV and LBP was first established in 1917 by Mario Bertolotti²⁰ and is known as Bertolotti syndrome. Tang et al¹⁵ found a significant relationship between LSTV and low back and buttock pain. The authors demonstrated odds ratios of 2.56 and 4.28 for LBP and 5.38 and 6.82 for gluteal pain in Castellvi–Chan types II and IV LSTV, respectively. While the exact mechanism is not understood, many studies have suggested that pain may be related to degeneration of the adjacent disc segments and facet joints due to hypermobility and increased forces at the level just cephalad to the transitional segment.^{12,16,21,22} As described previously, Mahato¹⁸ demonstrated numerous ways in which LSTV affects the load-bearing mechanics of the lumbosacral spine and contributes to mechan-

ical LBP, including hypolordosis associated with L5–S1 fusion and hyperlordosis associated with L5–S1 articulation. Farshad-Amacker et al²³ demonstrated a protective effect for the disc at the transitional segment and increased degenerative changes at the adjacent cephalad disc, particularly for Castellvi–Chan types III and IV LSTV. The relationship between LSTV and disc degeneration was further evaluated by Ahn et al,²² who determined that patients with LSTV undergoing microdiscectomy have clinically worse outcomes compared to those with “normal” lumbosacral anatomy. Illeez et al¹² evaluated the relationship between LSTV and sacroiliac joint dysfunction and LBP in 700 subjects. The authors found statistically significant associations between LSTV and sacroiliac dysfunction (28.5%) and between LSTV and LBP (26%). While the disc adjacent to the LSTV may experience increased stresses leading to degenerative change, the relative segmental innervation is unchanged in patients with sacral lumbralization compared to those with 5 lumbar vertebral bodies based on electromyographic findings.^{24,25}

SURGICAL IMPLICATIONS

LSTV has implications in the preoperative planning for surgical correction in spine deformity cases. Spinal correction surgery depends on accurate measurements of spinopelvic and global spinal parameters. A number of these parameters rely on the accurate identification of the sacral endplate and may be influenced by the relative height and translation of the sacral endplate. In patients with LSTV, the identification of the sacral endplate may not be entirely clear around the transitional segment. Zhou et al²⁵ measured these spinal parameters for patients with Castellvi–Chan types III and IV LSTV using both the cephalad and the caudal segments as the sacral endplate. The authors found that the pelvic incidence (PI) differed by $\sim 20^\circ$ between cephalad and caudal. Most of the parameters measured, including PI, lumbar lordosis (LL), PI–LL mismatch, pelvic tilt, sacral slope, sagittal vertical axis, and T1–pelvic angle, demonstrated significant differences between cephalad and caudal measurements. Thoracic kyphosis and spinal inclination (T1SPi) were not affected by cephalad and caudal positioning of the sacral endplate. The significant differences in the measurements may have adverse effects on the preoperative surgical planning for spine deformity cases. Furthermore,




Figure 6. The dysmorphic first sacral segment did not allow for a transiliac, transsacral screw at this level, as was used at S2.

Khalsa et al²⁶ found significant variability between surgeons in assessing spinopelvic parameters in patients with LSTV. While there are not yet standardized measures or agreed-on alignment goals for patients with LSTV, the variations caused by the deformity must be taken into consideration and play an important role in preoperative planning.

Pelvic ring injuries can cause widening of the sacroiliac joint or a sacral fracture, and treatment typically involves percutaneous reduction methods and fixation. The treating surgeon must understand the safe zones for screw placement in normal anatomy in order to avoid the risk of iatrogenic damage to nerves traversing through the sacral neural tunnels. Additionally, sacroiliac joint fusion utilizes similar safe zones in the placement of implants. Normal anatomy exhibits bilateral, ellipsoid, osseous areas bounded cranial-anteriorly by sacral alar cortical bone and caudal-posteriorly by first sacral neural tunnel. The most constrained area is between the sacral ala and the first sacral neural tunnel. On a true lateral radiograph, screw trajectory caudal and posterior relative to the sacral ala is important in order to avoid injury to the L5 nerve root.⁶

A dysmorphic sacrum has a distorted, misshapen, and ellipsoid upper sacral segment that limits the area available for screw placement. Oblique dysmorphic alar osteology makes transiliac, transsacral screw fixation unavailable in S1. The preferred safe zone for a sacroiliac screw in S1 is oblique from caudal to cranial and posterior to anterior, and there is not a safe corridor for horizontal screw placement in the AP plane. Alternatively, the safe zone is similar to normal in the S2 segment and is amenable to

transiliac, transsacral screw (Figure 6). Preoperative evaluation of the pelvic CT scan is important in planning the appropriate sacral segments and trajectories for safe percutaneous instrumentation. In addition, intraoperative CT navigation may be useful in avoiding potential complications for patients with sacral dysmorphism requiring SIJ fusion or posterior pelvic ring fixation.²⁷

REFERENCES

1. Brailsford JF. Deformities of the lumbosacral region of the spine. *Br J Surg.* 1929;16(64):562–627.
2. Hasner E, Jacobsen HH, Schalimtzek M, Snorrason E. Lumbosacral transitional vertebrae: a clinical and roentgenologic study of 400 cases of low back pain. *Acta Radiol.* 1953;39(3):225–230.
3. Mahato NK. Morphological traits in sacra associated with complete and partial lumbarization of first sacral segment. *Spine J.* 2010;10(10):910–915.
4. Bron JL, van Royen BJ, Wuisman PI. The clinical significance of lumbosacral transitional anomalies. *Acta Orthop Belg.* 2007;73(6):687–695.
5. Carapuço M, Nóvoa A, Bobola N, Mallo M. Hox genes specify vertebral types in the presomitic mesoderm. *Genes Dev.* 2005;19(18):2116–2121.
6. Miller AN, Routt ML. Variations in sacral morphology and implications for iliosacral screw fixation. *J Am Acad Orthop Surg.* 2012;20(1):8–16.
7. Abitbol MM. Evolution of the sacrum in hominoids. *Am J Phys Anthropol.* 1987;74(1):65–81.
8. Castellvi AE, Goldstein LA, Chan DP. Lumbosacral transitional vertebrae and their relationship with lumbar extradural defects. *Spine (Phila Pa 1976).* 1984;9(5):493–495.
9. O'Driscoll CM, Irwin A, Saifuddin A. Variations in morphology of the lumbosacral junction on sagittal MRI: correlation with plain radiography. *Skeletal Radiol.* 1996;25(3):225–230.
10. Konin GP, Walz DM. Lumbosacral transitional

vertebrae: classification, imaging findings, and clinical relevance. *Am J Neuroradiol.* 2010;31(10):1778–1786.

11. Apazidis A, Ricart PA, Diefenbach CM, Spivak JM. The prevalence of transitional vertebrae in the lumbar spine. *Spine J.* 2011;11(9):858–862.

12. Illeez OG, Atici A, Ulger EB, Kulcu DG, Ozkan FU, Aktas I. The transitional vertebra and sacroiliac joint dysfunction association. *Eur Spine J.* 2018;27(1):187–193.

13. Wu LP, Li YK, Li YM, Zhang YQ, Zhong SZ. Variable morphology of the sacrum in a Chinese population. *Clin Anat.* 2009;22(5):619–626.

14. Nardo L, Alizai H, Virayavanich W, et al. Lumbosacral transitional vertebrae: association with low back pain. *Radiology.* 2012;265(2):497–503.

15. Tang M, Yang XF, Yang SW, et al. Lumbosacral transitional vertebra in a population-based study of 5860 individuals: prevalence and relationship to low back pain. *Eur J Radiol.* 2014;83(9):1679–1682.

16. Luoma K, Vehmas T, Raininko R, Luukkonen R, Riihimäki H. Lumbosacral transitional vertebra: relation to disc degeneration and low back pain. *Spine (Phila Pa 1976).* 2004;29(2):200–205.

17. Gardner MJ, Morshed S, Nork SE, Ricci WM, Chip Routt ML. Quantification of the upper and second sacral segment safe zones in normal and dysmorphic sacra. *J Orthop Trauma.* 2010;24(10):622–629.

18. Mahato NK. Disc spaces, vertebral dimensions, and angle values at the lumbar region: a radioanatomical perspective in spines with L5-S1 transitions. Clinical article. *J Neurosurg Spine.* 2011;15(4):371–379.

19. Mahato NK. Pedicular anatomy of the first sacral segment in transitional variations of the lumbo-sacral junction. *Spine (Phila Pa 1976).* 2011;36(18):E1187–E1192.

20. Bertolotti M. Contributo alla conoscenza dei vizi di differenziazione regionale del rachide con speciale riguardo all'assimilazione sacrale della v. lombare [Contribution to the knowledge of the vices of regional differentiation of the spine with special regard to the sacral assimilation of the lumbar]. *Radiol Med.* 1917;4:113–144.

21. Quinlan JF, Duke D, Eustace S. Bertolotti's syndrome. A cause of back pain in young people. *J Bone Joint Surg Br.* 2006;88(9):1183–1186.

22. Ahn SS, Chin DK, Kim SH, Kim DW, Lee BH, Ku MG. The clinical significance of lumbosacral transitional vertebrae on the surgical outcomes of lumbar discectomy: a retrospective cohort study of young adults. *World Neurosurg.* 2017;99:745–750.

23. Farshad-Amacker NA, Herzog RJ, Hughes AP, Aichmair A, Farshad M. Associations between lumbosacral transitional anatomy types and degeneration at the transitional and adjacent segments. *Spine J.* 2015;15(6):1210–1216.

24. Hinterdorfer P, Parsaei B, Stieglbauer K, Sonnberger M, Fischer J, Wurm G. Segmental innervation in lumbosacral transitional vertebrae (LSTV): a comparative clinical and intraoperative EMG study. *J Neurol Neurosurg Psychiatry.* 2010;81(7):734–741.

25. Zhou PL, Moon JY, Tishelman JC, et al. Interpretation of spinal radiographic parameters in patients with transitional lumbosacral vertebrae. *Spine Deform.* 2018;6(5):587–592.

26. Khalsa AS, Mundis GM Jr, Yagi M, et al. Variability in assessing spinopelvic parameters with lumbosacral transitional vertebrae: inter- and intraobserver reliability among spine surgeons. *Spine (Phila Pa 1976).* 2018;43(12):813–816.

27. Khan JM, Lara DL, Marquez-Lara A, Rosas S, Hasty E, Pilson HT. Intraoperative CT and surgical navigation for iliosacral screws: technique for patients with sacral dysmorphism. *J Orthop Trauma.* 2018;32 Suppl 1:S24–S25.

Disclosures and COI: Each author certifies that he or she has no commercial associations (eg, consultancies, stock ownership, equity interest, patent/licensing arrangements, etc.) that might pose a conflict of interest in connection with the submitted work. This paper is exempt from institutional review board review, as it is not human subjects research. There was no external source of funding for this study.

Corresponding Author: Jonathan N. Sembrano, MD, Department of Orthopaedic Surgery, University of Minnesota, 2450 Riverside Avenue South, Suite R200, Minneapolis, MN 55454. Phone: (612) 273-7991; Email: sembr001@umn.edu.

Published 10 February 2020

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2020 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.