

Are Cervical Disc Arthroplasty Medicare Reimbursement Trends Sustainable?

Emily S. Mills, Cory K. Mayfield, Tara Shelby, Andy T. Ton, Raymond J. Hah and Ram K. Alluri

Int J Spine Surg 2023, 17 (2) 222-229 doi: https://doi.org/10.14444/8428 https://www.ijssurgery.com/content/17/2/222

This information is current as of May 9, 2025.

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at: http://ijssurgery.com/alerts



International Journal of Spine Surgery, Vol. 17, No. 2, 2023, pp. 222–229 https://doi.org/10.14444/8428 © International Society for the Advancement of Spine Surgery

Are Cervical Disc Arthroplasty Medicare Reimbursement Trends Sustainable?

EMILY S. MILLS, MD¹; CORY K. MAYFIELD, MD¹; TARA SHELBY, BS¹; ANDY T. TON, BS¹; RAYMOND J. HAH, MD¹; AND RAM K. ALLURI, MD¹

¹Keck School of Medicine, Department of Orthopaedic Surgery, University of Southern California, Los Angeles, CA, USA

ABSTRACT

Background: Cervical disc arthroplasty (CDA) was originally approved by the US Food and Drug Administration (FDA) in 2007 as a motion-sparing procedure to treat cervical degenerative disc disease. Since then, promising results from randomized control trials have led to increasing popularity. However, data discussing monetary trends are limited. The aim of this study was to determine how utilization, hospital charges, and Medicare physician reimbursement for CDA have changed over time.

Methods: In this retrospective cohort study, International Classification of Diseases procedure codes were used to identify all patients who underwent CDA from 2007 to 2017 in the National Inpatient Sample database. The Physician Fee Schedule Look-up Tool from the Centers for Medicare and Medicaid Services was queried for primary CDA using current procedural terminology codes to determine Medicare physician reimbursement from 2009 to 2021. Nominal monetary values were adjusted for inflation using the Consumer Price Index and inflation-adjusted data reported in 2021 US dollars.

Results: A total of 33,079 weighted patients who underwent CDA were included for analysis. CDA utilization increased by 183% from 2007 to 2017, with Medicare beneficiary utilization increasing 149%. Inflation-adjusted total hospital charges for CDA increased by 22.4%. However, inflation-adjusted Medicare physician reimbursement fell by 1.20% per year, demonstrating a total decrease of 12.9%, starting at \$1928 in 2009 and declining to \$1679 in 2021.

Conclusions: While utilization and total hospital charges for CDA continue to rise, Medicare physician reimbursement has not shown the same trend. In fact, inflation-adjusted reimbursement has seen a steady decline since FDA approval in 2007. If this trend persists, it may become unsustainable for physicians to continue offering CDA to Medicare patients. As disproportionate increases in hospital charges incentivize a transition to outpatient CDA, stricter patient selection criteria associated with outpatient procedures may create health care disparities for Medicare patients and those with higher comorbidity burden.

Clinical Relevance: This study shows the decreasing reimbursement trends for CDA, which may disproportionately affect Medicare patients and those with increased comorbidities.

Level of Evidence: 3.

Cervical Spine

Keywords: cervical spondylosis, disc degeneration, health expenditures

INTRODUCTION

Neck pain represents a large burden to the aging population and was recently found to be a leading cause of disability worldwide.¹ Cervical degenerative disc disease (DDD) is a known cause of neck pain, cervical radiculopathy, and myelopathy.² Historically, treatment for cervical DDD refractory to nonoperative management included anterior cervical discectomy and fusion (ACDF), which was originally described in the 1950s, and many studies have showed improved and sustained clinical outcomes after surgery.^{3–5} However, complications from fusion, including adjacent segment disease and pseudarthrosis, can lead to increased revision rates, particularly in multilevel surgery, and diminished clinical outcomes.^{6–11} In part, these limitations led to the advent and evolution of

motion-sparing technologies for the anterior column of the cervical spine.

Cervical disc arthroplasty (CDA) was initially approved by the US Food and Drug Administration (FDA) in 2007 and was introduced as a motion-sparing procedure to treat cervical DDD. Initial randomized controlled trials showed promising results, demonstrating mostly noninferior or superior results to ACDF with regard to clinical outcomes and lower rates of adjacent segment disease.^{12–16} Recent studies are now examining the efficacy of CDA with expanded indications in patients with cervical spondylosis, myelopathy, or kyphotic alignment, and the initial FDA investigational device exemption studies now have long-term follow-up with maintained results.^{17–20} Given the promising data, utilization of CDA has been increasing. A recent study by Niedzilak et al showed a 20% increase in utilization from 2005 to 2014 in a national database.²¹ However, data discussing monetary trends in CDA are limited. The aim of this study was to determine how utilization, cost, and Medicare physician reimbursement for CDA have changed over time.

METHODS

Database Description

The International Classification of Diseases (ICD) procedure codes were used to query the National Inpatient Sample (NIS) database to identify and analyze patients who underwent CDA.²² NIS is the largest publicly available all-payer inpatient health database, with a 20% stratified sample of all discharges from US community hospitals participating in the Healthcare Cost and Utilization Project, beginning from 2012. The large sample size of 35 million hospitalizations and 7 million hospital stays yearly allows for identification of uncommon treatments, special patient populations, and rare conditions. The NIS is the only national database that uses discharge weights, which allow accurate estimations of nationwide incidence of diseases and procedures.²²

Patient Selection

Procedure codes ICD-9-Procedure Coding System (PCS) 84.62 and ICD-10-PCS 0RR30JZ were used to identify patients undergoing CDA, and the number of cases was documented from 2007 to 2017. The NIS database contains data pertaining to patient disease status, procedure type, and demographics as well as hospital characteristics. Descriptive variables included in our analysis include total charges, age, sex, race, primary payer method, median household income, length of stay, bed size, ownership of hospital, and teaching status. Total charge is presented with inflation-adjusted 2021 US dollars (USD).²³ Each of the aforementioned elements was charted against the same time frame of 2007 to 2017.

Physician Reimbursement

The American Academy of Orthopedic Surgeons coding reference was queried to determine the CPT codes most frequently utilized in primary CDA.²⁴ Next, the Physician Fee Schedule Look-up Tool from the Centers for Medicare and Medicaid Services was

queried for primary CDA using CPT code 22856.²⁵ The reported reimbursement data average Medicare physician surgeon fees from over 100 different centers across the United States for each year between 2000 and 2021. While the Physician Fee Schedule Look-up Tool contains data from 2000 to 2021, CDA was not recognized as an individual CPT within this database until 2009; thus, only the years 2009 through 2021 were included in our analysis.

All nominal monetary values were adjusted for inflation using the Consumer Price Index, and inflationadjusted data were reported in 2021 USD. The latest Consumer Price Index data were acquired from the US Department of Labor's Bureau of Labor Statistics.²⁶ Physician fee and procedure charge trends were analyzed based on inflation-adjusted 2021 USD.²⁷ The average annual and total percentage change in reimbursement were calculated based on these trends.

All databases used in this study are publicly available national databases without patient identifiers. Therefore, institutional review board approval and informed consent were not required to complete the study.

Statistical Methods

We stratified the cohort by year, 2007 to 2017, and looked at various patient and hospital characteristics. Linear regression was then conducted to obtain P values, indicating whether there was a relationship between these characteristics and year of surgery. STATA version 16.1 (Stata Corporation) was used to perform all analyses, and a P value of <0.05 was considered statistically significant.

RESULTS

Demographic Data

Our query of the NIS database resulted in 33,079 weighted patients who underwent CDA from 2007 to 2017. The majority were Caucasian (72.92%, P < 0.001), had private insurance (64.8%, P < 0.001), and had a household income above the 75th percentile (29.0%, P < 0.001). Most of the procedures were performed at urban teaching hospitals (38.8%, P < 0.001).

Length of stay and female gender were both consistent from 2007 to 2017 (P = 0.185 and P = 0.26, respectively). Age significantly increased over the 10year period from 46.6 to 47.3 years (P = 0.0043). Other patient characteristics of race, insurance type, and patient zip code and hospital characteristics of teaching status, ownership, and region were all significantly associated with year of surgery (Table).

Age, y 46.60 ± 1.2 45 Length of stay, d 1.72 ± 0.2 1 Female 163 (51.4) 2 Caucasian 206 (65.0) 33 Medicare 29 (9.2) 3	5.76 ± 0.8 $.62 \pm 0.2$						-107	CT N7	0107	1707	TIMOT	1 10100
Length of stay, d 1.72 ± 0.2 1 Female 163 (51.4) 2 Caucasian 206 (65.0) 33 Medicare 29 (9.2) 3	$.62 \pm 0.2$	46.34 ± 0.7	45.79 ± 0.8	47.56 ± 0.9	45.86 ± 1.0	46.78 ± 0.9	47.29 ± 0.9	47.18 ± 0.8	46.63 ± 0.7	47.32 ± 0.7	46.7 ± 0.3	0.0043
Female 163 (51.4) 2 Caucasian 206 (65.0) 33 Medicare 29 (9.2) 3		1.74 ± 0.2	1.50 ± 0.2	1.72 ± 0.2	1.67 ± 0.2	1.67 ± 0.2	1.56 ± 0.4	1.65 ± 0.1	1.48 ± 0.1	1.59 ± 0.1	1.6 ± 0.1	0.1853
Caucasian 206 (65.0) 33 Medicare 29 (9.2) 3	(13 (49.9)	333 (49.9)	323 (52.1)	286 (48.8)	192 (45.6)	234 (49.1)	303 (50.2)	371 (53.7)	446 (52.5)	451 (53.1)	3375 (50.9)	0.26
Medicare 29 (9.2)	84 (70.2)	485 (72.7)	469 (75.7)	410 (70.0)	316 (75.1)	335 (70.2)	421 (69.8)	511 (74.0)	655 (77.1)	641 (77.1)	4833 (72.9)	<0.001
	24 (4.4)	49 (7.3)	42 (6.7)	56 (9.6)	34 (8.1)	45 (9.4)	66 (10.9)	84 (12.2)	88 (10.4)	107 (12.6)	624 (9.4)	<0.001
payment				t cu								
Private insurance 318 (68.8) 30	. (8.co) 00	397 (10.96)	(c./c) 0 c c	354 (60.7)	(0.20) 202	(0.69) 155	381 (6.50)	444 (64.3)	541 (63.7)	000 (04.8)	4294 (64.8)	
75 th -100 th zip 86 (27.1) 1 ⁴	46 (26.7)	149 (22.3)	148 (23.9)	196 (33.4)	126 (29.9)	151 (31.7)	182 (30.2)	205 (29.7)	257 (30.2)	274 (32.3)	1920 (29.0)	<0.001
code quartile												
Large bed size 150 (47.3) 3	17 (58.0)	390 (58.5)	332 (53.5)	388 (66.2)	247 (58.7)	295 (61.8)	273 (45.3)	328 (47.5)	417 (49.1)	433 (51.0)	3570 (53.86)	<0.001
Private, 65 (20.5) 10	07 (19.6)	86 (12.9)	129 (20.8)	127 (21.7)	90 (21.4)	127 (26.6)	131 (21.7)	154 (22.3)	187 (22.0)	176 (20.7)	1631 (24.6)	<0.001
investment												
hospital												
Urban 158 (49.8) 2:	50 (45.7)	373 (57.5)	302 (49.2)	219 (54.9)	199 (47.3)	239 (50.1)	157 (26.0)	208 (30.1)	254 (29.9)	214 (25.2)	2573 (38.8)	<0.001
Rural 10 (3.15) 3	32 (5.85)	23 (3.54)	20 (3.26)	13 (2.24)	17 (4.04)	11 (2.31)	27 (4.48)	35 (5.07)	26 (3.06)	26 (3.06)	240 (3.6)	
Western region 69 (21.77) 13	30 (23.77)	131 (19.64)	131 (21.13)	219 (37.37)	126 (29.93)	151 (31.66)	180 (29.85)	185 (26.77)	257 (30.24)	288 (33.92)	1867 (28.2)	<0.001

lisc arthroplasty per year, from 2007 to 2017.	10 2011 2012 201
ho underwent cervical c	2009 20
laracteristics of patients w	2007 2008
Table. Baseline ch	Demographic

	Downloaded from https://www.ijssurgery.com/ by guest on May 9, 2025
224	International Journal of Spine Surgery, Vol. 17, No. 2



Figure 1. Total cervical disc arthroplasty utilization from 2007 to 2017.

Utilization

The weighted total number of CDA procedures is represented in Figure 1. Over the entire study period, there was a 183% increase in the total number of CDA procedures, from 1502 in 2007 to 4245 in 2017. From 2007 to 2009, there was a sharp increase in case volume to 3428 annual cases followed by a steady decline in case volume to 2105 in 2012. More recent years have demonstrated an increase in annual cases to 4245 in 2017. Medicare beneficiary utilization has increased by 149% over the study period, starting at 43 per 100,000 in 2007 and increasing to 107 per 100,000 in 2017 (Figure 2).

Cost Data

Inflation-adjusted total hospital charges for CDA demonstrated an overall increase of 22.4% from 2007 to 2017, starting at \$76,117 in 2007 and increasing to \$93,172 in 2017 (Figure 3). Prior to inflation adjustment, physician reimbursement remained constant throughout the study period, from \$1556 in 2009 to \$1679 in 2021, with an average change of +0.62% per year. When correcting for inflation, the physician reimbursement for CDA has seen a steady decline from 2009 to 2021 (Figure 4). Values either consistently fell or remained similar from year to year, starting at \$1928 in 2009 and declining to \$1679 in 2021. Over the study period, there was an average decrease in reimbursement of 1.20% per year and a total decrease of 12.9%.



Figure 2. Medicare beneficiary cervical disc arthroplasty utilization from 2007 to 2017.



Figure 3. Inflation-adjusted hospital charges for cervical disc arthroplasty from 2007 to 2017. USD, US dollars.

DISCUSSION

In the present study, we found that CDA utilization increased by 183% from 2007 to 2017. Specifically, Medicare beneficiary utilization increased by 149%. The total hospital charge for the procedure increased by 22.4%, while inflation-adjusted Medicare physician reimbursement decreased by 12.9% from 2009 to 2021.

Several randomized trials have shown the benefit of CDA. Initial investigation device exemption studies showed noninferior or superior outcomes when compared with ACDF for both single-level and 2-level CDA.^{12,14,28,29} A 7-year follow-up study published in 2021 showed decreased adjacent segment disease in 2-level CDA vs 2-level ACDF and noninferior outcomes of 1-level CDA vs 1-level ACDF.³⁰ A similar study investigating single-level CDA vs ACDF found a lower rate of reoperation in the CDA cohort compared with the ACDF cohort at 7 years.¹³ Furthermore, a 10-year follow-up study confirmed a lower rate of reoperation and concluded that CDA is a safe and effective alternative to ACDF for cervical DDD.¹⁵ In addition, a meta-analysis of 14 randomized



Figure 4. Inflation-adjusted Medicare physician reimbursement from 2009 to 2021. USD, US dollars.

controlled trials found a decreased risk of adjacent segment disease in CDA compared with ACDF.¹⁶ The cost-effectiveness of CDA has also been studied, with several recent reports finding CDA is more cost-effective than ACDF.^{31,32}

These favorable results, in part, explain the increasing utilization of CDA. The increase in utilization demonstrated in the current study is similar to several other studies that analyzed national trends. A 692% and 223% increase in 2-level CDA utilization from 2015 to 2018 in the outpatient and inpatient setting, respectively, were found using the National Surgical Quality Improvement Program database.³³ There was an increase of 150% in CDA utilization from 2006 to 2013 in the NIS database.³⁴ Niedzelak et al found a 20% increase in CDA between 2005 and 2014 using the Medicare PearlDiver database.²¹ Similarly, Jain et al found an increase of 17% per year from 2009 to 2014 in the Humana PearlDiver database.³⁵

Despite the promising outcomes of CDA, the increase in utilization, and an increase in hospital charges for the procedure, we found a decrease in physician Medicare reimbursement. While this is the first study to report on Medicare reimbursement for CDA, recent studies have published on Medicare reimbursement for other common orthopedic procedures. Lopez et al recently published a similar study on ACDF procedures.³⁶ They found an increase in annual procedure volume of 24.2% in the Medicare Part A and B database, while inflation-adjusted physician reimbursement fell 4.3% over the same period. Mayfield et al studied Medicare reimbursement trends for hip and knee arthroplasty and found that inflation-adjusted reimbursement fell by 31.9% and 33.3% for all hip and knee arthroplasty procedures, respectively, from 2000 to 2019.³⁷ They concluded that this trend in reimbursement was not sustainable for physician practice.

The disproportionate increase in hospital charges compared with physician reimbursement may incentivize surgeons to transition to outpatient CDA, as outpatient procedures are more cost-effective and can result in increased reimbursement.³⁸ The safety profile of outpatient CDA has been assessed by several recent studies.^{33,39} A large national database study found that after propensity score matching, there was no difference in the inpatient vs outpatient CDR 30-day complication profile and readmission rate, citing decreasing reimbursement as a driving factor in increasing outpatient CDA.⁴⁰ Cuellar et al found that single- and multilevel CDA in the outpatient

setting were both safe and efficient, although the conclusions did caution that careful patient selection was an important consideration in the low complication risk.⁴¹ As reimbursement decreases and an increased emphasis is placed on cost savings, shifts to outpatient surgery centers may lead to stricter patient selection, which will likely disproportionately affect older patients with increased comorbidities.

Limitations

This study has several limitations. It is a national database study, so it lacks granular patient data and is susceptible to bias and medical coding errors. In addition, we utilized the NIS database for utilization trends and the Centers for Medicare and Medicaid Services database for monetary data. This resulted in 2 different time periods, 2007 to 2017 and 2009 to 2021, used in the study. However, we do not believe this affects our results or conclusions. Finally, the NIS database is an inpatient-only database, so we did not capture outpatient procedures in our utilization trend data.

CONCLUSION

In conclusion, CDA utilization has increased by 183% from 2007 to 2017, and the total hospital charges for the procedure have increased by 22.4% over the same period. Medicare beneficiaries increased utilization by 149%. However, physician Medicare reimbursement has decreased by 12.9% from 2009 to 2021. If this course continues, physicians may be unable to sustain offering CDA to Medicare patients. In addition, patients who meet criteria for outpatient surgery may be more likely to be selected for surgery. This has the danger of creating a health care disparity for Medicare patients as well as for patients with increased comorbidities. Policy makers, hospitals, and physicians should be aware of this to ensure equitable access to health care.

REFERENCES

1. Global Burden of Disease Study 2013 Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2015;386(9995):743–800. doi:10.1016/S0140-6736(15)60692-4

2. Rao RD, Currier BL, Albert TJ, et al. Degenerative cervical spondylosis: clinical syndromes, pathogenesis, and management. *J Bone Joint Surg Am.* 2007;89(6):1360–1378. doi:10.2106/00004623-200706000-00026

3. Smith GW, Robinson RA. The treatment of certain cervicalspine disorders by anterior removal of the intervertebral disc and interbody fusion. *J Bone Joint Surg Am.* 1958;40-A(3):607–624.

4. Buttermann GR. Anterior cervical discectomy and fusion outcomes over 10 years: a prospective study. *Spine (Phila Pa 1976)*. 2018;43(3):207–214. doi:10.1097/BRS.00000000002273

5. Stull JD, Goyal DKC, Mangan JJ, et al. The outcomes of patients with neck pain following ACDF: a comparison of patients with radiculopathy, myelopathy, or mixed symptomatology. *Spine (Phila Pa 1976)*. 2020;45(21):1485–1490. doi:10.1097/ BRS.00000000003613

6. Cho SK, Riew KD. Adjacent segment disease following cervical spine surgery. *J Am Acad Orthop Surg.* 2013;21(1):3–11. doi:10.5435/JAAOS-21-01-3

7. Epstein NE. A review of complication rates for anterior cervical diskectomy and fusion (ACDF). *Surg Neurol Int*. 2019;10:100. doi:10.25259/SNI-191-2019

8. Shriver MF, Lewis DJ, Kshettry VR, Rosenbaum BP, Benzel EC, Mroz TE. Pseudoarthrosis rates in anterior cervical discectomy and fusion: a meta-analysis. *Spine J*. 2015;15(9):2016–2027. doi:10.1016/j.spinee.2015.05.010

9. Crawford CH 3rd, Carreon LY, Mummaneni P, Dryer RF, Glassman SD. Asymptomatic ACDF nonunions underestimate the true prevalence of radiographic pseudarthrosis. *Spine (Phila Pa 1976)*. 2020;45(13):E776–E780. doi:10.1097/BRS.00000000003444

10. Hilibrand AS, Carlson GD, Palumbo MA, Jones PK, Bohlman HH. Radiculopathy and myelopathy at segments adjacent to the site of a previous anterior cervical arthrodesis. *J Bone Joint Surg Am.* 1999;81(4):519–528. doi:10.2106/00004623-199904000-00009

11. Bydon M, Xu R, Macki M, et al. Adjacent segment disease after anterior cervical discectomy and fusion in a large series. *Neurosurgery*. 2014;74(2):139–146. doi:10.1227/ NEU.000000000000204

12. Riew KD, Buchowski JM, Sasso R, Zdeblick T, Metcalf NH, Anderson PA. Cervical disc arthroplasty compared with arthrodesis for the treatment of myelopathy. *J Bone Joint Surg Am*. 2008;90(11):2354–2364. doi:10.2106/JBJS.G.01608

13. Janssen ME, Zigler JE, Spivak JM, Delamarter RB, Darden BV, Kopjar B. ProDisc-C total disc replacement versus anterior cervical discectomy and fusion for single-level symptomatic cervical disc disease: seven-year follow-up of the prospective randomized U.S. food and drug administration investigational device exemption study. *J Bone Joint Surg Am.* 2015;97(21):1738–1747. doi:10.2106/JBJS.N.01186

14. Gornet MF, Burkus JK, Shaffrey ME, Argires PJ, Nian H, Harrell FE Jr. Cervical disc arthroplasty with prestige Lp disc versus anterior cervical discectomy and fusion: a prospective, multicenter investigational device exemption study. *J Neurosurg Spine*. 2015;23(5):558–573. doi:10.3171/2015.1.SPINE14589

15. Gornet MF, Lanman TH, Burkus JK, et al. Two-level cervical disc arthroplasty versus anterior cervical discectomy and fusion: 10-year outcomes of a prospective, randomized investigational device exemption clinical trial. *J Neurosurg Spine*. 2019;31(4):508–518. doi:10.3171/2019.4.SPINE19157

16. Findlay C, Ayis S, Demetriades AK. Total disc replacement versus anterior cervical discectomy and fusion: a systematic review with meta-analysis of data from a total of 3160 patients across 14 randomized controlled trials with both short- and medium- to long-term outcomes. *Bone Joint J*. 2018;100-B(8):991–1001. doi:10.1302/0301-620X.100B8.BJJ-2018-0120.R1

17. Tu T-H, Lee C-Y, Kuo C-H. Cervical disc arthroplasty for less-mobile discs. *J Neurosurg Spine*. 2019;31(3):310–316. doi:10. 3171/2019.2.SPINE181472

18. Hu X, Liu H, Wang B. Cervical disc arthroplasty versus anterior cervical discectomy and fusion for the treatment of single-level disc degenerative disease with preoperative reversible kyphosis. *Clin Neurol Neurosurg.* 2021;202:106493. doi:10.1016/j. clineuro.2021.106493

19. Xu H, Liu H, Hong Y. Clinical and radiological outcomes of single-level cervical disc arthroplasty in the patients with preoperative reversible kyphosis: a matched cohort study. *Clin Neurol Neurosurg*. 2020;198:106247. doi:10.1016/j.clineuro.2020.106247

20. Chang C-C, Huang W-C, Wu J-C, Mummaneni PV. The option of motion preservation in cervical spondylosis: cervical disc arthroplasty update. *Neurospine*. 2018;15(4):296–305. doi:10.14245/ns.1836186.093

21. Niedzielak TR, Ameri BJ, Emerson B, Vakharia RM, Roche MW, Malloy JP. Trends in cervical disc arthroplasty and revisions in the medicare database. *J Spine Surg.* 2018;4(3):522–528. doi:10.21037/jss.2018.09.04

22. HCUP National Inpatient Sample (NIS). *Healthcare Cost and Utilization Project (HCUP)*. 2007-2017. Rockville, MD: Agency for Healthcare Research and Quality; date unknown. www. hcup-us.ahrq.gov/nisoverview.jsp.

23. CPI Inflation Calculator. https://www.bls.gov/data/inflation_calculator.htm. Accessed September 4, 2021.

24. Aaos. *Aaos Musculoskeletal Coding Guide 2020*. American Academy of Orthopaedic Surgeons; 2020.

25. Physician Fee Schedule Look-Up Tool. https://www.cms. gov/Medicare/Medicare-Fee-for-Service-Payment/PFSlookup. Accessed September 4, 2021.

26. CPI Home. 2017. https://www.bls.gov/cpi/. Accessed September 4, 2021.

27. CPI Inflation Calculator. https://www.bls.gov/data/inflation_calculator.htm. Accessed September 4, 2021.

28. Ament JD, Yang Z, Nunley P, Stone MB, Kim KD. Costeffectiveness of cervical total disc replacement vs fusion for the treatment of 2-level symptomatic degenerative disc disease. *JAMA Surg.* 2014;149(12):1231–1239. doi:10.1001/jamasurg.2014.716

29. Kim JS, Dowdell J, Cheung ZB, et al. The seven-year costeffectiveness of anterior cervical discectomy and fusion versus cervical disc arthroplasty: a Markov analysis. *Spine (Phila Pa 1976)*. 2018;43(22):1543–1551. doi:10.1097/BRS.000000000002665

30. Doan MK, Chung AS, Makovicka JL, Hassebrock JD, Polveroni TM, Patel KA. Comparison of two-level cervical disc replacement versus two-level anterior cervical discectomy and fusion in the outpatient setting. *Spine (Phila Pa 1976)*. 2021;46(10):658–664. doi:10.1097/BRS.00000000003871

31. Saifi C, Fein AW, Cazzulino A, et al. Trends in resource utilization and rate of cervical disc arthroplasty and anterior cervical discectomy and fusion throughout the united states from 2006 to 2013. *Spine J.* 2018;18(6):1022–1029. doi:10.1016/j. spinee.2017.10.072

32. Jain NS, Nguyen A, Formanek B, et al. Cervical disc replacement: trends, costs, and complications. *Asian Spine J*. 2020;14(5):647–654. doi:10.31616/asj.2019.0246

33. Gornet MF, Lanman TH, Burkus JK, et al. Cervical disc arthroplasty with the prestige Lp disc versus anterior cervical discectomy and fusion, at 2 levels: results of a prospective, multicenter randomized controlled clinical trial at 24 months. *J Neurosurg Spine*. 2017;26(6):653–667. doi:10.3171/2016.10.SPINE16264

34. Heller JG, Sasso RC, Papadopoulos SM, et al. Comparison of Bryan cervical disc arthroplasty with anterior cervical decompression and fusion: clinical and radiographic results of a randomized, controlled, clinical trial. *Spine (Phila Pa 1976)*. 2009;34(2):101–107. doi:10.1097/BRS.0b013e31818ee263

35. Radcliff K, Davis RJ, Hisey MS, et al. Long-term evaluation of cervical disc arthroplasty with the mobi-C[©] cervical disc: a randomized, prospective, multicenter clinical trial with seven-year follow-up. *Int J Spine Surg.* 2017;11(4):31. doi:10.14444/4031

36. Lopez CD, Boddapati V, Lombardi JM, et al. Recent trends in medicare utilization and reimbursement for anterior cervical discectomy and fusion. *Spine J*. 2020;20(11):1737. doi:10.1016/j. spinee.2020.06.010

37. Mayfield CK, Haglin JM, Levine B, Della Valle C, Lieberman JR, Heckmann N. Medicare reimbursement for hip and knee arthroplasty from 2000 to 2019: an unsustainable trend. *J Arthroplasty*. 2020;35(5):1174–1178. doi:10.1016/j.arth.2019.12.008

38. Munnich EL, Parente ST. Procedures take less time at ambulatory surgery centers, keeping costs down and ability to meet demand up. *Health Aff (Millwood)*. 2014;33(5):764–769. doi:10.1377/hlthaff.2013.1281

39. Wang X, Meng Y, Liu H, et al. Comparison of the safety of outpatient cervical disc replacement with inpatient cervical disc replacement: a systematic review and meta-analysis. *Global Spine J*. 2021;11(7):1121–1133. doi:10.1177/2192568220959265

40. Bovonratwet P, Fu MC, Tyagi V, Ondeck NT, Albert TJ, Grauer JN. Safety of outpatient single-level cervical total disc replacement: a propensity-matched multi-institutional study. *Spine (Phila Pa 1976)*. 2019;44(9):E530–E538. doi:10.1097/BRS.00000000002884

41. Cuéllar JM, Lanman TH, Rasouli A. The safety of single and multilevel cervical total disc replacement in ambulatory surgery centers. *Spine (Phila Pa 1976)*. 2020;45(8):512–521. doi:10.1097/ BRS.00000000003307

Funding: The authors received no financial support for the research, authorship, and/or publication of this article.

Declaration of Conflicting Interests: There are no direct conflicts of interest in this study. Emily S. Mills, Cory K. Mayfield, Tara Shelby, and Andy T. Ton have no conflicts to declare. Raymond J. Hah is a paid consultant for Nuvasive. He is on the editorial boards for Global Spine Journal, Clinical Spine Surgery, and The Spine Journal. Ram K. Alluri is a paid consultant and receives royalties from HIA Technologies Incorporated.

IRB Approval: Institutional review board (IRB) approval and informed consent were not required to complete the study given that all databases used in this study are publicly available national databases without patient identifiers.

Corresponding Author: Emily S. Mills, Keck School of Medicine, Department of Orthopaedic Surgery, University of Southern California, 1200 N. State St, GH 3900, Los Angeles, CA 90033, USA; Emily.mills@med.usc.edu

Published 16 March 2023

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2023 ISASS. To see more or order reprints or permissions, see http:// ijssurgery.com.