

Biportal Endoscopic TLIF With an Expandable Cage: Technical Note and Preliminary Results in Terms of Segmental Lordosis Achievement and Disc Height Elevation

Tae Hoon Kang, Minjoon Cho and Jae Hyup Lee

Int J Spine Surg 2024, 18 (5) 571-581

doi: <https://doi.org/10.14444/8680>

<https://www.ijssurgery.com/content/18/5/571>

This information is current as of May 4, 2025.

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://ijssurgery.com/alerts>

Biportal Endoscopic TLIF With an Expandable Cage: Technical Note and Preliminary Results in Terms of Segmental Lordosis Achievement and Disc Height Elevation

TAE HOON KANG, MD^{1,2}; MINJOON CHO, MD^{1,2}; AND JAE HYUP LEE, MD, PhD^{1,2}

¹Department of Orthopedic Surgery, SMG-SNU BRM Medical Center, Seoul, South Korea; ²Department of Orthopedic Surgery, Seoul National University, College of Medicine, Seoul, South Korea

ABSTRACT

Background: Biportal endoscopic transforaminal lumbar interbody fusion (BE-TLIF) is a minimally invasive TLIF (MIS-TLIF) technique, commonly performed with various cage types. Expandable cages are particularly effective in achieving segmental lordosis (SL) and disc height (DH) elevation in minimally invasive TLIF. However, the published literature lacks details regarding how these outcomes can be accomplished using BE-TLIF with an expandable cage.

Methods: Nine cases (10 levels) of BE-TLIF with an expandable cage were reviewed. Procedures including unilateral laminotomy and bilateral decompression, cage expansion trials, and bilateral facetectomies were carried out under biportal endoscopy to achieve SL and DH elevation. Postoperative standing x-ray images at 3 months and reconstructed computed tomography images were analyzed. The sublamina decompression angle—measured as the angle between the spinous process and the sublamina decompression line on axial computed tomography—was used to evaluate contralateral sublamina decompression.

Results: All procedures were completed without changes to the surgical methods. Eight patients underwent single-level fusion, with 4 of them receiving additional decompression at adjacent levels. One patient underwent a 2-level fusion. Four cases utilized 12° lordotic cages, while the rest employed 20° hyperlordotic cages. The total time for each fusion was 152.5 ± 38.5 minutes. Segmental lordosis increased by 5.1° , with anterior and posterior DH elevations of 4.8 ± 1.7 mm and 3.1 ± 1.8 mm, respectively. No endplate injuries or early cage subsidence occurred. The mean sublamina decompression angle was $31.8^\circ \pm 7.0^\circ$.

Conclusions: BE-TLIF with an expandable cage may offer benefits in SL correction and DH elevation. These advantages are attributed to the use of more lordotic expandable cages, combined with contralateral facetectomies and careful endplate preparation—key features of the BE-TLIF technique.

Clinical Relevance: SL correction and DH elevation can be achieved through BE-TLIF, which helps to reduce the recurrence of symptoms and improves the lumbar lordotic curve.

Level of Evidence: 4.

Endoscopic Minimally Invasive Surgery

Keywords: biportal endoscopic TLIF, expandable cage, biportal endoscopic spine surgery, minimally invasive TLIF, hyperlordotic expandable cages

INTRODUCTION

Lumbar interbody fusion is a surgical technique used to treat lumbar stenosis and instability between vertebral segments when nonsurgical options are ineffective. Achieving adequate segmental lordosis (SL) and disc height (DH) during fusion surgery is critical for successful outcomes.¹ Among the various approaches to accessing the disc space, the posterior techniques, such as posterior lumbar interbody fusion and transforaminal lumbar interbody fusion (TLIF), allow for direct decompression of the central canal.² However, since

the cage is inserted posteriorly, achieving sufficient SL and DH elevation can be challenging.³ As a solution, expandable cages, which can adequately elevate the anterior DH after insertion, have been developed.⁴

Minimally invasive spine surgeries (MISS) have gained popularity due to the reduced invasiveness compared with traditional spine surgeries. Since the early 2000s, minimally invasive TLIF (MIS-TLIF), performed with microscopes and tubular retractors, has become a common practice.^{5,6} To achieve SL and DH elevation with smaller incisions and less muscle

detachment, expandable cages have been introduced in MIS-TLIF procedures.^{7,8} However, there is ongoing debate about the effectiveness of expandable cages in MIS-TLIF. While some studies report improvements in radiographic parameters such as SL and DH,⁷⁻⁹ others have noted concerns about cage subsidence, questioning its long-term effectiveness.¹⁰

Biportal endoscopic spine surgery (BESS), a type of MISS, has gained popularity due to its familiarity for surgeons and ease of use. It offers a similar field of view to a microscope and allows for greater freedom of instrument movement through dual incisions.¹¹ As more surgeons adopt unilateral laminotomy with bilateral decompression (ULBD) using BESS, the biportal endoscopic TLIF (BE-TLIF) has also emerged as a viable technique.¹²⁻¹⁴ BE-TLIF provides a similar surgical view to microscopic MIS-TLIF and allows for meticulous endplate preparation through direct visualization of the intervertebral space. Additionally, it facilitates direct decompression of the bilateral canal space using a 30° scope from a unilateral approach.¹⁵⁻¹⁸

Various cage types have been used during BE-TLIF, and the expandable cage is a promising option. Although some studies have reported on the surgical outcomes of BE-TLIF with expandable cages, no study has thoroughly described the procedure in detail, particularly regarding its potential to achieve SL and DH elevation.^{19,20}

The purpose of this study is to review the procedure of BE-TLIF with an expandable cage at our hospital, analyze the outcomes with a focus on SL and DH elevation, and provide surgical tips to enhance the effectiveness of the procedure.

METHODS

Research Ethics

As this was a retrospective case series study based on the review of patients' medical records after surgery, consent to participate was exempted. The study was approved by the Institutional Review Board (IRB) at our hospital (30-2024-49, SMG-SNU BMC IRB).

Materials

This review included 9 patients (10 levels) who underwent BE-TLIF with expandable cages at our hospital between June 2023 and June 2024. The indications for BE-TLIF with an expandable cage are the same as those for conventional MIS-TLIF. Similarly, the absolute contraindications are identical to those of conventional MIS-TLIF.

Expandable Cage

We used Excender expandable cages (CG Bio, South Korea). These cages have a width of 11 mm and are available in 2 lengths: 28 and 32 mm. The cages offer fixed angles with 2 options: 12° (lordotic) and 20° (hyperlordotic). The initial cage height is either 8 or 10 mm, and it can be expanded by up to 4 mm using a torque driver. A trial implant is available to test the cage's expansion capability. After the final cage insertion, additional bone grafts, such as hydroxyapatites or rhBMP2, can be injected through a posterior injection channel.

Surgical Procedure

The instruments used for BE-TLIF are the same as those employed in conventional BE-TLIF. At our hospital, arthroscopic devices (Arthrex, Florida, USA) and 30° scopes were used, along with radiofrequency ablaters (Delphi, South Korea) for ablation and coagulation. Other instruments were standard for MIS-TLIF procedures.

All surgeries were performed under general anesthesia, with the patient positioned prone on a spine table. Standard skin preparation and draping were carried out. The approach direction was based on the side with the more severe radiating pain, although, due to the benefits of BESS in performing ULBD, a left-sided approach was generally preferred. The surgeon stood on the left side of the patient, while the devices, including arthroscopic tools and the C-arm, were positioned on the right side.

This procedure is a variation of the posterolateral trans-Kambin interlaminar TLIF. Three incisions are made as part of the process (Figure 1).

First, after identifying the midline and disc space via C-arm, 2 longitudinal 1-cm incisions are made lateral to the upper and lower lumbar pedicle margins. These incisions, commonly used for BE-TLIF and percutaneous pedicle screws, are referred to as cranial and caudal biportal incisions.^{11,13,21} Using these 2 incisions, the periosteal muscle detachment of the lamina and ipsilateral facet joint is performed with a periosteal elevator. After serial dilation using a tubular dilator to widen the fascia, a 30° scope is inserted through the cranial biportal incision. Saline, maintained at a pressure of 30 mmHg using a pressure-controlled arthroscopic pump, is used for fluid management.

A high-speed diamond burr is then employed to perform a unilateral laminotomy and resection of the deep portion of the interspinous ligament while preserving the superficial spinous process. Next, contralateral




Figure 1. Incisions for biportal endoscopic transforaminal lumbar interbody fusion with an expandable cage. Red solid lines: 2 longitudinal 1-cm incisions lateral to the pedicle margin for biportal endoscopic procedures and percutaneous pedicle screw placement. These are referred to as the cranial and caudal biportal incisions. Blue arrow line: a less than 1 cm longitudinal incision just lateral to the midline, known as the quarterback incision. Green dotted lines: 2 longitudinal 1-cm incisions lateral to the pedicle margin for contralateral percutaneous pedicle screw insertions.

sublaminar bony resection is performed to achieve bilateral decompression, with complete resection of the ligamentum flavum (ULBD) until both traversing nerve roots are visualized and free. During this process, the 30° scope is rotated toward the contralateral side,

providing a wide view of the epidural space beneath the midline. Once full decompression of the dura is confirmed, bilateral facetectomies are performed with reduced risk of neural tissue damage. This procedure includes total ipsilateral synovectomy, complete resection of the ipsilateral inferior articular process (IAP), partial resection of the ipsilateral superior articular process (SAP), partial contralateral synovectomy, and partial resection of both the contralateral IAP and SAP. The scope is first rotated toward the ipsilateral side to inspect the remaining facet joint after laminotomy. Using tools such as a chisel or osteotome, the remaining IAP and partial SAP are carefully removed to ensure sufficient space for smooth cage insertion without bony obstructions, as excessive resection of the SAP can lead to bleeding. The annulotomy and discectomy are then performed via the caudal biportal incision. The end-plate preparation is continuously monitored through the cranial biportal portal with the endoscope.

If necessary, an additional small incision, called the “quarterback incision,” can be made near the midline. This incision is less than 1 cm in size and allows for the insertion of additional instruments, such as root retractors. It can be used for additional endoscopic view, additional ipsilateral discectomy, dura retraction, and drain insertion (Figures 2 and 3).

The cage trial is then inserted into the intervertebral space. After trial insertion, possible expansion is verified using the C-arm. If the expansion is insufficient, additional contralateral facetectomy is performed until adequate release between the 2 vertebral bodies is confirmed, which is checked again using the cage trial. Following this, the conventional BE-TLIF procedures are completed. Bone grafting is done via a funnel, using a mixture of autologous bone from the facetectomy and other bone substitutes.




Figure 2. (A) Covered field of view of the 30° scope before laminotomy. (B) Covered field of view of the 30° scope after unilateral laminotomy and bilateral decompression. (C) Red arrow indicates additional facetectomy, discectomy, and cage insertion. (D) Blue line marks the quarterback incision, which can be used for the endoscope, additional ipsilateral discectomy, dura retraction, and drain insertion.




Figure 3. Endoscopic view. (A) Decompressed central canal after unilateral laminotomy. (B) Contralateral sublaminar bony resection while preserving the spinous process. (C) Contralateral sublaminar partial facetectomy performed using an osteotome. (D) After ipsilateral total inferior articular process and partial superior articular process resection. (E) Discectomy performed through the caudal biportal incision and quarterback incision. (F) Endoscope insertion into the disc space to inspect endplate preparation via the quarterback incision.

The real expandable cage is inserted via the caudal biportal incision from a lateral-to-medial and posterior-to-anterior direction to ensure it reaches the anterior epiphyseal ring, providing sufficient anterior positioning and epiphyseal ring support. During cage insertion, root retractors can be positioned via the quarterback incision to prevent dura injuries. Once the cage is fully expanded, additional bone grafts, such as rhBMP2, can be delivered through the posterior injection channel. Meticulous hemostasis is achieved, and a surgical drain can be placed via the quarterback incision. Percutaneous pedicle screw insertion and rod fixation are performed through the previous biportal endoscopy incisions and new contralateral percutaneous pedicle fixation incisions (Figure 4).

Outcome Measurements

The patients' demographics (age, sex, height, weight, body mass index, American Society of Anesthesiologists score, presence of osteoporosis, and HbA1c) and clinical factors (preoperative diagnosis, surgical level, preoperative DH, spondylolisthesis slip, presence of kissing spine, and spinopelvic parameters including pelvic incidence [PI], pelvic tilt [PT], and lumbar lordosis [LL]) were reviewed. Additionally, surgical factors such as total operation time, time spent on ULBD and cage insertion, cage size, cage angle, and final cage height were examined. Postoperative radiological

factors were assessed using standing radiography at 3 months postoperatively and computed tomography (CT). These included SL correction, anterior, posterior, and mean DH elevation, slip reduction, final PI-LL, postoperative PT decrease, incidence of endplate injury, and the sublaminar decompression angle. The mean DH was calculated as the average of the anterior and posterior DHs.²² The sublaminar decompression angle is a novel concept that refers to the angle between the spinous process and the decompressed contralateral sublaminar space's bony contour, measured on axial CT (Figure 5).

Statistical Analysis

All data were analyzed using REX version 1.0 (Rex-software, Korea).

RESULTS

All surgeries were completed without deviations from the surgical plan. The diagnoses included 7 levels of degenerative spondylolisthesis with severe spinal stenosis (Schizas grade D) and 3 levels of foraminal stenosis that had failed conservative treatment. Two of the foraminal stenosis cases were revision surgeries. Patients' demographics and clinical factors are summarized in Table 1.




Figure 4. (A and B) C-arm images showing the cage expansion trial. (C and D) C-arm images of the real cage insertion and expansion. (E) Real cage insertion with dura protection using a root retractor via the quarterback incision. (F) Real cage expansion using a torque driver.

One patient underwent a 2-level BE-TLIF with expandable cages, while 4 patients had additional decompression at adjacent levels. The average total surgery time was 286 minutes, with an average of 153 minutes from ULBD to cage insertion. All cases included a contralateral partial facetectomy and a cage expansion trial.

Postoperative standing x-ray images showed a mean segmental lordotic gain of 5.4° and a mean DH increase of 3.5 mm, with an anterior DH increase of 4.8 mm and posterior height increase of 3.1 mm. Consequently, lumbar lordosis improved by 3.0° . The mean final PI-LL mismatch was 1.2° , and the PT decreased by 2.2° . Two patients did not achieve a PI-LL mismatch of less than




Figure 5. (A) The sublaminar decompression angle: the angle between the spinous process and the bony contour of the decompressed contralateral sublaminar space. (B) Axial computed tomography (CT) image of biportal endoscopic transforaminal lumbar interbody fusion (TLIF) at L4–L5 with an expandable cage. The sublaminar decompression angle is 36°. (C) Axial CT image of tubular minimally invasive TLIF at L4–L5 with an expandable cage. Only ipsilateral facetectomy was performed, and the sublaminar decompression angle is 8°.

10 degrees, as the procedures involved primarily 1-level fusions.

Postoperative CT images revealed no iatrogenic endplate injuries. The sublaminar decompression angle averaged 31.6°, providing sufficient bilateral decompression without resecting the spinous process or requiring a contralateral approach (Table 2).

For example, the patient in case 2 underwent BE-TLIF L4 to L5 with expandable cage and BE decompression L3 to L4. The SL correction was 5.5°. The postoperative CT image of this case is shown in Figure 5. Postoperative 6-week magnetic resonance images showed a fully decompressed central canal (Figure 6).

DISCUSSION

This study presents the technique of BE-TLIF using a lordotic expandable cage and evaluates the outcomes in terms of SL achievement and DH elevation. Concerns have been raised that MIS-TLIF with expandable cages may achieve greater SL and DH elevation compared with static cages.^{23,24} However, our study, though small, demonstrated an average SL correction of 5.1° and a DH elevation of 3.5 mm.

The SL correction observed in this study is comparable to the 5.04° reported in a meta-analysis of MIS-TLIF using expandable cages.²⁵ However, the DH elevation in our study (3.5 mm) exceeds the 1.14 mm reported in that same meta-analysis.²⁵ When considering anterior and posterior DH separately, our results align with findings from other studies.⁹ These data suggest that BE-TLIF with an expandable cage can be a favorable option for achieving SL restoration and DH elevation.

Previous studies have highlighted some limitations of MIS-TLIF with expandable cages. One of the key limitations is that bilateral annulotomies and anterior longitudinal ligament resection are

more critical for SL correction.²⁶ Another issue is that cage expansion without sufficient release of the intervertebral space can lead to cage subsidence and loss of SL and DH.^{27,28} While our procedure shares the first limitation as a posterior approach, the outcomes differ from previous MIS-TLIF procedures for a few reasons.

Key Factors for SL Achievement and DH Elevation in BE-TLIF With Expandable Cages

Lordotic Cages

In our study, we used lordotic cages with angles of 12° and 20°. Past studies have suggested that the degree of cage lordosis was not critical to SL, but those studies involved cages with less than 12° of lordosis.^{29,30} Recent findings indicate that hyperlordotic expandable cages (12+ degrees) should be used to achieve SL correction and DH elevation.^{9,24,30} These cages, with more aggressive angles, are relatively difficult to be inserted due to its initial height in conventional MIS-TLIF. However, their use in BE-TLIF is further facilitated by the nature of the procedure, including better endplate preparation and bilateral partial facetectomies, which are discussed below.

Endplate Preparation

During BE-TLIF, endoscopic visualization allows for thorough cartilaginous endplate curettage, which reduces the risk of acute endplate injury, a key factor in preventing cage subsidence.³¹ Cage expansion using a torque-limited driver, combined with proper endplate preparation, ensures maintenance of SL correction and DH elevation.

Table 1. The preoperative demographic and radiographic data.

Case	Level	Age, y	Sex	Height, cm	Weight, kg	BMI	ASA	Osteoporosis	HbA1c	Preoperative Diagnosis	Revision	Disc Height, mm	Lithesis Grade, mm	Kissing Spine	PI, °	PT, °	LL, °	PI-LL Mismatch, °
1	1	68	F	151.3	55.60	24.29	2	O	6.70	Degenerative spondylolisthesis L4–L5 with spinal stenosis (schiza D)	X	6.5	1 (6)	O	39.8	13.8	52.0	–12.2
2	2	81	M	161	65.00	25.10	2	O	5.90	Degenerative spondylolisthesis L4–L5 with spinal stenosis (schiza D)	X	10.9	1 (5)	O	47.2	15.0	50.0	–2.8
3	3	72	M	164.6	64.25	24.40	2	X	6.50	Degenerative spondylolisthesis L4–L5 with spinal stenosis (schiza D)	X	10.9	1 (5)	O	43.6	11.5	54.0	–10.4
4	4	70	M	168.9	83.70	29.30	2	X	6.80	Foraminal stenosis, L5–S1	X	12.4	0	O	47.1	24.0	44.6	2.5
5	5	79	F	147.3	41.80	19.27	2	O	6.20	Degenerative spondylolisthesis L3–L4 with spinal stenosis (schiza D)	X	10.1	1 (3)	O	52.3	23.8	54.2	–1.9
6	6	65	F	157	49.10	19.92	1	X	5.50	Foraminal stenosis, L4–L5	O	2.7	0	O	38.4	12.3	31.9	6.5
7	7	69	F	148	51.70	23.60	2	X	5.20	Foraminal stenosis, L5–S1	O	12.0	0	O	44.1	28.3	20.1	24.0
8	8	78	F	147.8	43.80	20.05	2	O	5.80	Degenerative spondylolisthesis L3–L4 with spinal stenosis (schiza D)	X	9.6	1 (2)	O	57.7	26.9	36.1	21.6
9	9	72	F	153	68.7	29.35	2	X	6.20	Degenerative spondylolisthesis with L2–L3 spinal stenosis (schiza D)	X	7.9	1 (5)	O	35.3	12.7	29.2	6.1
10										Degenerative spondylolisthesis with L3–L4 spinal stenosis (schiza D)		11.1	1 (7)	O				6.1
Total ^a		72.6 ± 5.2	M (3)/F (6)	155.2 ± 7.5	59.2 ± 13.1	24.5 ± 3.9	1 (1)/2 (8)	O (4)/X (5)	6.1 ± 0.5	Spondylolisthesis (7)/foraminal stenosis (3)	O (2)/X (7)	9.4 ± 3.0		O (10)	44.1 ± 12.4	18.1 ± 6.8	40.1 ± 12.4	4.0 ± 11.9

Abbreviations: ASA, American Society of Anesthesiologists; BMI, body mass index; HbA1c, hemoglobin A1c; LL, lumbar lordosis; PI, pelvic incidence; PT, pelvic tilt.

Note: O indicates present; X indicates absent.

^aData presented as mean ± SD or (n).

Table 2. The operation related data and postoperative radiographic data.

Case	Level	Fusion Level	Adjacent Decompression Operation	Total Operation Time, min	Total Time for ULBD and Cage Insertion, min	Cage Size, mm	Cage Angle, °	Segmental Lordosis Correction, °	LL Increase, °	Anterior Disc Height Elevation, mm	Posterior Disc Height Elevation, mm	Disc Height Elevation, mm	Slip Reduction, mm	Final PI-LL, °	PT Change, °	Sublaminar Decompression Angle, °	Endplate Injury
1	1	L4-L5	X	295	192	32	12	1.0	1.0	5.30	6.30	4.40	5.2	-13.2	-2.0	31	X
2	2	L4-L5	O (L3-L4)	287	165	28	12	5.5	4.8	4.10	1.60	3.20	4.0	-7.6	1.1	36	X
3	3	L4-L5	X	213	162	28	20	1.0	5.3	4.10	1.40	2.60	1.5	-15.7	0.4	36	X
4	4	L5-S1	X	282	209	28	20	-1.0	0.1	4.30	2.80	1.00	0	2.4	-5.1	14	X
5	5	L3-L4	O (L2-L3)	235	128	28	12	6.0	2.5	5.40	1.80	2.90	1.7	-4.4	1.4	33	X
6	6	L4-L5	O (L3-L4)	260	152	28	12	8.0	3.6	9.40	5.20	8.00	0	2.9	-2.8	32	X
7	7	L5-S1	O (L3-L4)	318	186	28	20	9.3	1.5	3.60	2.80	3.20	0	22.5	-5.1	38	X
8	8	L3-L4	X	275	136	28	20	4.0	0.3	3.50	1.00	1.70	4.0	21.3	-1.4	38	X
9	9	L2-L3	X	348	89	28	20	3.2	5.4	3.80	4.30	4.30	3.4	0.7	-2.0	32	X
10	10	L3-L4	X		106	28	20	14.0		4.60	3.50	3.40	3.7			28	X
Total		L2-L3 (1) L3-L4 (3) L4-L5 (4) L5-S1 (2)	O (4)/X (6)	286.1 ± 44.1	152.5 ± 38.2	28 (9)/32 (1)	12 (4), 20 (6)	5.1 ± 4.5	3.0 ± 2.2	4.8 ± 1.7	3.1 ± 1.8	3.5 ± 1.5	2.9 ± 2.3	1.0 ± 12.8	-1.8 ± 2.3	31.8 ± 7.0	X (10)

Abbreviations: LL, lumbar lordosis; PI, pelvic incidence; PT, pelvic tilt; ULBD, unilateral laminotomy with bilateral decompression.

Note: O indicates present, X indicates absent.




Figure 6. (A and B) Preoperative and postoperative 3-month lumbar standing lateral x-ray images. (C and D) Preoperative and postoperative 3-month lumbar standing anteroposterior x-ray images. (E and F) Preoperative and postoperative 6 weeks magnetic resonance images (MRIs), sagittal cut. (G and H) Preoperative and postoperative 6 weeks MRIs, axial cut of endoscopic fusion level, L4–L5.

Bilateral Facetectomies

Bilateral partial facetectomies in BE-TLIF, as demonstrated by the sublaminar decompression angles, facilitate the resection of the contralateral facet joint and partial removal of the SAP. This is made more effective with the use of a 30° endoscope and helps avoid resecting the spinous process. While contralateral facetectomies can be performed in MIS-TLIF using tubular retractors, they are more challenging due to limitations posed by the spinous process, often requiring additional incisions.^{32–34} A cadaver study has shown that adequate decompression of the posterior complex, including bilateral facetectomies, is important for achieving SL.³⁵ Furthermore, bilateral facetectomies can aid in inserting hyperlordotic cages and avoiding endplate breakages by providing better release of the posterior disc space.²⁷ Conversely, relying on unilateral facetectomy alone carries the risk of endplate breakage during cage expansion.²⁸

Lumbar Lordosis and PI-LL Mismatch

Our study showed an increase in lumbar lordosis of 3.0°, and 2 patients failed to achieve a PI-LL mismatch correction of less than 10°, similar to other studies involving hyperlordotic cages.^{9,28,36} Most of the procedures were single-level fusions, and we performed decompression only at the levels of spinal stenosis due to insurance coverage limitations.

Surgical Approach and Incisions

We sometimes used the quarterback incisions for several reasons. The midline incisions allow for versatility, such as performing adjacent level decompressions, inserting the endoscope for cage insertion, and preventing dura injury during cage placement by allowing root retractors. After cage insertion, a temporary drain can be placed via the midline incisions to prevent

hematoma formation. The quarterback incisions are small, and it does not cause muscle belly injury because it is near the spinous process.

Sublaminar Decompression Angle

The concept of the sublaminar decompression angle, typically over 30° in most cases, was introduced to assess the effectiveness of ULBD. This angle represents that the BE-TLIF facilitates bilateral synovectomies and resections of both side IAP and SAP, enabling full contralateral decompression and foraminotomy, which may help prevent complications such as contralateral radiculopathy infrequently seen in MIS-TLIF procedures.^{37,38} In our study, no cases of contralateral radiculopathy occurred, likely due to thorough decompression and facetectomies.

Limitations

A significant limitation of this technique is the longer surgical time, which is a critical factor in MISS. Shorter surgical times reduce the burden on elderly patients and those with comorbidities.³⁹ Moreover, further reductions in surgical time may enable awake spine fusion surgeries performed with regional anesthesia.^{13,40,41} As the author gains more experience with BE-TLIF, surgical times may decrease, potentially enabling awake spine fusion techniques.

This study itself has several limitations. First, it is a technical note with preliminary results on SL achievement and DH elevation, with only a 3-month follow-up. Long-term follow-up is needed to assess potential complications, such as late cage subsidence or low fusion rates. Additionally, the small number of cases limits the generalizability of the results. Finally, the study does not prove that BE-TLIF can achieve better SL correction than tubular MIS-TLIF, nor does it suggest

superiority over other fusion techniques. A comparative study between BE-TLIF and tubular MIS-TLIF is planned.

CONCLUSION

BE-TLIF with an expandable cage may offer benefits in SL correction and DH elevation. These advantages are attributed to the use of hyperlordotic expandable cages, combined with contralateral facetectomies and careful endplate preparation—key features of the BE-TLIF technique.

REFERENCES

1. Lee JH, Lee J-H, Yoon K-S, Kang S-B, Jo CH. Effect of intraoperative position used in posterior lumbar interbody fusion on the maintenance of lumbar lordosis. *SPI*. 2008;8(3):263–270. doi:10.3171/SPI/2008/8/3/263
2. Mobbs RJ, Phan K, Malham G, Seex K, Rao PJ. Lumbar interbody fusion: techniques, indications and comparison of interbody fusion options including PLIF. *LLIF ALIF J Spine Surg Dec*. 2015;1(1):2–18. doi:10.3978/j.issn.2414-469X.2015.10.05
3. Carlson BB, Saville P, Dowdell J, et al. Restoration of lumbar lordosis after minimally invasive transforaminal lumbar interbody fusion: a systematic review. *Spine J*. 2019;19(5):951–958. doi:10.1016/j.spinee.2018.10.017
4. Chang SY, Kang D-H, Cho SK. Innovative developments in lumbar interbody cage materials and design: a comprehensive narrative review. *Asian Spine J*. 2024;18(3):444–457. doi:10.31616/asj.2023.0407
5. Phan K, Rao PJ, Kam AC, Mobbs RJ. Minimally invasive versus open transforaminal lumbar interbody fusion for treatment of degenerative lumbar disease: systematic review and meta-analysis. *Eur Spine J*. 2015;24(5):1017–1030. doi:10.1007/s00586-015-3903-4
6. Foley KT, Lefkowitz MA. Advances in minimally invasive spine surgery. *Clin Neurosurg*. 2002;49:499–517.
7. Hawasli AH, Khalifeh JM, Chatrath A, Yarbrough CK, Ray WZ. Minimally invasive transforaminal lumbar interbody fusion with expandable versus static interbody devices: radiographic assessment of sagittal segmental and pelvic parameters. *Neurosurg Focus*. 2017;43(2):E10. doi:10.3171/2017.5.FOCUS17197
8. Ledesma JA, Lambrechts MJ, Dees A. Static versus expandable interbody fusion devices: a comparison of 1-year clinical and radiographic outcomes in minimally invasive transforaminal lumbar interbody fusion. *Asian Spine J*. 2023;17(1):61–74. doi:10.31616/asj.2021.0486
9. Jitpakdee K, Sommer F, Gouveia E. Expandable cages that expand both height and lordosis provide improved immediate effect on sagittal alignment and short-term clinical outcomes following minimally invasive transforaminal lumbar interbody fusion (MIS TLIF). *J Spine Surg*. 2024;10(1):55–67. doi:10.21037/jss-23-106
10. Subramanian T, Merrill RK, Shahi P, et al. Predictors of subsidence and its clinical impact after expandable cage insertion in minimally invasive transforaminal interbody fusion. *Spine*. 2023;48(23):1670–1678. doi:10.1097/BRS.0000000000004619
11. Kim J-E, Choi D-J, Park EJJ, et al. Biportal endoscopic spinal surgery for lumbar spinal stenosis. *Asian Spine J*. 2019;13(2):334–342. doi:10.31616/asj.2018.0210
12. Park S-M, Park J, Jang HS, et al. Biportal endoscopic versus microscopic lumbar decompressive laminectomy in patients with spinal stenosis: a randomized controlled trial. *Spine J*. 2020;20(2):156–165. doi:10.1016/j.spinee.2019.09.015
13. Kang TH, Kim WJ, Lee JH. Efficacy of the erector spinae plane block with sedation for unilateral biportal endoscopic spine surgery and comparison with other anesthetic methods. *Acta Neurochir*. 2023;165(9):2651–2663. doi:10.1007/s00701-023-05643-1
14. Park S-M, Lee H-J, Park H-J, et al. Biportal endoscopic versus microscopic discectomy for lumbar herniated disc: a randomized controlled trial. *Spine J*. 2023;23(1):18–26. doi:10.1016/j.spinee.2022.09.003
15. Heo DH, Hong YH, Lee DC, Chung HJ, Park CK. Technique of biportal endoscopic transforaminal lumbar interbody fusion. *Neurospine*. 2020;17(Suppl 1):S129–S137. doi:10.14245/ns.2040178.089
16. Kang MS, You KH, Choi JY, Heo DH, Chung HJ, Park HJ. Minimally invasive transforaminal lumbar interbody fusion using the biportal endoscopic techniques versus microscopic tubular technique. *Spine J*. 2021;21(12):2066–2077. doi:10.1016/j.spinee.2021.06.013
17. Liang J, Lian L, Liang S, et al. Efficacy and complications of unilateral biportal endoscopic spinal surgery for lumbar spinal stenosis: a meta-analysis and systematic review. *World Neurosurg*. 2022;159:e91–e102. doi:10.1016/j.wneu.2021.12.005
18. Lin GX, Yao ZK, Zhang X, Chen CM, Rui G, Hu BS. Evaluation of the outcomes of biportal endoscopic lumbar interbody fusion compared with conventional fusion operations: a systematic review and meta-analysis. *World Neurosurg*. 2022;160:55–66. doi:10.1016/j.wneu.2022.01.071
19. Park DY, Heo DH. The use of dual direction expandable titanium cage with biportal endoscopic transforaminal lumbar interbody fusion: a technical consideration with preliminary results. *Neurospine*. 2023;20(1):110–118. doi:10.14245/ns.2346116.058
20. Cao S, Fan B, Song X, Wang Y, Yin W. Expandable versus static cages in unilateral biportal endoscopy lumbar interbody fusion (ULIF) for treating degenerative lumbar spondylolisthesis (DLS): comparison of clinical and radiological results. *J Orthop Surg Res*. 2023;18(1):505. doi:10.1186/s13018-023-03979-z
21. Kang MS, Chung HJ, Jung HJ, Park HJ. How I do it? extraforaminal lumbar interbody fusion assisted with biportal endoscopic technique. *Acta Neurochir*. 2021;163(1):295–299. doi:10.1007/s00701-020-04435-1
22. Iii WS, Orías AAE, Shifflett GD. Image-based markers predict dynamic instability in lumbar degenerative spondylolisthesis. *Neurospine*. 2020;17(1):221–227. doi:10.14245/ns.1938440.220
23. Canseco JA, Karamian BA, DiMaria SL, et al. Static versus expandable polyether ether ketone (PEEK) interbody cages: a comparison of one-year clinical and radiographic outcomes for one-level transforaminal lumbar interbody fusion. *World Neurosurg*. 2021;152:e492–e501. doi:10.1016/j.wneu.2021.05.128
24. Yee TJ, Joseph JR, Terman SW, Park P. Expandable vs static cages in transforaminal lumbar interbody fusion: radiographic comparison of segmental and lumbar sagittal angles. *Neurosurgery*. 2017;81(1):69–74. doi:10.1093/neuros/nyw177
25. Alvi MA, Kurian SJ, Wahood W, Goyal A, Elder BD, Bydon M. Assessing the difference in clinical and radiologic outcomes between expandable cage and nonexpandable cage among

patients undergoing minimally invasive transforaminal interbody fusion: a systematic review and meta-analysis. *World Neurosurg.* 2019;127:596–606. doi:10.1016/j.wneu.2019.03.284

26. Macki M, Hamilton T, Haddad YW, Chang V. Expandable cage technology-transforaminal, anterior, and lateral lumbar interbody fusion. *Oper Neurosurg.* 2021;21(Suppl 1):S69–S80. doi:10.1093/ons/opaa342

27. Chang C-C, Chou D, Pennicooke B, et al. Long-term radiographic outcomes of expandable versus static cages in transforaminal lumbar interbody fusion. *J Neurosurg.* 2021;34(3):471–480. doi:10.3171/2020.6.SPINE191378

28. Armocida D, Pesce A, Cimatti M, Proietti L, Santoro A, Frati A. Minimally invasive transforaminal lumbar interbody fusion using expandable cages: increased risk of late postoperative subsidence without a real improvement of perioperative outcomes: a clinical monocentric study. *World Neurosurg.* 2021;156:e57–e63. doi:10.1016/j.wneu.2021.08.127

29. Cho JH, Hwang CJ, Lee DH, Lee CS. Using lordotic cages at the L5-S1 level does not guarantee the improvement of sagittal alignment in patients who underwent posterior lumbar interbody fusion. *Asian Spine J.* 2023;17(3):477–484. doi:10.31616/asj.2022.0228

30. Lee JH, Lee DO, Lee JH, Shim HJ. Effects of lordotic angle of a cage on sagittal alignment and clinical outcome in one level posterior lumbar interbody fusion with pedicle screw fixation. *Biomed Res Int.* 2015;2015:523728. doi:10.1155/2015/523728

31. Kim CW, Doerr TM, Luna IY, et al. Minimally invasive transforaminal lumbar interbody fusion using expandable technology: a clinical and radiographic analysis of 50 patients. *World Neurosurg.* 2016;90:228–235. doi:10.1016/j.wneu.2016.02.075

32. Chang C-C, Chou D, Pennicooke B, et al. Long-term radiographic outcomes of expandable versus static cages in transforaminal lumbar interbody fusion. *J Neurosurg.* 2020;34(3):471–480. doi:10.3171/2020.6.SPINE191378

33. Hyun SJ, Kim YB, Kim YS, et al. Postoperative changes in paraspinal muscle volume: comparison between paramedian interfascial and midline approaches for lumbar fusion. *J Korean Med Sci.* 2007;22(4):646–651. doi:10.3346/jkms.2007.22.4.646

34. Ge DH, Stekas ND, Varlotta CG, et al. Comparative analysis of two transforaminal lumbar interbody fusion techniques. *Spine.* 2019;44(9):E555–E560. doi:10.1097/BRS.0000000000002903

35. Ochtman AEA, Bisschop A, Bleys R, Öner FC, van Gaalen SM. Surgical techniques in restoration lumbar lordosis: a biomechanical human cadaveric study. *Spine Deform.* 2023;11(1):35–40. doi:10.1007/s43390-022-00549-x

36. Mathew J, Cerpa M, Lee NJ, et al. Comparing hyperlordotic and standard lordotic cages for achieving segmental lumbar lordosis during transforaminal lumbar interbody fusion in adult spinal deformity surgery. *J Spine Surg.* 2021;7(3):318–325. doi:10.21037/jss-21-15

37. Akbary K, Kim J-S, Park CW, Jun SG, Hwang JH. Biportal endoscopic decompression of exiting and traversing nerve roots through a single interlaminar window using a contralateral approach: technical feasibilities and morphometric changes of the lumbar canal and foramen. *World Neurosurg.* 2018;117:153–161. doi:10.1016/j.wneu.2018.05.111

38. Chen Y-L, Hu X-D, Wang Y, Jiang W-Y, Ma W-H. Contralateral radiculopathy after unilateral transforaminal lumbar interbody fusion: causes and prevention. *J Int Med Res.* 2021;49(8). doi:10.1177/03000605211037475

39. Cheng H, Clymer JW, Po-Han Chen B, et al. Prolonged operative duration is associated with complications: a systematic review and meta-analysis. *J Surg Res.* 2018;229:134–144. doi:10.1016/j.jss.2018.03.022

40. Lee JK, Park JH, Hyun SJ, Hodel D, Hausmann ON. Regional anesthesia for lumbar spine surgery: can it be a standard in the future? *Neurospine.* 2021;18(4):733–740. doi:10.14245/ns.2142584.292

41. Kang TH, Kim WJ. Awake unilateral biportal endoscopic thoracic decompression - technical note -. *J Korean Soc Spine Surg.* 2022;29(4):143. doi:10.4184/jkss.2022.29.4.143

Funding: This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT; No. 2022R1A2C1010962).

Declaration of Conflicting Interests: The authors report no conflicts of interest in this work.

Research Ethics: As this was a retrospective case-control study reviewing patients' medical records after surgeries, consents to participate in this study were exempted. This study was reviewed by the IRB in our hospital (30-2024-49, SMG-SNU BMC IRB).

Corresponding Author: Jae Hyup Lee, Professor, Department of Orthopedic Surgery, Seoul National University College of Medicine, President, SMG-SNU Boramae Medical Center 20 Boramae-ro 5-gil, Dongjak-gu, Seoul, Republic of Korea; spinelee@snu.ac.kr

Published 05 November 2024

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2024 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.