

The Role of Psychosocial Screening in Patient Selection for Spine Surgery: A Review

Ethan R. Faye, Jacob R. Ball, Emily S. Mills, Jeffrey Wang, Raymond J. Hah and Ram K. Alluri

Int J Spine Surg published online 8 March 2023
<http://ijssurgery.com/content/early/2023/03/08/8429>

This information is current as of March 27, 2023.

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://ijssurgery.com/alerts>

The Role of Psychosocial Screening in Patient Selection for Spine Surgery: A Review

ETHAN R. FAYE, BA¹; JACOB R. BALL, MD¹; EMILY S. MILLS, MD¹; JEFFREY WANG, MD¹; RAYMOND J. HAH, MD¹; AND RAM K. ALLURI, MD¹

¹Department of Orthopaedic Surgery, University of Southern California Keck School of Medicine, Los Angeles, CA, USA

ABSTRACT

Background: The purpose of this review is to provide a brief history of previous attempts at developing patient screening tools and to further examine the definitions of these psychological concepts, relevance to clinical outcomes, and implications for spine surgeons during preoperative patient assessments.

Methods: A literature review was performed by 2 independent researchers to identify original manuscripts related to spine surgery and novel psychological concepts. The history of presurgical psychology screening was also studied, and definitions of frequently utilized metrics were detailed.

Results: Seven manuscripts were identified that utilized psychological metrics for preoperative risk assessments and correlated outcomes with these scores. The metrics most frequently used in the literature included resilience, patient activation, grit, and self-efficacy.

Discussion: Current literature favors resilience and patient activation as important metrics for preoperative patient screening. Available studies demonstrate significant associations between these character traits and patient outcomes. Further research is warranted to investigate the roles of preoperative psychological screening to optimize patient selection in spine surgery.

Clinical Relevance: The purpose of this review is to provide clinicians with a reference for available psychosocial screening tools and their relevance to patient selection. This review also serves to guide future research directions given the importance of this topic.

Level of Evidence: 4.

Other and Special Categories

Keywords: minimally invasive surgery, spine surgery, psychology, grit, resilience, patient activation, psychosocial screening

INTRODUCTION

One of the most important challenges that spine surgeons encounter is the appropriate selection of surgical patients. Research efforts have been directed toward identifying psychological factors that can predict patient outcomes following surgery. Early attempts employed full psychological screening profiles meant to identify high-risk personality traits for patients with organic vs functional low back pain.¹⁻⁴ Hanvik was the first to utilize the Minnesota Multiphasic Personality Inventory (MMPI) and identified hypochondriasis and hysteria as personality traits frequently observed in patients with functional low back pain.¹ These personality traits combined to form a conversion-V personality type and became an area of focus until challenged by studies demonstrating significant obstacles with utilizing this tool to predict outcomes.^{3,4} A more nuanced approach to predicting outcomes arose following the development of the biopsychosocial theory of pain.^{5,6} In addition to underlying pathology, it is well studied that worker's compensation,^{7,8} working status at the time of surgery,⁹

social support,¹⁰ education level,¹¹ and other social factors are predictive of surgical outcomes. Outcomes of interest are typically patient-focused, such as pain, disability, and work capacity.⁵

Significant efforts continue to be underway to develop a preoperative screening tool that can predict which patients are most likely to succeed following spine surgery, but there remains no widely accepted measure.^{12,12-16} A promising new area of research focuses on resilience and grit of orthopedic surgical candidates.¹⁷ Interestingly, studies have demonstrated that these traits can be improved through guided therapy, which could allow physicians to intervene preoperatively if they are found to improve surgical outcomes, such as adherence to physical therapy and health-related quality of life.¹⁸ The purpose of this review is to provide a brief history of previous attempts at developing patient screening tools and to further examine the definitions of these psychological concepts, relevance to clinical outcomes, and implications for spine surgeons during preoperative patient assessments.

PREVIOUS EFFORTS IN PATIENT SELECTION IN SPINE SURGERY

The intersection of psychology, societal factors, and underlying pathology presents a unique challenge for clinicians in determining whether a patient is a good surgical candidate. Given this complexity, there has been an attempt to develop presurgical psychological screening (PPS) score cards that can simplify the process of patient selection for the physician.¹⁹ Epker and Block's review of screening tools details the importance of personality/emotional, cognitive/behavioral, and environmental/historical factors that must be accounted for in any clinically meaningful screening tool.¹⁹

The early screening tool developed by Finneson and Cooper specifically for lumbar discectomy made use of biopsychosocial-type questions to determine which patients may benefit the most from surgery.¹⁶ In addition to physical examination and radiographic findings, their survey asked whether patients had "poor psychological backgrounds," "secondary gain," and a "history of previous lawsuits."¹⁶ Interestingly, the authors report that their score card could predict which patients would benefit most from surgery in the near term but failed to be predictive 5 years postoperatively.¹⁶ Spengler et al also attempted to simplify patient selection for lumbar discectomy by utilizing neurological signs, sciatic tension signs, MMPI scores, and lumbar myelography or computed tomography data.²⁰ After applying their scorecard to 84 patients, they determined that they could predict 40% of outcome variability, and 26% was due to psychological profile, leaving them to conclude that patients with scores less than 50/100 should not be considered for surgery.²⁰

More recent attempts at creating a PPS include a scorecard created by Block et al in 2001, which utilized semistructured interviews.¹³ Questions included medical risk factors as well as psychological risk factors, including family reinforcement of pain, marital satisfaction, coping mechanisms, and MMPI scores. Furthermore, social risk factors such as pending litigation, job satisfaction, and substance abuse were included. Interestingly, their score card was 82.8% accurate in predicting good outcomes and 83% accurate in predicting poor outcomes.¹³ After performing post hoc analysis, they determined that psychological testing alone could predict 78.4% of patient outcomes, with the psychological interview only adding an additional 5% and medical risk factors contributing 1% toward prognosis.¹³

There continues to be significant research toward developing a clinically relevant tool for identifying

which patients are likely to benefit most from undergoing spine surgery.¹⁵ The PPS discussed previously is not routinely used in clinical practice mainly due to the challenges associated with conducting psychological interviews and uncertainty of long-term predictive abilities of these metrics. Although PPS seems promising, long-term follow-up with objective functional outcome measures is required before routinely implementing such a tool into clinical practice.¹³ With further investigation of novel parameters, such as grit and resilience, PPS can continue to be refined for clinical use.

PSYCHOSOCIAL METRICS

Psychological concepts such as resilience, self-efficacy, and grit are distinct and complex with many social, cultural, and biological factors interacting to determine an individual's response to a stressful experience such as surgery.^{21,22} The defined personality trait of grit is somewhat new,²² and there has been some variability in the reporting of resilience in the literature, with no universally agreed upon definition.^{23,24} As such, it is important to examine the most up-to-date definitions and use of these concepts.

DEFINITIONS

Resilience

The American Psychological Association defines resilience as "the process and outcome of successfully adapting to difficult or challenging life experiences, especially through mental, emotional, and behavioral flexibility and adjustment to external and internal demands."²⁵

Additionally, a recent literature review proposed an expansion of the definition to include "the ability to maintain the persistence of one's orientation toward existential purposes."²³ Resilience is beneficial in the face of major life difficulties and is associated with positive interpersonal relationships and social competence²⁶ as well as lower overall health care utilization and improved self-rated health.²⁷ Resilience is common in the general population and allows individuals to continue functioning in other areas of their life despite hindrance from a major stressor.²⁸ Additionally, resilience has been shown to be a dynamic process that is capable of increasing or decreasing in response to a stressful situation.²⁹ Resilience may be conditioned through therapy,³⁰ and targeted interventions are capable of building resilience in pain management,³¹ although the source of resilience may be changed over time and situation.^{26,32} Some of the confusion regarding the definition of resilience may result from the interplay between resilience and other psychological factors. For example, positive mental health

has been associated with high resilience in older adults,³³ and high self-efficacy is correlated with increased resilience behaviors in victims of spinal cord injury.³⁴

Patient Activation

Related to the psychosocial construct of resilience is the model of patient activation, which was created by Hibbard et al in recognition of the vital role patient's play in managing their health. Patient activation comprises 4 domains: (1) believing the patient's role is important, (2) having the confidence and knowledge necessary to take action, (3) actually taking action to maintain and improve one's health, and (4) staying the course even under stress.³⁵ Patient activation has been recognized as a potent indicator of health outcomes in chronic disease and has specific potential in spinal surgery.³⁶

Grit

Grit is a personality trait defined as an individual's perseverance and passion for achieving long-term goals in spite of failures and plateaus of progress.³⁷ It was developed and continues to be studied as a measured trait to account for the difference in success among high-achieving individuals independent of talent and intelligence.^{22,38} Grit has been associated with individuals who seek happiness from engagement and meaning rather than the pursuit of pleasure.³⁹ However, the validity of grit as an independent trait has been called into question, with recent research suggesting that the perseverance aspect of grit plays a far greater role in predicting success than consistency or passion⁴⁰ and that there is construct overlap with the trait of self-control.⁴¹

Self-Efficacy

Self-efficacy "is an individual's subjective perception of his or her capability to perform in a given setting or to attain desired results."⁴² Bandura originally described self-efficacy as the belief that one carries out specific behaviors under specific circumstances.⁴³ Self-efficacy may be able to influence behaviors, activity level, and achievements.⁴⁴

MEASUREMENTS

Resilience

Given the complex nature of resilience, a reliable measurement of this psychological construct is difficult to achieve. There are more than 15 measures of resilience currently used in the published literature. The 3 highest psychometrically rated scales are the 6-item Brief Resilience Scale (BRS),⁴⁵ the 25-item Connor-Davidson Resilience Scale (CD-RISC),⁴⁶ and

the 37-item Resilience Scale for adults (RSA),⁴⁷ all of which were developed originally for use in adult populations. While top-rated among available measurements, these scales were described as being of only "moderate" quality.⁴⁸

The BRS considers resilience in its most basic definition—"the ability to bounce back from stress." It is the only scale that does not investigate the availability of any resources or characteristics known to facilitate resilience and focuses exclusively on personal agency.⁴⁵ The BRS has good internal consistency, interpretability, and construct validity, and it is the shortest assessment commonly used.⁴⁸

The CD-RISC is a widely used scale in published literature and has been translated into more than 25 languages.⁴⁶ The CD-RISC has 5 factors: (1) personal competence, high standards, and tenacity; (2) trust in one's instincts, tolerance of negative affect, and strengthening effects of stress; (3) positive acceptance of change and secure relationships; (4) control; and (5) spiritual influences. In a recent study of patients rehabilitating from spinal cord injury, the abbreviated 10-item CD-RISC showed the highest reliability, validity, and practicality of available CD-RISC scales.⁴⁹ A broad criticism of the available resilience scales is a focus on individual over environmental factors in assessing resilience.⁵⁰

The RSA thoroughly examines the interpersonal (eg, family connections and social support) and intrapersonal (eg, personal strength, social competence, and structured style factors), contributing to an individual's resilience. The RSA has recently demonstrated high validity and consistency in the prediction of depression and anxiety in the face of stressful life events.⁵¹ Otlans et al report that the RSA, CD-RISC, CD-RISC-10, and BRS have been used in the literature regarding resilience in orthopedic patients¹⁷ (see summary of relevant measurement tools in Table 1).

Patient Activation

The measure of patient activation, which is focused on the engagement and self-confidence that patients have in their care, captures many of the characteristics of the concept of resilience. It measures a patient's willingness to work with a physician to overcome illness as well as their ability to follow through on treatment plans in the face of the stress. Such properties are highly valuable in the assessment of psychological fitness for spinal surgery.³⁶ The 22-item patient activation measure (PAM-22) and subsequent PAM-13 were developed to conceptualize and measure this property and have both been proven to be valid and reliable.^{35,54}

Table 1. Summary of measurement tools for psychosocial traits.

Measurement Tool	Number of Items	Internal Consistency (Cronbach's α)	Test-Retest Reliability (r Value)	Validity
Brief Resilience Scale ^{45,48}	6	0.7–0.95	0.69	Construct
Connor-Davidson Resilience Scale-10 ⁴⁹	10	0.86	0.87	Construct
Resilience Scale for Adults	37	0.9	0.84	Construct
Pain self-efficacy questionnaire ⁵²	10	0.92	0.73	Construct
Short Grit Scale ⁵³	8	0.73–0.83	0.68	Criterion
Patient activation measure-13 ³⁵	22	0.87	0.85	Criterion

Grit

Grit is commonly measured using the 12-item Grit Scale²² or the revised 8-item Short Grit Scale (Grit-S),⁵³ both of which demonstrated sufficient psychometric properties in diverse samples in their development. The use of the Grit Scale is somewhat controversial due to its overlap with concepts of perseverance and self-control. In a recent psychometric evaluation of the Grit-S performed by Gonzalez et al, the Grit-S was found to focus more on consistency of interest rather than perseverance, limiting its assessment of the original definition of grit, as well as finding no difference between the grit subscales compared with the total score.⁴¹ A 5-item Grit Scale was adapted from the Grit-S, focusing primarily on the questions relating to perseverance. The scale displayed high psychometric properties and was found to be a good predictor of academic and career outcomes.⁵⁵

Self-Efficacy

As self-efficacy is a widely researched construct, there are many metrics available for a range of conditions. Of 14 scales assessed in a systematic review, no current scales assessing self-efficacy were suited to musculoskeletal rehabilitation.⁵⁶ In the current review evaluating spinal surgery and self-efficacy, the pain self-efficacy questionnaire (PSEQ) was the most widely used. The PSEQ is a 10-item questionnaire that assesses the confidence that people with ongoing pain have in performing activities with pain.⁵²

SPINE SURGERY AND RESILIENCE, PATIENT ACTIVATION, GRIT, AND SELF-EFFICACY

Resilience in patients undergoing spinal surgery for a non-spinal cord injury has been an understudied area, considering the complexity of spine pathology and surgical sequelae.¹⁷ In terms of presurgical evaluation, Ahmed et al sampled 195 patients undergoing evaluation for back and neck pain at an orthopedic spine clinic and were surveyed for resilience (BRS), pain self-efficacy (PSEQ), and disability (Neck Disability Index

and Oswestry Disability Index). The study revealed independent associations of low resilience and pain self-efficacy scores with greater functional disability in patients with low back and neck pain.⁵⁷ To build upon this finding, a prospective study of 248 patients undergoing laminectomy had patients complete resilience (BRS) and pain self-efficacy (PSEQ) surveys at 6 weeks and 12 months postoperatively.⁵⁸ Patients were assessed for physical function, pain interference, social participation, and disability at both time points, revealing a significant association between resilience and all listed outcomes as well as an association between pain self-efficacy and disability at 12 months. Interestingly, pain self-efficacy was a greater predictor of lower patient-reported lifestyle measures at 6 weeks than resilience, a finding paralleled in the previous study by Ahmed et al.^{57,58} These results, while limited by the number of studies, are a promising indication for the use of resilience and pain self-efficacy metrics in patient selection for spinal surgery.

Patient activation has been more directly investigated in the context of patient selection for spinal surgery. In a prospective study, 65 patients undergoing lumbar spine surgery were preoperatively assessed for level of patient activation using PAM-13 and subsequently evaluated at 12 and 24 months following surgery for functional disability and mental health. A high level of patient activation was significantly associated with improved functional improvement and engagement in physical therapy after surgery but not with improved mental health.^{59,60} Block et al expanded upon these findings in a larger study of 581 patients who underwent spinal surgery or spinal cord stimulator implantation.⁶¹ Again, higher levels of preoperative patient activation were associated with improved levels of pain and functional disability. Analyses also revealed that while low baseline levels of patient activation were associated with negative psychosocial risk factors, patient activation mediated the association between these negative risk factors and poorer surgical outcomes.⁶¹ Thus, patients with otherwise concerning psychosocial qualities, such as

depression and negative affect, may have better outcomes in spinal surgery than otherwise expected if they also possess a high level of patient activation. In contrast, a recent retrospective study of patient activation in 64 patients undergoing anterior cervical discectomy and fusion revealed conflicting results.⁶² Patients were grouped based on preoperative PAM-10 scores and surveyed for pain and functional disability at 6 weeks, 3 months, 6 months, and 12 months. The results demonstrated no association between patient activation and any measure of postoperative recovery.⁶² While this finding somewhat limits the indication of patient activation scores for assessment in neck surgery, this is a retrospective study with a relatively small sample size and is the only published study focusing on patient activation and neck surgery.⁶² Patient activation has potential as a tool for preoperative assessment of spinal surgery due to positive correlations to outcomes in large samples. Given the relatively few studies done in this space, further prospective studies on the effect of patient activation on elective spinal surgery are called for.

The psychosocial trait of grit has also been studied in a limited fashion in spine surgery. Mok et al retrospectively assessed grit (Grit-S) and self-control (Self-Scoring Self-Control Scale) in a sample of 69 patients who underwent elective spinal surgery. Patient-reported outcome measures were completed, and patients were subsequently grouped by whether they had achieved significant improvements in functional disability. No significant difference in either grit or self-control was found.⁶³ This study is the only study performed to our knowledge that focused on the effect of grit on spinal surgery outcomes. It is limited by its retrospective nature and lack of multiple or uniform follow-up periods for patient subgroups. Considering the prior research discussed on the effect of grit on predicting performance, this is still a worthwhile metric to consider in the area of spinal surgery.

Self-efficacy, while intuitively promising as a predictor of outcomes in spinal surgery, has been studied in a very limited fashion. A randomized control trial studying the effect of postoperative physical therapy in surgically corrected cervical radiculopathy found that self-efficacy scores (Swedish Self-Efficacy Scale) did not differ between those who did and did not receive the intervention.⁶⁴ While self-efficacy improved in both groups from preoperative assessment, self-efficacy was not studied as a modifier of functional outcomes, so conclusions regarding its effect on patient recovery are limited.

DISCUSSION

Resilience, patient activation, grit, and self-efficacy all have potential to predict success following spine surgery. Current evidence favors resilience and patient activation as the strongest metrics for use in preoperative evaluation for spine pathology. Resilience, while complex, has been well studied and may be assessed on a personal level in short scales, such as the BRS and the CD-RISC-10, which have strong predictive evidence in orthopedic surgical outcomes.¹⁷ If a surgeon wishes to capture a broader picture of a patient's resilience, including external social and cultural factors, the longer form RSA may be better suited.^{48,51} Patient activation, while including aspects of resilience, is a unique measure that is specific to the health care setting and may offer more practical insight for an individual patient. The PAM-13 has been the most used survey and has demonstrated successful correlation with patient outcomes.³⁵

Accurate collection of these measures may present challenges, with 1 recent study reporting significant changes in patient-reported outcome measurement information system scores between the preoperative visit and the day of surgery.⁶⁵ This finding indicates a need to standardize collection of data at a set time point before surgery, though more research is needed to determine the ideal time of assessment of character traits. To optimize the collection rates for patient-reported outcomes (PROs), clinics should increase physician training in PRO platforms, engage administrative support, and consider appointing a physician primary contact for a PRO program.⁶⁶ Prior experience in collecting patient outcomes has also been associated with higher collection rates.⁶⁶

Many of the psychological traits discussed in this review have been studied in greater detail in other orthopedic subspecialties as well as other chronic medical conditions.⁶⁷ The results from these studies have been promising, as Otlans et al report in their review of resilience in orthopedic patients, though the area of research is new and lacks many long-term, prospective studies.¹⁷ For example, Lim et al demonstrated that patients with hip fractures having higher preoperative resilience measured by CD-RISC-10 have improvements in physical function postoperatively unrelated to general mental health scores.⁶⁸ As another example, Tokish et al showed that higher postoperative resilience measured by BRS was significantly associated with improvements in American Shoulder and Elbow Surgeons and Penn scores in patients undergoing total shoulder arthroplasty.⁶⁷ There are numerous other studies with

Table 2. Summary of psychosocial screening.

Article	Study Design	Psychosocial Scales Used	Population	Number of Patients	Time From Injury or Follow-Up Period	Preoperative Collection	Level of Evidence ^a	Key Findings
Ahmed et al (2019) ⁵⁷	Cross-sectional survey	BRS, PSEQ	Low back pain, neck pain	195	NA	NA	IV	Independent negative associations of resilience and pain self-efficacy scores with ODI and NDI in patients with low back and neck pain
Coronado et al (2021) ⁵⁸	Retrospective cohort	BRS, PSEQ	Laminectomy	248	6 wk and 12 mo	No	III	Association with resilience and physical function, pain interference, social participation, and disability at both time points; association with pain self-efficacy and disability at 12 mo
Mok et al (2021) ⁶³	Retrospective medical record review, abstract	Short Grit Scale, Self-Scoring Self-Control Scale	ACDF, MI-D, MI-TLIF	69	6 mo for ACDF, 2 wk for MI-D, and 6 wk for MI-TLIF	No	III	No significant difference in either grit or self-control and ODI and NDI scores
Wibault et al (2017) ⁶⁹	Randomized controlled trial	Swedish version of the Self-Efficacy Scale	ACDF, PCF	202	6 mo	Yes	I	Self-efficacy improved in patients exposed to physical therapy and those not exposed to physical therapy postoperatively; however, self-efficacy was not studied as a modifier of functional outcomes
Block et al (2019) ⁶¹	Prospective cohort	PAM-13	Spinal surgery, spinal cord stimulation	581	mean 180 d (SD 79.1)	Yes	II	Patient activation mediated the association between negative risk factors and pain and disability
Skolasky et al (2011) ⁶⁰	Prospective cohort	PAM-13	Lumbar fusion	65	12 and 24 mo	Yes	II	Positive association with patient activation and functional improvement and engagement in physical therapy after surgery
Patel et al (2019) ⁶²	Retrospective cohort	PAM-10	ACDF	64	6 wk, 3 mo, 6 mo, and 12 mo	Yes	III	No association between patient activation and any measure of postoperative recovery

Abbreviations: ACDF, anterior cervical discectomy and fusion; BRS, Brief Resilience Scale; MI-D, minimally invasive lumbar decompression; MI-TLIF, minimally invasive transforaminal lumbar interbody fusion; NA, not available; NDI, Neck Disability Index; ODI, Oswestry Disability Index; PAM, patient activation measure; PCF, posterior cervical fusion; PSEQ, pain self-efficacy questionnaire.

^aLevel of evidence based upon the *Journal for Bone and Joint Surgery* criteria and focused on the psychosocial outcome of interest.⁷⁰

similar results; however, complex spine pathology requiring surgical intervention can often result in a prolonged recovery process with long-term implications that may differ from other orthopedic procedures. As such, the role of resilience may play a greater role in spine surgery and may be well suited as an area of study for preoperative risk stratification.

This review highlights the need for high-quality research in this area as well as a potential opportunity to improve patient outcomes through appropriate selection and targeted interventions. Future research should collect psychological data preoperatively, with validated measures as mentioned in this article. To definitively investigate the role of these traits in spine surgery, long-term studies with a large patient population are required. The research reviewed in this article on resilience, patient activation, grit, and self-efficacy in spine surgery only spans 7 articles, as shown in Table 2, 5 of which have been published since 2019. Follow-up has been mostly short term, and few studies have collected data preoperatively. Importantly, resilience can be improved with validated interventions focusing on a mixture of cognitive behavioral therapy and mindfulness.¹⁸ The recent excitement surrounding these psychological metrics and their potential to identify preoperative risk factors that can be intervened upon provides support for the continued study of this emerging field.

REFERENCES

1. MMPI Profiles in Patients with Low-Back Pain - ProQuest. www.proquest.com/docview/614263510?accountid=14749. Accessed January 15, 2022.
2. Pichot P, Perse J, Lekeous MO, Dureau JL, Perez CI, Rychewaert A. La personalite des sujets presentant des douleurs dorsales fonchonnelles valeur de l'inventaire multiphasique de personalite du minnesota. *Revue de Psychologie Applique*. 1972;22:145–172.
3. Vendrig AA. The minnesota multiphasic personality inventory and chronic pain: a conceptual analysis of a long-standing but complicated relationship. *Clinical Psychology Review*. 2000;20(5):533–559. doi:10.1016/S0272-7358(00)00053-2
4. Love AW, Peck CL. The MMPI and psychological factors in chronic low back pain: a review. *Pain*. 1987;28(1):1–12. doi:10.1016/0304-3959(87)91054-2
5. den Boer JJ, Oostendorp RAB, Beems T, Munneke M, Oerlemans M, Evers AWM. A systematic review of bio-psychosocial risk factors for an unfavourable outcome after lumbar disc surgery. *Eur Spine J*. 2006;15(5):527–536. doi:10.1007/s00586-005-0910-x
6. Weiner BK. Spine update: the biopsychosocial model and spine care. *Spine*. 2008;33(2):219–223. doi:10.1097/BRS.0b013e3181604572
7. Cheriyan T, Harris B, Cheriyan J, et al. Association between compensation status and outcomes in spine surgery: a meta-analysis of 31 studies. *The Spine Journal*. 2015;15(12):2564–2573. doi:10.1016/j.spinee.2015.09.033
8. Daniels AH, Kuris EO, Kleinhenz DT, Palumbo MA. Spine surgery outcomes in workers' compensation patients. *J Am Acad Orthop Surg*. 2017;25(10):e225–e234. doi:10.5435/JAAOS-D-16-00895
9. Anderson PA, Schwaegler PE, Cizek D, Leverson G. Work status as a predictor of surgical outcome of discogenic low back pain. *Spine (Phila Pa 1976)*. 2006;31(21):2510–2515. doi:10.1097/01.brs.0000239180.14933.b7
10. Laxton AW, Perrin RG. The relations between social support, life stress, and quality of life following spinal decompression surgery. *Spinal Cord*. 2003;41(10):553–558. doi:10.1038/sj.sc.3101432
11. Johnson BC, Vasquez-Montes D, Steinmetz L, et al. Association between nonmodifiable demographic factors and patient satisfaction scores in spine surgery clinics. *Orthopedics*. 2019;42(3):143–148. doi:10.3928/01477447-20190424-05
12. Junge A, Fröhlich M, Ahrens S, et al. Predictors of bad and good outcome of lumbar spine surgery. A prospective clinical study with 2 years' follow up. *Spine (Phila Pa 1976)*. 1996;21(9):1056–1064. doi:10.1097/00007632-199605010-00013
13. Block AR, Ohnmeiss DD, Guyer RD, Rashbaum RF, Hochschuler SH. The use of presurgical psychological screening to predict the outcome of spine surgery. *Spine J*. 2001;1(4):274–282. doi:10.1016/s1529-9430(01)00054-7
14. Junge A, Dvorak J, Ahrens S. Predictors of bad and good outcomes of lumbar disc surgery. A prospective clinical study with recommendations for screening to avoid bad outcomes. *Spine (Phila Pa 1976)*. 1995;20(4):460–468. doi:10.1097/00007632-199502001-00009
15. Marek RJ, Block AR, Ben-Porath YS. Validation of a psychological screening algorithm for predicting spine surgery outcomes. *Assessment*. 2019;26(5):915–928. doi:10.1177/1073191117719512
16. Finneson BE, Cooper VR. A lumbar disc surgery predictive score card. A retrospective evaluation. *Spine (Phila Pa 1976)*. 1979;4(2):141–144. doi:10.1097/00007632-197903000-00008
17. Otlans PT, Szukics PF, Bryan ST, Tjoumakaris FP, Freedman KB. Resilience in the orthopaedic patient. *Journal of Bone and Joint Surgery*. 2021;103(6):549–559. doi:10.2106/JBJS.20.00676
18. Joyce S, Shand F, Tighe J, Laurent SJ, Bryant RA, Harvey SB. Road to resilience: a systematic review and meta-analysis of resilience training programmes and interventions. *BMJ Open*. 2018;8(6):e017858. doi:10.1136/bmjopen-2017-017858
19. Epker J, Block AR. Presurgical psychological screening in back pain patients: a review. *The Clinical Journal of Pain*. 2001;17(3):200–205. doi:10.1097/00002508-200109000-00003
20. Spengler DM, Ouellette EA, Battié M, Zeh J. Elective discectomy for herniation of a lumbar disc. additional experience with an objective method. *The Journal of Bone & Joint Surgery*. 1990;72(2):230–237. doi:10.2106/00004623-199072020-00010
21. Southwick SM, Bonanno GA, Masten AS, Panter-Brick C, Yehuda R. Resilience definitions, theory, and challenges: interdisciplinary perspectives. *Eur J Psychotraumatol*. 2014;5. doi:10.3402/ejpt.v5.25338
22. Duckworth AL, Peterson C, Matthews MD, Kelly DR. Grit: perseverance and passion for long-term goals. *J Pers Soc Psychol*. 2007;92(6):1087–1101. doi:10.1037/0022-3514.92.6.1087
23. Sisto A, Vicinanza F, Campanozzi LL, Ricci G, Tartaglino D, Tambone V. Towards A transversal definition of

- psychological resilience: A literature review. *Medicina (Kaunas)*. 2019;55(11):E745. doi:10.3390/medicina55110745
24. Aburn G, Gott M, Hoare K. What is resilience? An integrative review of the empirical literature. *J Adv Nurs*. 2016;72(5):980–1000. doi:10.1111/jan.12888
 25. Resilience – APA Dictionary of Psychology. <https://dictionary.apa.org/resilience>. Accessed January 17, 2022.
 26. Luthar SS, Cicchetti D. The construct of resilience: implications for interventions and social policies. *Dev Psychopathol*. 2000;12(4):857–885. doi:10.1017/S0954579400004156
 27. Ezeamama AE, Elkins J, Simpson C, Smith SL, Allegra JC, Miles TP. Indicators of resilience and healthcare outcomes: findings from the 2010 health and retirement survey. *Qual Life Res*. 2016;25(4):1007–1015. doi:10.1007/s11136-015-1144-y
 28. Bonanno GA. Loss, trauma, and human resilience: have we underestimated the human capacity to thrive after extremely aversive events? *Am Psychol*. 2004;59(1):20–28. doi:10.1037/0003-066X.59.1.20
 29. Magaldi RJ, Staff I, Stovall AE, Stohler SA, Lewis CG. Impact of resilience on outcomes of total knee arthroplasty. *The Journal of Arthroplasty*. 2019;34(11):2620–2623. doi:10.1016/j.arth.2019.06.008
 30. Padesky CA, Mooney KA. Strengths-based cognitive-behavioural therapy: a four-step model to build resilience. *Clin Psychol Psychother*. 2012;19(4):283–290. doi:10.1002/cpp.1795
 31. Hassett AL, Finan PH. The role of resilience in the clinical management of chronic pain. *Curr Pain Headache Rep*. 2016;20(6):39. doi:10.1007/s11916-016-0567-7
 32. Bolton KW, Praetorius RT, Smith-Osborne A. Resilience protective factors in an older adult population: a qualitative interpretive meta-synthesis. *Social Work Research*. 2016;40(3):171–182. doi:10.1093/swr/svw008
 33. Färber F, Rosendahl J. Trait resilience and mental health in older adults: a meta-analytic review. *Personality and Mental Health*. 2020;14(4):361–375. doi:10.1002/pmh.1490
 34. Kilic SA, Dorstyn DS, Guiver NG. Examining factors that contribute to the process of resilience following spinal cord injury. *Spinal Cord*. 2013;51(7):553–557. doi:10.1038/sc.2013.25
 35. Hibbard JH, Stockard J, Mahoney ER, Tusler M. Development of the patient activation measure (PAM): conceptualizing and measuring activation in patients and consumers. *Health Serv Res*. 2004;39(4 Pt 1):1005–1026. doi:10.1111/j.1475-6773.2004.00269.x
 36. Block AR. Demoralization, patient activation, and the outcome of spine surgery. *Healthcare (Basel)*. 2016;4(1):11. doi:10.3390/healthcare4010011
 37. Grit – APA Dictionary of Psychology. <https://dictionary.apa.org/grit>. Accessed January 17, 2022.
 38. Louie PK, McCarthy MH, Albert TJ, Kim HJ. Reaching for peak performance during surgical training: the value in assessment tools and critical performance measures. *J Am Acad Orthop Surg*. 2020;28(17):e744–e751. doi:10.5435/JAAOS-D-20-00172
 39. Von Culin KR, Tsukayama E, Duckworth AL. Unpacking grit: motivational correlates of perseverance and passion for long-term goals. *J Posit Psychol*. 2014;9(4):306–312. doi:10.1080/17439760.2014.898320
 40. Credé M, Tynan MC, Harms PD. Much ado about grit: a meta-analytic synthesis of the grit literature. *Journal of Personality and Social Psychology*. 2017;113(3):492–511. doi:10.1037/pspp0000102
 41. Gonzalez O. A psychometric evaluation of the short grit scale: A closer look at its factor structure and scale functioning. *Eur J Psychol Assess*. 2020;36(4):646–657. doi:10.1027/1015-5759/a000535
 42. Self-Efficacy – APA Dictionary of Psychology. <https://dictionary.apa.org/self-efficacy>. Accessed January 17, 2022.
 43. Bandura A. *Self-Efficacy: The Exercise of Control*. Macmillan; 1997.
 44. Eklund RC, Tenenbaum G. *Encyclopedia of Sport and Exercise Psychology*. SAGE Publications; 2013.
 45. Smith BW, Dalen J, Wiggins K, Tooley E, Christopher P, Bernard J. The brief resilience scale: assessing the ability to bounce back. *Int J Behav Med*. 2008;15(3):194–200. doi:10.1080/10705500802222972
 46. Burns RA, Anstey KJ. The connor-davidson resilience scale (CD-RISC): testing the invariance of a uni-dimensional resilience measure that is independent of positive and negative affect. *Personality and Individual Differences*. 2010;48(5):527–531. doi:10.1016/j.paid.2009.11.026
 47. Friborg O, Barlaug D, Martinussen M, Rosenvinge JH, Hjemdal O. Resilience in relation to personality and intelligence. *Int J Methods Psychiatr Res*. 2005;14(1):29–42. doi:10.1002/mpr.15
 48. Windle G, Bennett KM, Noyes J. A methodological review of resilience measurement scales. *Health Qual Life Outcomes*. 2011;9:8. doi:10.1186/1477-7525-9-8
 49. Kuiper H, van Leeuwen CCM, Stolwijk-Swüste JM, Post MWM. Measuring resilience with the connor-davidson resilience scale (CD-RISC): which version to choose? *Spinal Cord*. 2019;57(5):360–366. doi:10.1038/s41393-019-0240-1
 50. Straus E, Dev SI, Moore RC. The measurement of resilience and grit: room for improvement. *Psychiatry Res*. 2020;285:112791. doi:10.1016/j.psychres.2020.112791
 51. Morote R, Hjemdal O, Kryszinska K, Martinez Uribe P, Corveleyn J. Resilience or hope? Incremental and convergent validity of the resilience scale for adults (RSA) and the herth hope scale (HHS) in the prediction of anxiety and depression. *BMC Psychol*. 2017;5(1):36. doi:10.1186/s40359-017-0205-0
 52. Nicholas MK. The pain self-efficacy questionnaire: taking pain into account. *Eur J Pain*. 2007;11(2):153–163. doi:10.1016/j.ejpain.2005.12.008
 53. Duckworth AL, Quinn PD. Development and validation of the short grit scale (grit-S). *J Pers Assess*. 2009;91(2):166–174. doi:10.1080/00223890802634290
 54. Hibbard JH, Mahoney ER, Stockard J, Tusler M. Development and testing of a short form of the patient activation measure. *Health Serv Res*. 2005;40(6 Pt 1):1918–1930. doi:10.1111/j.1475-6773.2005.00438.x
 55. Lechner CM, Danner D, Rammstedt B. Grit (effortful persistence) can be measured with a short scale, shows little variation across socio-demographic subgroups, and is associated with career success and career engagement. *PLoS One*. 2019;14(11):e0224814. doi:10.1371/journal.pone.0224814
 56. Picha KJ, Jochimsen KN, Heebner NR, et al. Measurements of self-efficacy in musculoskeletal rehabilitation: a systematic review. *Musculoskeletal Care*. 2018;16(4):471–488. doi:10.1002/msc.1362
 57. Ahmed SA, Shantharam G, Eltorai AEM, Hartnett DA, Goodman A, Daniels AH. The effect of psychosocial measures of resilience and self-efficacy in patients with neck and lower back pain. *Spine J*. 2019;19(2):232–237. doi:10.1016/j.spinee.2018.06.007
 58. Coronado RA, Robinette PE, Henry AL, et al. Bouncing back after lumbar spine surgery: early postoperative resilience is associated with 12-month physical function, pain interference,

social participation, and disability. *Spine J.* 2021;21(1):55–63. doi:10.1016/j.spinee.2020.07.013

59. Skolasky RL, Mackenzie EJ, Wegener ST, Riley LH. Patient activation and adherence to physical therapy in persons undergoing spine surgery. *Spine (Phila Pa 1976)*. 2008;33(21):E784–E791. doi:10.1097/BRS.0b013e31818027f1

60. Skolasky RL, Mackenzie EJ, Wegener ST, Riley LH. Patient activation and functional recovery in persons undergoing spine surgery. *J Bone Joint Surg Am.* 2011;93(18):1665–1671. doi:10.2106/JBJS.J.00855

61. Block AR, Marek RJ, Ben-Porath YS. Patient activation mediates the association between psychosocial risk factors and spine surgery results. *J Clin Psychol Med Settings.* 2019;26(2):123–130. doi:10.1007/s10880-018-9571-x

62. Patel DV, Yoo JS, Block AM, Karmarkar SS, Lamoutte EH, Singh K. Patient activation is not associated with postoperative outcomes following anterior cervical discectomy and fusion. *Clin Spine Surg.* 2019;32(10):E453–E456. doi:10.1097/BSD.0000000000000847

63. Mok J, Vaishnav AS, Louie P, et al. 222. do grit and self-control impact outcomes following spine surgery: a retrospective analysis. *The Spine Journal.* 2021;21(9):S114. doi:10.1016/j.spinee.2021.05.429

64. Jing D, Jia L. Assessment of patients' psychological state and self-efficacy associated with postoperative constipation after thoracolumbar fracture surgery. *J Int Med Res.* 2019;47(9):4215–4224. doi:10.1177/0300060519859732

65. Cross AG, Yedulla NR, Ziedas AC, et al. Trends in patient-reported outcomes measurement information system scores exist between day of surgical scheduling and day of surgery. *Arthroscopy: The Journal of Arthroscopic & Related Surgery.* 2022;38(1):139–144. doi:10.1016/j.arthro.2021.05.063

66. Sisodia RC, Dankers C, Orav J, et al. Factors associated with increased collection of patient-reported outcomes within a large health care system. *JAMA Network Open.* 2020;3(4):e202764. doi:10.1001/jamanetworkopen.2020.2764

67. Dombrowsky AR, Kirchner G, Isbell J, et al. Resilience correlates with patient reported outcomes after reverse total shoulder arthroplasty. *Orthopaedics & Traumatology: Surgery & Research.* 2021;107(1):102777. doi:10.1016/j.otsr.2020.102777

68. Lim KK, Matchar DB, Tan CS, et al. The association between psychological resilience and physical function among

older adults with hip fracture surgery. *Journal of the American Medical Directors Association.* 2020;21(2):260–266. doi:10.1016/j.jamda.2019.07.005

69. Wibault J, Öberg B, Dederig Å, et al. Neck-related physical function, self-efficacy, and coping strategies in patients with cervical radiculopathy: A randomized clinical trial of postoperative physiotherapy. *J Manipulative Physiol Ther.* 2017;40(5):330–339. doi:10.1016/j.jmpt.2017.02.012

70. JBJS, Inc. *Journals Level of Evidence: JBJS.* <https://journals-lww-com.libproxy2.usc.edu/jbjsjournal/pages/journals-level-of-evidence.aspx>. Accessed March 30, 2022.

Funding: The authors received no financial support for the research, authorship, and/or publication of this article.

Declaration of Conflicting Interests:

Jeffrey Wang reports payment for expert testimony from law firms; a leadership role and expense reimbursement from the AO Foundation Board; stock/stock options from Bone Biologics, Pearldiver, and Electrocore; and fellowship funding from AO Spine North America (paid to institution). The remaining authors have nothing to disclose.

IRB Approval: This work was exempt from IRB approval.

Corresponding Author: Jacob R. Ball, Department of Orthopaedic Surgery, Department of Orthopaedic Surgery, 1520 San Pablo St, Los Angeles, CA 90033, USA; Jacob.ball@med.usc.edu

Published 08 March 2023

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2023 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.