

Decompression Alone Compared to Decompression With Fusion in Patients With Lumbar Spondylolisthesis: Systematic Review, Meta-Analysis, and Meta-Regression

Raymond Pranata, Michael Anthonius Lim, Rachel Vania and Tjokorda Gde Bagus Mahadewa

Int J Spine Surg 2022, 16 (1) 71-80

doi: <https://doi.org/10.14444/8179>

<https://www.ijssurgery.com/content/16/1/71>

This information is current as of May 17, 2025.

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://ijssurgery.com/alerts>

Decompression Alone Compared to Decompression With Fusion in Patients With Lumbar Spondylolisthesis: Systematic Review, Meta-Analysis, and Meta-Regression

RAYMOND PRANATA, MD¹; MICHAEL ANTHONIUS LIM, MD, MHSM¹; RACHEL VANIA, MD¹; AND TJOKORDA GDE BAGUS MAHADEWA, MD, PHD²

¹Faculty of Medicine, Universitas Pelita Harapan, Tangerang, Indonesia; ²Department of Surgery, Division of Neurosurgery, Faculty of Medicine, Udayana University, Denpasar, Bali, Indonesia

ABSTRACT

Background: We aimed to synthesize the latest evidence on the efficacy and safety of decompression alone compared to decompression with fusion in patients with lumbar spondylolisthesis. We also aimed to evaluate factors affecting the efficacy and complications.

Methods: A systematic literature search was conducted using PubMed, Scopus, Europe PMC, Cochrane Central Database, and ClinicalTrials.gov. The main outcome was improvement in Oswestry Disability Index (ODI). The secondary outcome was back pain and leg pain improvement, complications, reoperation rate, duration of surgery, length of hospital stay, and blood loss.

Results: There were 3993 patients from 13 studies. Decompression with fusion was associated with greater reduction in ODI (mean difference 4.04 [95% CI 0.95, 7.13], $P = 0.01$) compared to decompression alone. Greater reduction in back (standardized mean difference [SMD] 0.27 [95% CI 0.00, 0.53], $P = 0.05$) and leg pain (SMD 0.13 [95% CI 0.06, 0.21], $P < 0.001$) was observed in the decompression with fusion group. Complications were similar in the 2 groups (OR 0.60 [95% CI 0.34, 1.04], $P = 0.07$). The reoperation rate was similar in both groups ($P = 0.54$). Decompression alone resulted in shorter duration of surgery (mean difference -85.18 minutes [95% CI -122.79 , -47.57], $P < 0.001$), less blood loss (mean difference -262.65 mL [95% CI -313.45 , -211.85], $P < 0.001$), and shorter hospital stay (mean difference -2.64 days [95% CI -3.58 , -1.70], $P < 0.001$). Empirical Bayes random-effects meta-regression showed that the rate of complication was influenced by age (coefficient 0.172, $P = 0.004$).

Conclusion: Decompression with fusion had greater efficacy than decompression alone but was associated with more blood loss, lengthier surgery, and hospitalization. In terms of complications, decompression alone may be beneficial in younger patients. (PROSPERO CRD42020211904)

Level of Evidence: 2A.

Lumbar Spine

Keywords: decompression, spinal fusion, spondylolisthesis, lumbar vertebra

INTRODUCTION

Lumbar spondylolisthesis is a degenerative condition in which one vertebra slips forward relative to the vertebra below, resulting in a spondylotic and narrowed spinal segment. This pathology typically causes low back pain, radiating to the buttocks and lower extremities, and neurogenic claudication due to concomitant lumbar spinal stenosis.^{1,2} As lumbar spinal stenosis symptoms stem from the dural sac or nerve root compression, the need for decompressing the neural structures becomes apparent.³ Conservative treatment is initially offered to symptomatic patients, but if the nonsurgical approach is no longer successful, surgical management in the form of decompression alone or decompression in combination with a spinal fusion is considered.⁴

To date, the selection of a surgical technique for lumbar spondylolisthesis remains debatable. Many studies

comparing the outcomes of 2 different surgical procedures have often had conflicting results. In this systematic review and meta-analysis, we aimed to synthesize the latest evidence on the efficacy and safety of decompression alone compared to decompression with fusion in patients with lumbar spondylolisthesis. We also aimed to evaluate factors affecting the efficacy and complications.

METHODS

This study follows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) reporting guideline. The protocol was registered in PROSPERO database (CRD42020211904; https://www.crd.york.ac.uk/prospero/display_record.php?RecordID=211904).

Eligibility Criteria

We included research articles including randomized controlled trials (RCTs) and observational studies (prospective or retrospective cohorts) that investigate the outcome of interest between patients with degenerative lumbar spondylolisthesis that underwent decompression alone or decompression with fusion with at least 12 months follow-up. Only published studies have been included in this study. We excluded abstract-only publications, letters, preprints, review articles, editorial/commentaries, case reports/case series/studies <20 patients, animal/cadaveric studies, nonEnglish language articles, and studies that did not report key exposures or outcomes of interest.

Search Strategy and Study Selection

A systematic literature search was conducted using PubMed, Scopus, Europe PMC, Cochrane Central Database, and Clinicaltrials.gov with keywords ((decompression alone) OR (decompression only) OR (decompression)) AND ((decompression plus fusion) OR (fusion)) AND (spondylolisthesis) for records published from inception up until September 29, 2020. Duplicates were removed, and the titles and abstracts of the remaining articles were independently assessed by 2 authors using the inclusion and exclusion criteria.

Data Extraction

Two authors performed data extraction independently using standardized extraction forms that included author, year, study design, age, gender, body mass index, baseline Oswestry Disability Index (ODI), length of follow-up, and the outcome of interest.

The main outcome of interest was the improvement in ODI postoperatively, reported as the mean change from baseline or ODI at the last follow-up. If a study reported both, mean change from baseline was preferred for analysis. Minimal clinically important difference (MCID) in the ODI improvement was ODI improvement above the cut-off point that the respective studies used.

The secondary outcome of interest includes improvement in back pain and leg pain, measured as improvement in either Numerical Rating Scale (NRS) or visual analog scale (VAS) postoperatively. Complications and reoperation rate were reported in the number of events per total sample. Duration of surgery was the length of the procedure which was reported in minutes. The length of hospital stay was the number of hospitalization days before the patient was discharged. Blood loss

was the volume of intraoperatively blood loss in milliliters.

The between-group differences in ODI, length of hospital stay, duration of surgery, and blood loss were reported in mean differences. Back pain and leg pain were reported as standardized mean differences (SMDs), because they include 2 different scale, NRS and VAS. MCID, complications, and reoperation rate were reported in odds ratios (ORs).

Risk of Bias Assessment

Risk of bias assessment was performed by 2 independent authors. Cochrane risk of bias assessment tools was used to evaluate RCTs and the Newcastle-Ottawa Scale (NOS) was used to assess observational studies.

Statistical Analysis

Review Manager 5.3 (Cochrane Collaboration) and Stata 16 (StataCorp LLC, Texas, USA) were used to perform meta-analysis in this study. We used the inverse variance method to pool continuous variables and obtain mean differences/SMD along with its standard deviations (SDs). To obtain OR along with its 95% confidence intervals (CIs), we used the Maentel-Haenszel formula. Random-effects model was used to pool the effect estimates regardless of heterogeneity. *P* values in this study were two-tailed, and a value of ≤ 0.05 was considered statistically significant. Chi-squared and Cochrane *Q* test were used to assess heterogeneity, *I* squared (*I*²) value of >50% and/or a *P* value of <0.10 was statistically significant for substantial heterogeneity. We performed regression-based Egger's test to evaluate the presence of small-study effects for the outcomes. The risk of publication bias was assessed using inverted-funnel plot. Empirical Bayes random-effects meta-regression analysis was performed using age, female sex, and ODI as covariates. Subgroup analyses were performed for RCTs, grade I spondylolisthesis, and single-vertebral level surgery.

RESULTS

Study Selection and Baseline Characteristics

The PRISMA flowchart is presented in Figure 1. We included 13 studies comprising 3993 patients in the qualitative and quantitative syntheses.¹⁻¹⁵ Three studies were RCTs, and 10 studies were cohorts. The baseline characteristics of the studies can be seen in Table 1 and Table 2. The mean age of the pooled analysis was 67.85 ± 18.93 .




Figure 1. PRISMA flowchart.

Oswestry Disability Index

Decompression with fusion was associated with greater reduction in ODI (mean difference 4.04 [95% CI 0.95, 7.13], $P = 0.01$; $I^2: 80\%$, $P < 0.001$) compared to decompression alone (Figure 2). The mean changes in ODI at the last follow-up was greater in the decompression with fusion group (mean difference 5.74 [95% CI 1.65, 9.82], $P = 0.006$; $I^2: 42\%$, $P = 0.16$). The MCID for ODI

improvement was similar (OR 0.76 [95% CI, 0.57, 1.07], $P = 0.12$; $I^2: 0\%$, $P = 0.40$).

Pain Scores

Meta-analysis showed that a greater reduction in back pain (SMD 0.27 [95% CI 0.00, 0.53], $P = 0.05$; $I^2: 90\%$, $P < 0.001$) (Figure 3A) and leg pain (SMD 0.13 [95% CI 0.06, 0.21], $P < 0.001$; $I^2: 0\%$, $P = 0.49$) (Figure 3B) was

Table 1. Baseline characteristics of included studies.

| Authors | Design | Sample, n | Age, y, mean ± SD | Female, % | BMI | Baseline ODI, mean ± SD | Last Follow-up, mo | NOS |
|--------------------------------|--------------------------------------|-----------|-------------------|-----------|------------|-------------------------|----------------------|------------------|
| Austevoll 2020 ² | Multicenter prospective cohort (PSM) | 570 | 64.7 ± 9.5 | 72.5 | 26.9 ± 4.5 | 41.1 ± 14.9 | 12 | 8 |
| Chan 2019 ⁴ | Multicenter prospective cohort | 426 | 62.5 ± 11.5 | 59.2 | 30.5 ± 6.5 | 47.7 ± 16.9 | 12 | 7 |
| Försth 2013 ¹⁶ | Multicenter prospective cohort | 1306 | 69.4 ± 29.6 | 65.7 | NR | 44.4 ± 15.8 | 25 | 6 |
| Försth 2016 ⁶ | Multicenter RCT | 135 | 67.5 ± 7.0 | 79.2 | NR | 41.0 ± 13.5 | 24 | RoB ^a |
| Ghogawala 2004 ⁷ | Multicenter prospective cohort | 34 | 68.8 ± 8.0 | 68.0 | NR | 41.2 | 12 | 6 |
| Ghogawala 2016 ⁸ | Multicenter RCT | 66 | 66.6 ± 7.6 | 80.3 | NR | 37.5 ± 15.5 | 48 (24 for analysis) | RoB ^a |
| Inose 2018 ³ | Multicenter RCT | 60 | 62.3 ± 7.7 | 53.3 | NR | NR | 60 | RoB ^a |
| Kim 2018 ⁹ | Prospective cohort | 129 | 66.1 ± 8.9 | 77.5 | 21.4 ± 5.2 | NR | 24 | 7 |
| Kleinstueck 2011 ¹⁰ | Prospective cohort | 213 | 68.9 ± 9.3 | 72.7 | NR | NR | 12 | 7 |
| Matsudaira 2005 ¹¹ | Retrospective cohort | 39 | 67.5 ± 6.9 | 59.5 | NR | NR | 24 | 7 |
| Park 2012 ¹² | Retrospective cohort | 45 | 64.3 ± 19.6 | 82.2 | 25.8 ± 4.7 | 26.9 ± 5.6 | 63 | 7 |
| Sigmundsson 2015 ¹³ | Retrospective cohort | 839 | 70.3 ± 9.5 | 76.9 | NR | 44.9 ± 14.8 | 36 | 6 |
| Ulrich 2017 ¹⁵ | Retrospective cohort | 131 | 72.3 ± 8.4 | 58.0 | NR | NR | 36 | 7 |

Abbreviations: BMI, body mass index; NOS, Newcastle-Ottawa Scale; NR, not reported; ODI, Oswestry Disability Index; PSM, propensity-score matching; RoB, Cochrane risk of bias assessment tool.

^aSee Figure 5A.

observed in the decompression with fusion compared to decompression alone.

Complications

The rate of postoperative complications was similar (OR 0.60 [95% CI 0.34, 1.04], $P = 0.07$; I^2 : 35%, $P = 0.13$) (Figure 4) compared to the decompression with fusion group. The reoperation rate was similar in both groups (OR 1.17 [95% CI 0.71, 1.93], $P = 0.54$; I^2 : 52%, $P = 0.08$).

Duration of Surgery

The duration of surgery was shorter in the decompression alone group (mean difference -85.18 minutes [95% CI -122.79, -47.57], $P < 0.001$; I^2 : 99%, $P < 0.001$) compared to decompression with fusion group. Decompression alone group was associated with less blood loss (mean difference -262.65 mL [95% CI -313.45, -211.85], $P < 0.001$; I^2 : 87%, $P < 0.001$).

Length of Hospital Stay

The length of hospital stay was shorter in the decompression alone group (mean difference -2.64 days [95% CI -3.58, -1.70], $P < 0.001$; I^2 : 91%, $P < 0.001$).

Risk of Bias Assessment

The risk of bias assessment for RCTs can be seen in Figure 5A, and the NOS for observational studies can be seen in Table 1. Funnel plot analysis showed a relatively symmetrical distribution for the ODI (Figure 5B) and

complications (Figure 5C). Regression-based Egger's test showed no indication of small-study effects for the ODI ($P = 0.219$), leg pain ($P = 0.257$) improvement, and number of complications ($P = 0.810$). Back pain improvement was statistically significant for small-study effects ($P = 0.009$).

Meta-Regression

Empirical Bayes random-effects meta-regression showed that the difference in ODI improvement between decompression alone and decompression with fusion group was not influenced by preoperative ODI (coefficient -0.31, $P = 0.322$), age (coefficient -0.519, $P = 0.619$), and female sex (coefficient 0.309, $P = 0.352$). Age (coefficient -0.533, $P = 0.411$), female sex (coefficient -0.174, $P = 0.449$), and preoperative ODI (coefficient -0.022, $P = 0.948$) did not significantly influence the difference in back pain improvement between the 2 groups. Difference in leg pain improvement was not significantly affected by age (coefficient -0.020, $P = 0.904$), female sex (coefficient -0.026, $P = 0.672$), and preoperative ODI (coefficient 0.005, $P = 0.974$). Meanwhile, the rate of complications varied significantly by age (coefficient 0.172, $P = 0.004$) but not by female sex (coefficient 0.041, $P = 0.115$). We further performed subgroup analysis in studies with mean age less than 67.8 years old (the mean age of the pooled analysis) and found that complications were significantly decreased in the decompression alone group (OR 0.46 [95% CI 0.23, 0.92], $P = 0.03$; I^2 : 36%, $P = 0.15$).

Table 2. Inclusion criteria and types of operation of included studies.

| Authors | Inclusion Criteria | No. of Vertebral Levels | Types of Operation | |
|--------------------------------|--|-------------------------|---|--|
| | | | Decompression Alone | Decompression + Fusion |
| Austevoll 2020 ² | Diagnosed both spinal stenosis and spondylolisthesis, and then operated with decompression or with decompression + fusion | 1–2 | Microdecompression with midline preservation | Instrumented fusion |
| Chan 2019 ⁴ | Diagnosed grade one spondylolisthesis, and then underwent posterior-only surgery at the index level of lumbar spondylolisthesis (decompression alone or fusion with or without decompression), or underwent single-segment surgery (single-level disc decompression [eg, L4 or L4–five decompressive laminectomy]) or a 2-level vertebral fusion (eg, L4–five posterior spinal fusion) | 1 | Laminectomy | Instrumented fusion |
| Försth 2013 ^{16a} | Aged >50 y at the time of surgery, diagnosed spinal stenosis on 1–2 adjacent levels between L2 and L5, with or without spondylolisthesis, and then operated with decompression or with decompression + fusion | 1–2 | Decompression | Instrumented or noninstrumented fusion |
| Försth 2016 ^{6a} | Aged between 50 and 80 y, diagnosed spinal stenosis on 1–2 adjacent levels between L2 and sacrum, with or without spondylolisthesis, symptomatic (pseudoclaudication in one or both legs and back pain [VAS > 30]) for >6 mo, and then operated with decompression or with decompression + fusion | 1–2 | Decompression | Instrumented or noninstrumented fusion |
| Ghogawala 2004 ⁷ | Aged between 50 and 81 y, diagnosed grade one spondylolisthesis, and then operated with decompression or with decompression + fusion | - | Decompression with aggressive facet-sparing technique | Posterolateral instrumented fusion |
| Ghogawala 2016 ⁸ | Aged between 50 and 81 y, diagnosed stable grade one spondylolisthesis with spinal stenosis, symptomatic (neurogenic claudication with or without lumbar radiculopathy), and then operated with decompression or with decompression + fusion | 1 | Complete laminectomy with partial removal of medial facet joint | Posterolateral instrumented fusion |
| Inose 2018 ³ | Diagnosed spinal stenosis on one level at the L4-5, with spondylolisthesis, and then operated with decompression or decompression + fusion or decompression + stabilization | 1 | Decompression | Posterolateral instrumented fusion with autogenous iliac bone graft and pedicle screw |
| Kim 2018 ⁹ | Diagnosed grade one spondylolisthesis at L4-5, symptomatic, and then operated with decompression or with decompression + fusion | - | Decompressive laminectomy, partial facetectomy, <50% of the inferior articular processes in the transverse dimension, foraminotomy according to the extent of radiculopathy, with both facet joints preserved | Circumferential fusion with pedicle screws, posterolateral fusion with autogenous iliac bone graft, and posterior lumbar interbody fusion using a titanium cage filled with a local bone graft |
| Kleinstueck 2011 ¹⁰ | Diagnosed spondylolisthesis with a maximum of 3 levels affected, and then operated with decompression or with decompression + fusion | 1–3 | Decompression | Instrumented fusion |
| Matsudaira 2005 ¹¹ | Diagnosed both spinal stenosis and spondylolisthesis on single level at L4/5, and then operated with decompression or with decompression + fusion | 1 | Laminoplasty to preserve the integrity of midline structure | Posterolateral fusion and pedicle screw instrumentation |
| Park 2012 ¹² | Diagnosed stable grade one spondylolisthesis, and then operated with decompression or with decompression + fusion | 1 | Laminectomy | Instrumented fusion |
| Sigmundsson 2015 ¹³ | Diagnosed both spinal stenosis and spondylolisthesis on single level, and then operated with decompression or with decompression + fusion | 1 | Decompression | Posterolateral instrumented fusion |
| Ulrich 2017 ¹⁵ | Diagnosed both spinal stenosis and spondylolisthesis, and then operated with decompression or with decompression + fusion | 1–2 | Standard open or microscopic decompression of the lateral recess and the foramina | Implantation of pedicle screws with rods and intersomatic fusion and cage(s) at the affected level(s) |

^aOnly patients with spondylolisthesis were included in the analysis.

Subgroup Analysis for RCTs

Decompression alone and decompression with fusion resulted in a similar rate of ODI improvement (mean difference 1.78 [95% CI -10.34, 13.91], $P = 0.77$; $I^2: 80\%$, $P = 0.03$), reoperation (OR 1.39 [95% CI 0.33, 5.93], $P = 0.66$; $I^2: 80\%$, $P = 0.03$), complications (OR 0.49 [95% CI 0.11, 2.23], $P = 0.36$; $I^2: 68\%$, $P = 0.04$). Decompression alone was associated with reduced blood loss (mean difference -345.00 mL [95% CI -458.89, -231.11], $P < 0.001$; $I^2: 71\%$, $P = 0.03$),

length of surgery (mean difference -104.40 [95% CI -168.75, -40.04], $P = 0.001$; $I^2: 96\%$, $P < 0.001$), and length of stay (mean difference -2.14 d [95% CI -3.12, -1.16], $P < 0.001$; $I^2: 49\%$, $P = 0.14$). There was only one RCT that evaluated back and leg pain improvement; hence, it was not included in the subgroup analysis.

Subgroup Analysis for Grade I Spondylolisthesis

Meta-analysis showed that decompression with fusion was associated with greater ODI improvement




Figure 2. Oswestry Disability Index (ODI) outcomes. DCM, decompression.

(mean difference 6.48 [95% CI 1.17, 11.79], $P < 0.001$; I^2 : 72%, $P = 0.003$). There was similar improvement in terms of back (SMD 0.26 [95% CI -0.58, 1.10], $P = 0.55$; I^2 : 95%, $P < 0.001$) and leg pain (SMD 0.02 [95% CI -0.17, 0.20], $P = 0.86$; I^2 : 0%, $P = 0.68$). Both have similar complications (OR 0.77 [95% CI 0.45, 1.33], $P = 0.35$; I^2 : 0%, $P = 0.92$) and reoperation rate (OR 1.35 [95% CI 0.56, 3.23], $P = 0.50$; I^2 : 61%, $P = 0.08$). Decompression alone was associated with reduced blood loss (mean difference -267.15 mL [95% CI -326.17, -208.14], $P < 0.001$; I^2 : 90%, $P < 0.001$), length of surgery (mean difference -81.21 minutes [95% CI -129.40, -33.03], $P = 0.001$; I^2 : 98%, $P < 0.001$), and length of stay (mean difference -2.09 days [95% CI -2.69, -1.49], $P < 0.001$; I^2 : 62%, $P = 0.05$).

Subgroup Analysis for Single-Vertebral Level Surgery

Subgroup analysis showed that decompression with fusion was associated with greater ODI improvement (mean difference 4.15 [95% CI 1.48, 6.82], $P = 0.002$; I^2 : 0%, $P = 0.50$) compared to decompression only. Similar improvement in terms of back (SMD 0.15 [95% CI -0.20, 0.51], $P = 0.39$; I^2 : 76%, $P = 0.01$) and leg pain (SMD 0.13 [95% CI -0.02, 0.28], $P = 0.08$; I^2 : 0%, $P = 0.65$) was observed. The 2 procedures have similar complications (OR 0.42 [95% CI 0.09, 1.94], $P = 0.27$; I^2 : 37%, $P = 0.19$) and reoperation rate (OR 2.05 [95% CI 0.94, 4.45], $P = 0.07$; I^2 : 10%, $P = 0.29$). Decompression alone resulted in less blood loss (mean difference -277.09

mL [95% CI -395.15, -159.03], $P < 0.001$; I^2 : 89%, $P < 0.001$), shorter length of surgery (mean difference -115.38 minutes [95% CI -160.05, -70.71], $P < 0.001$; I^2 : 91%, $P < 0.001$), and shorter length of stay (mean difference -2.03 days [95% CI -2.62, -1.44], $P < 0.001$; I^2 : 70%, $P = 0.04$).

DISCUSSION

This meta-analysis showed that decompression with fusion resulted in a more significant improvement in ODI and pain scale compared to decompression alone. However, the duration of surgery, operative blood loss, and the length of hospital stay were lesser in the decompression alone group. Although the trend for complications was lower in decompression alone group, it was not statistically significant. The summary of meta-analysis is available in Table 3.

Consideration remains regarding the extent of surgery required in each individual case and whether analysis of the preoperative symptoms can assist in selecting the most appropriate procedure. Currently, there is no clear guidelines about which surgery is best for any given patient. In a clinical practice, patient selection is typically based on the symptoms arising from mechanical low back pain and radiating buttock and leg pain and their relationship with each other. A patient presenting with nerve compression due to stenosis as the main symptom, without the presence of major low back symptoms, may be advised to undergo a simple decompression rather than a more extensive fusion procedure




Figure 3. (A) Back pain and (B) leg pain outcomes. DCM, decompression.

even though there is an underlying slippage, which may indicate an inherent instability.¹⁰

Decompression has become the treatment of choice for severe spinal stenosis and provides better outcomes

than nonoperative management, but it is often associated with postoperative instability.¹⁶ Several factors may increase the likelihood of instability after laminectomy for spondylolisthesis, such as facet angle, disc




Figure 4. Complications. DCM, decompression.




Figure 5. Risk of bias assessment. Risk of bias assessment for randomized controlled trials (A). Funnel plot analysis for Oswestry Disability Index (B) and complications (C).

height, and motion on flexion-extension radiographs.⁸ Some surgeons consider spondylolisthesis with a 3-mm translation as a sign of instability; therefore, some facilities recommend performing instrumentation surgery regardless of the severity of spondylolisthesis. Decompression with fusion technique has since become the standard treatment for degenerative spondylolisthesis.³

Many believe that adding fusion to decompression will result in a systematically better outcome. One plausible explanation for this assumption is that decompression alone does not adequately help patients with predominantly stenotic symptoms, because the underlying cause of the stenosis, such as instability, is left untreated. Another explanation is that more stenosis would present due to the slippage, and this condition requires more extensive resection, which is often impossible without creating more instability, and therefore this cannot be resolved by decompression alone.¹⁰

Performing simultaneous fusion in elderly population, who generally have multiple comorbidities, may lead to a greater risk of life-threatening perioperative complications (ie, pneumonia, respiratory failure, and acute myocardial infarct) and a higher rate of death.^{10,16} Osteopenia or osteoporosis is a condition that commonly coexists in the elderly that increases the risk of screw loosening and sinking of the intersomatic cage.¹⁵ Although age may be a factor limiting the effectiveness of lumbar surgery, it is not associated with worse outcomes when fusion is used to complement decompression compared to decompression alone.⁷

Even though the use of pedicle screws may result in higher fusion rates, it does not necessarily lead to improvements in low back pain and radiculopathy; this raises the question of which procedure is superior to others, and also whether localized spinal instability due to spondylolisthesis causes lower back pain and/or decreased quality of life.³ Decompression alone is performed with less resection of the bony elements and thus in a more limited space compared to decompression with fusion, where the exposure is usually more extensive and resection can be carried out more freely without the risk of creating increased instability.¹⁰ Our meta-analysis showed that the improvement in ODI and pain scale is better in patients receiving decompression with fusion. However, the need for instrumentation in fusion surgery carries the risk of infection due to osteosynthesis material and the risk of hardware malposition and failure, necessitating reoperation.^{4,15} Our study showed that the reoperation rate was similar in the 2 groups. The longer duration of decompression with fusion procedure compared to decompression alone also increases the risks of anesthesia and their consequences

Table 3. Summary of findings.

| Outcomes | Primary Analysis | RCT Subgroup | Grade I Spondylolisthesis | Single-Vertebral Level Surgery |
|---------------------------|---|---|---|---|
| Oswestry Disability Index | MD 4.04 [0.95, 7.13], $P = 0.01$; $I^2: 80\%$ | MD 1.78 [-10.34, 13.91], $P = 0.77$; $I^2: 80\%$ | MD 6.48 [1.17, 11.79], $P < 0.001$; $I^2: 72\%$ | MD 4.15 [1.48, 6.82], $P = 0.002$; $I^2: 0\%$ |
| MCID ODI | OR 0.76 [0.57, 1.07], $P = 0.12$; $I^2: 0\%$ | Insufficient data | Insufficient data | Insufficient data |
| Back pain | SMD 0.27 [0.00, 0.53], $P = 0.05$; $I^2: 90\%$ | Insufficient data | SMD 0.26 [-0.58, 1.10], $P = 0.55$; $I^2: 95\%$ | SMD 0.15 [-0.20, 0.51], $P = 0.39$; $I^2: 76\%$ |
| Leg pain | SMD 0.13 [0.06, 0.21], $P < 0.001$; $I^2: 0\%$ | Insufficient data | SMD 0.02 [-0.17, 0.20], $P = 0.86$; $I^2: 0\%$ | SMD 0.13 [-0.02, 0.28], $P = 0.08$; $I^2: 0\%$ |
| Complications | OR 0.60 [0.34, 1.04], $P = 0.07$; $I^2: 35\%$ | OR 0.49 [0.11, 2.23], $P = 0.36$; $I^2: 68\%$ | OR 0.77 [0.45, 1.33], $P = 0.35$; $I^2: 0\%$ | OR 0.42 [0.09, 1.94], $P = 0.27$; $I^2: 37\%$ |
| Reoperation | OR 1.17 [0.71, 1.93], $P = 0.54$; $I^2: 52\%$ | OR 1.39 [0.33, 5.93], $P = 0.66$; $I^2: 80\%$ | Insufficient data | Insufficient data |
| Duration of surgery | MD -85.18 min [-122.79, -47.57], $P < 0.001$; $I^2: 99\%$ | MD -104.40 min [-168.75, -40.04], $P = 0.001$; $I^2: 96\%$ | MD -81.21 min [-129.40, -33.03], $P = 0.001$; $I^2: 98\%$ | MD -115.38 min [-160.05, -70.71], $P < 0.001$; $I^2: 91\%$ |
| Blood loss | MD -262.65 mL [-313.45, -211.85], $P < 0.001$; $I^2: 87\%$ | MD -345.00 mL [-458.89, -231.11], $P < 0.001$; $I^2: 71\%$ | MD -267.15 mL [-326.17, -208.14], $P < 0.001$; $I^2: 90\%$ | MD -277.09 mL [-395.15, -159.03], $P < 0.001$; $I^2: 89\%$ |
| Length of hospital stay | MD -2.64 d [-3.58, -1.70], $P < 0.001$; $I^2: 91\%$ | MD -2.14 d [-3.12, -1.16], $P < 0.001$; $I^2: 49\%$ | MD -2.09 d [-2.69, -1.49], $P < 0.001$; $I^2: 62\%$ | MD -2.03 d [-2.62, -1.44], $P < 0.001$; $I^2: 70\%$ |

Data in brackets represent 95% CIs.

in the elderly population.¹⁵ Furthermore, there is a risk of accelerated degenerative changes adjacent to a lumbar fusion in elderly individuals.¹⁶

Meta-regression analysis showed that in terms of complications, decompression alone may be beneficial in younger patients. Increasing age, although controversial, has been shown to affect the prognosis in lumbar spinal surgery negatively.^{14,17-20} One of the plausible explanations for the possible benefit in the younger age is the rate of complication is higher in the elderly due to age or other medical comorbidities. Thus, the complications are more avoidable in the younger groups.

One study provided no standard deviation, P values, or other alternative variables required to calculate effect estimate; the study reported that both methods result in satisfactory clinical outcomes.²¹ One study was excluded because of the unclear number of samples at final follow-up; it showed that both decompression alone and decompression with fusion resulted in similar 5-year reoperation rates.²² Studies showed that both surgical modalities resulted in similar improvement in quality of life measured by EuroQol 5-dimensions (EQ-5D) and Short Form-36 (SF-36).^{1,6-9,12,13,16} However, Kleinstueck et al reported that the improvement in Core Outcome Measures Index was significantly greater in the decompression with fusion group.¹⁰ In terms of expense, Yagi et al showed that both decompression alone and decompression with fusion is equally cost-effective.²³

Clinical Implications

This meta-analysis indicates that decompression with fusion has greater efficacy than decompression alone but was associated with lengthier surgery and hospitalization. In terms of complications, decompression alone may be beneficial in younger patients. Thus, decompression alone might be more valuable in younger patients, although further RCT in younger patients is needed. Nevertheless,

in the presence of instability, decompression alone may have limited success. Thus, in this subset of patients, decompression with fusion might be a procedure of choice. The studies included involvement of 1-3 vertebral levels, patients with >3 vertebral involvements may benefit more from the decompression with fusion as opposed to decompression alone.

Limitations

The primary limitation of this meta-analysis is that the several included studies have a moderate-high risk of bias. Additionally, there were only 3 RCTs available for analysis. Several studies did not report their spondylolisthesis grade, which may influence the surgical outcome.

CONCLUSIONS

Decompression with fusion results in a greater improvement in ODI and pain scale compared to decompression alone. Duration of surgery, operative blood loss, and the length of hospital stay were lesser in the decompression alone group. In terms of complications, decompression alone may be beneficial in younger patients.

REFERENCES

1. Austevoll IM, Gjestad R, Brox JJ, et al. The effectiveness of decompression alone compared with additional fusion for lumbar spinal stenosis with degenerative spondylolisthesis: a pragmatic comparative non-inferiority observational study from the Norwegian Registry for Spine Surgery. *Eur Spine J.* 2017;26(2):404-413. doi:10.1007/s00586-016-4683-1
2. Austevoll IM, Gjestad R, Solberg T, et al. Comparative effectiveness of microdecompression alone vs decompression plus instrumented fusion in lumbar degenerative spondylolisthesis. *JAMA Netw Open.* 2020;3(9):e2015015. doi:10.1001/jamanetworkopen.2020.15015
3. Inose H, Kato T, Yuasa M, et al. Comparison of decompression, decompression plus fusion, and decompression plus

- stabilization for degenerative spondylolisthesis: a prospective, randomized study. *Clin Spine Surg.* 2018;31(7):E347–E352. doi:10.1097/BSD.0000000000000659
4. Chan AK, Bisson EF, Bydon M, et al. Laminectomy alone versus fusion for grade 1 lumbar spondylolisthesis in 426 patients from the prospective Quality Outcomes Database. *J Neurosurg Spine.* 2019;30(2):234–241. doi:10.3171/2018.8.SPINE17913
 5. Parsons N, Smith RC, Griffin XL, Stengel D. Research fraud and the bone. *Joint Journal J Bone Jt Surgery-British.* 2013;95(7):866–867. doi:10.1302/0301-620X.95B7
 6. Försth P, Ólafsson G, Carlsson T, et al. A randomized, controlled trial of fusion surgery for lumbar spinal stenosis. *N Engl J Med.* 2016;374(15):1413–1423. doi:10.1056/NEJMoa1513721
 7. Ghogawala Z, Benzel EC, Amin-Hanjani S, et al. Prospective outcomes evaluation after decompression with or without instrumented fusion for lumbar stenosis and degenerative Grade I spondylolisthesis. *J Neurosurg Spine.* 2004;1(3):267–272. doi:10.3171/spi.2004.1.3.0267
 8. Ghogawala Z, Dziura J, Butler WE, et al. Laminectomy plus fusion versus laminectomy alone for lumbar spondylolisthesis. *N Engl J Med.* 2016;374(15):1424–1434. doi:10.1056/NEJMoa1508788
 9. Kim S-I, Ha K-Y, Kim Y-H, Kim Y-H, Oh I-S. A comparative study of decompressive laminectomy and posterior lumbar interbody fusion in grade I degenerative lumbar spondylolisthesis. *Indian J Orthop.* 2018;52(4):358–362. doi:10.4103/ortho.IJOrtho_330_16
 10. Kleinstueck FS, Fekete TF, Mannion AF, et al. To fuse or not to fuse in lumbar degenerative spondylolisthesis: do baseline symptoms help provide the answer? *Eur Spine J.* 2012;21(2):268–275. doi:10.1007/s00586-011-1896-1
 11. Matsudaira K, Yamazaki T, Seichi A, et al. Spinal stenosis in grade I degenerative lumbar spondylolisthesis: a comparative study of outcomes following laminoplasty and laminectomy with instrumented spinal fusion. *J Orthop Sci.* 2005;10(3):270–276. doi:10.1007/s00776-005-0887-7
 12. Park JH, Hyun SJ, Roh SW, Rhim SC. A comparison of unilateral laminectomy with bilateral decompression and fusion surgery in the treatment of grade I lumbar degenerative spondylolisthesis. *Acta Neurochir (Wien).* 2012;154(7):1205–1212. doi:10.1007/s00701-012-1394-1
 13. Sigmundsson FG, Jönsson B, Strömquist B. Outcome of decompression with and without fusion in spinal stenosis with degenerative spondylolisthesis in relation to preoperative pain pattern: a register study of 1,624 patients. *Spine J.* 2015;15(4):638–646. doi:10.1016/j.spinee.2014.11.020
 14. Okuda S, Oda T, Miyauchi A, Haku T, Yamamoto T, Iwasaki M. Surgical outcomes of posterior lumbar interbody fusion in elderly patients. *J Bone Joint Surg Am.* 2006;88(12):2714–2720. doi:10.2106/JBJS.F.00186
 15. Ulrich NH, Burgstaller JM, Pichierri G, et al. Decompression surgery alone versus decompression plus fusion in symptomatic lumbar spinal stenosis. *Spine.* 2017;42(18):E1077–E1086. doi:10.1097/BRS.0000000000002068
 16. Försth P, Michaëlsson K, Sandén B. Does fusion improve the outcome after decompressive surgery for lumbar spinal stenosis? *The Bone & Joint Journal.* 2013;95-B(7):960–965. doi:10.1302/0301-620X.95B7.30776
 17. Little DG, MacDonald D. The use of the percentage change in Oswestry disability index score as an outcome measure in lumbar spinal surgery. *Spine.* 1994;19(19):2139–2142. doi:10.1097/00007632-199410000-00001
 18. Arinzon ZH, Fredman B, Zohar E, et al. Surgical management of spinal stenosis: a comparison of immediate and long term outcome in two geriatric patient populations. *Arch Gerontol Geriatr.* 2003;36(3):273–279. doi:10.1016/s0167-4943(02)00172-3
 19. Ragab AA, Fye MA, Bohlman HH. Surgery of the lumbar spine for spinal stenosis in 118 patients 70 years of age or older. *Spine (Phila Pa 1976).* 2003;28(4):348–353. doi:10.1097/01.BRS.0000048494.66599.DF
 20. Fredman B, Arinzon Z, Zohar E, et al. Observations on the safety and efficacy of surgical decompression for lumbar spinal stenosis in geriatric patients. *Eur Spine J.* 2002;11(6):571–574. doi:10.1007/s00586-002-0409-7
 21. Ha D-H, Kim T-K, Oh S-K, Cho H-G, Kim K-R, Shim D-M. Results of decompression alone in patients with lumbar spinal stenosis and degenerative spondylolisthesis: a minimum 5-year follow-up. *Clin Orthop Surg.* 2020;12(2):187–193. doi:10.4055/cios19110
 22. Vorhies JS, Hernandez-Boussard T, Alamin T. Treatment of degenerative lumbar spondylolisthesis with fusion or decompression alone results in similar rates of reoperation at 5 years. *Clin Spine Surg.* 2018;31(1):E74–E79. doi:10.1097/BSD.0000000000000564
 23. Yagi M, Fujita N, Okada E, et al. Comparisons of direct costs, outcomes, and cost-utility of decompression surgery with fusion versus decompression alone for degenerative lumbar spondylolisthesis. *J Orthop Sci.* 2018;23(4):653–657. doi:10.1016/j.jos.2018.04.001

Funding: The authors received no financial support for the research, authorship, and/or publication of this article.

Declaration of Conflicting Interests:

All authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers' bureaus; membership, employment, consultancies, stock ownership, or other equity interest; and expert testimony or patent-licensing arrangements), or nonfinancial interest (such as personal or professional relationships, affiliations, knowledge, or beliefs) in the subject matter or materials discussed in this manuscript.

Corresponding Author:

Tjokorda Gde Bagus Mahadewa, Department of Surgery, Division of Neurosurgery, Faculty of Medicine, Udayana University, Denpasar, Bali, Indonesia; tjokmahadewa@unud.ac.id

Published 09 March 2022

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2022 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.