

## Visibility of Prevertebral Soft Tissues in the Neck Using Ultrasonography: A Feasibility Study

O. Kenechi Nwawka, Noah Cha, Lydia M. Ko and Vincent R. LaSala

*Int J Spine Surg* published online 9 October 2023

<https://www.ijssurgery.com/content/early/2023/10/09/8546>

This information is current as of May 4, 2025.

---

**Email Alerts** Receive free email-alerts when new articles cite this article. Sign up at:  
<http://ijssurgery.com/alerts>

# Visibility of Prevertebral Soft Tissues in the Neck Using Ultrasonography: A Feasibility Study

O. KENECHI NAWKA, MD<sup>1</sup>; NOAH CHA, BA<sup>1</sup>; LYDIA M. KO, MPH<sup>1</sup>; AND VINCENT R. LASALA, MD<sup>2</sup>

<sup>1</sup>Department of Radiology and Imaging, Hospital for Special Surgery, New York, NY, USA; <sup>2</sup>Department of Anesthesiology, Hospital for Special Surgery, New York, NY, USA

## ABSTRACT

**Objective:** To aid emergent diagnosis of postoperative retropharyngeal hematoma in anterior cervical spine surgery patients, this study investigates ultrasonography's potential role by evaluating the visibility of retropharyngeal and prevertebral soft tissues in the neck using ultrasound and potential correlations with body habitus.

**Methods:** The visibility of the anterior vertebral bodies and the prevertebral soft tissues in both sides of the neck was recorded and analyzed with 95% CI using the Wilson method. Body mass index, neck circumference, and neck length were measured. A point-biserial correlation was performed to compare body habitus with visibility of vertebrae and prevertebral tissues.

**Results:** Longus colli muscle and C3 to C6 were successfully visualized in all 10 (100%) cases on both sides. C2 was only visible in 6 (60%) on both sides. C7 was visible in 9 (90%) on the right and 7 (70%) on the left. The esophagus was visible in 7 (70%) on the right and 10 (100%) on the left. There was a significant negative correlation with neck circumference and C2 visibility on the right side,  $r(8) = -0.76$ ,  $P = 0.011$ .

**Conclusions:** Ultrasonography was successful in visualizing prevertebral tissues, with a trend of obstructed visibility with wider and longer necks.

**Clinical Relevance:** Ultrasonography has potential to aid early detection of postoperative retropharyngeal hematoma after cervical spine surgery.

**Level of Evidence:** 4.

New Technology


Keywords: retropharyngeal, prevertebral tissues, ultrasound, imaging, anterior cervical spine surgery

## INTRODUCTION

A postoperative retropharyngeal hematoma may lead to airway compromise, a rare but life-threatening complication in patients who have undergone procedures such as anterior cervical discectomy and fusion.<sup>1–3</sup> However, imaging screening protocols for postoperative retropharyngeal hematoma are poorly defined.<sup>4</sup> This potentially life-threatening condition can be difficult to diagnose clinically and by computed tomography (CT), and although magnetic resonance imaging (MRI) can depict the extent of retropharyngeal lesions, it requires more patient cooperation, making it difficult to perform in acutely ill patients who may not have stable airways.<sup>5</sup> Sonography may potentially fill that role in cervical patients, given its portability and high-resolution imaging capabilities. This is a feasibility study with the purpose of evaluating the visibility of retropharyngeal and prevertebral soft tissues in the neck using ultrasonography (US) and determining the potential correlations with the body habitus.

## METHODS

This prospective feasibility study was approved by the institutional review board and compliant with the Health Insurance Portability and Accountability Act. It was conducted between 8 August 2020 and 13 April 2021. Healthy adult volunteers aged 18 years or older were recruited from our institution and underwent US of the neck. All participants provided informed consent. All US examinations were performed and interpreted by a single fellowship-trained musculoskeletal radiologist. Visibility of the anterior vertebral bodies and the prevertebral soft tissues in the neck (esophagus and longus colli muscle [LCM]; Figure) was recorded and analyzed with 95% CIs using the Wilson method. Body mass index, neck circumference, and length were measured in all participants. A point-biserial correlation was performed to compare body habitus measurements (neck circumference and length, LCM thickness and cross-sectional area, and body mass index) with visibility of retropharyngeal and prevertebral tissues in the neck (C2, C3, C4, C5, C6, C7, esophagus, and LCM).



**Figure.** Sonography of the prevertebral soft tissues in the neck. Transverse (A) and longitudinal (B) images of the prevertebral tissues in the cervical spine from the left. The longus colli muscle overlies the anterior/anterolateral vertebral bodies. (C and D) Transverse sonographic images from the right and left of the neck demonstrating visualization of the esophagus only from the left. Abbreviations: L, left; R, right.

The analysis was stratified into right- and left-sided visibility.

## RESULTS

Ten volunteers, 4 men and 6 women, were enrolled. Participants' mean age was 41 years (range 31–65). The mean neck length and circumference were 15.25 cm (SD = 2.4) and 37.55 cm (SD = 2.0), respectively. LCM and C3 to C6 of the anterior cervical vertebral body were successfully visualized in all 10 (100%) cases on both sides. C2 was only visible in 60% of the cases on both sides. C7 was visible in 90% on the right and 70% on the left. The esophagus was visible 100% of the time on the left but only 70% on the right (Table; Figure). There was a significant negative correlation with neck

circumference and C2 visibility on the right side,  $r(8) = -0.76$ ,  $P = 0.011$ . Visibility of left and right C2, right esophagus, and right C7 displayed similar negative correlations with neck circumference and length but were not significant. Because LCM, C3 to C6, and the esophagus (on the left) were visible in all cases, correlations could not be completed due to the constancy of the variable.

## DISCUSSION

In this feasibility study, US was successful in the visualization of anterior vertebral bodies and prevertebral soft tissues in the neck in most cases. There was a trend of obstructed visibility with wider and longer necks, but only C2 level visibility was significantly decreased

**Table.** Visualization of soft tissues in neck with ultrasound imaging.

Soft Tissue	Cases Visualized on Ultrasonography		Proportion (95% CI) <sup>a</sup>	
	Right	Left	Right	Left
Esophagus	7/10	10/10	70% (40%–89%)	100% (72%–100%)
Longus colli muscle	10/10	10/10	100% (72%–100%)	100% (72%–100%)
C2	6/10	6/10	60% (31%–83%)	60% (31%–83%)
C3	10/10	10/10	100% (72%–100%)	100% (72%–100%)
C4	10/10	10/10	100% (72%–100%)	100% (72%–100%)
C5	10/10	10/10	100% (72%–100%)	100% (72%–100%)
C6	10/10	10/10	100% (72%–100%)	100% (72%–100%)
C7	9/10	7/10	90% (60%–98%)	70% (40%–89%)

<sup>a</sup>The proportion (Wilson 95% CI) of cases with successful visualization of the target structure.

in participants with wider necks. Decreased rates of esophageal visualization on the right makes anatomical sense, given its leftward location in the neck. Although visibility for the cervical prevertebral soft tissues using US may not be as good as on MRI or CT, the portability of US allows it to be applied at the bedside. This portability would allow for early diagnosis of complications such as prevertebral hematoma; such early diagnosis would be particularly useful in critically ill patients who are not stable enough to undergo MRI or CT.

After anterior cervical spine surgery, dysphagia is the most common complication.<sup>2,3</sup> However, airway compromise is of the most critical adverse events, caused by retropharyngeal edema or hematoma. Emergent treatment of airway compromise requires stabilization of the airway and identification of the cause of obstruction. Angioedema is the earliest cause of airway compromise, seen within 6 to 12 hours. Retropharyngeal hematoma typically occurs between 6 and 24 hours postoperatively<sup>3</sup> while the patient is in postoperative recovery and usually before the patient is discharged from the hospital. Bedside US evaluation for the presence, size, and location of hematoma could be beneficial in this period, guiding the need for surgical reintervention in real time.

There is a fairly long literature history documenting the use of US in the spine,<sup>6,7</sup> with studies using US to characterize intramedullary lesions<sup>8,9</sup> and studies evaluating sonography to confirm successful cervical spinal decompression.<sup>10,11</sup> Several reports have demonstrated the benefits of US-guided intervention of the cervical spine and nerves.<sup>12,13</sup> The previously published findings and the current study showing feasible sonographic visibility of the prevertebral soft tissues of the neck suggest the potential of US in the diagnosis of postoperative anterior cervical hematoma.

Some limitations of our study are our small sample size and the use of healthy volunteers as study subjects. Future studies should focus on assessing the efficacy of cervical prevertebral US in the acute postoperative period. A musculoskeletal radiologist performed and

interpreted the US examinations in this study, an unlikely scenario in an emergent moment of need. However, there is routine and successful application of emergency US evaluation by nonradiologists in clinical practice among emergency department practitioners.<sup>14–17</sup> Thus, providers on the inpatient wards (eg, spine surgeons and critical care clinicians) can be taught to successfully use the appropriate US techniques.

## CONCLUSION

This feasibility study establishes the viability of US as an imaging modality to visualize soft tissue structures in the anterior neck. The portability and ability of bedside application of US makes it particularly useful for critically ill patients who are not stable enough to undergo MRI or CT. This suggests the use of US in the early detection and intervention of complicated postoperative issues such as retropharyngeal hematoma following anterior cervical spine surgery.

## REFERENCES

1. Ren H, Wang J, Yu L. Retropharyngeal hematoma following anterior cervical spine surgery: lessons from a case report (CARE-compliant). *Medicine (Baltimore)*. 2019;98(38):e17247. doi:10.1097/MD.00000000000017247
2. Yee TJ, Swong K, Park P. Complications of anterior cervical spine surgery: a systematic review of the literature. *J Spine Surg*. 2020;6(1):302–322. doi:10.21037/jss.2020.01.14
3. Debkowska MP, Butterworth JF, Moore JE, Kang S, Appelbaum EN, Zuelzer WA. Acute post-operative airway complications following anterior cervical spine surgery and the role for cricothyrotomy. *J Spine Surg*. 2019;5(1):142–154. doi:10.21037/jss.2019.03.01
4. Muñoz A, Fischbein NJ, de Vergas J, Crespo J, Alvarez-Vincent J. Spontaneous retropharyngeal hematoma: diagnosis by MR imaging. *AJNR Am J Neuroradiol*. 2001;22(6):1209–1211.
5. Debnam JM, Guha-Thakurta N. Retropharyngeal and prevertebral spaces: anatomic imaging and diagnosis. *Otolaryngol Clin North Am*. 2012;45(6):1293–1310. doi:10.1016/j.otc.2012.08.004
6. Ganau M, Syrmos N, Martin AR, Jiang F, Fehlings MG. Intraoperative ultrasound in spine surgery: history, current applications,



future developments. *Quant Imaging Med Surg*. 2018;8(3):261–267. doi:10.21037/qims.2018.04.02

7. Dohrmann GJ, Rubin JM. History of intraoperative ultrasound in neurosurgery. *Neurosurg Clin N Am*. 2001;12(1):155–166.

8. Dohrmann GJ, Rubin JM. Intraoperative ultrasound imaging of the spinal cord: syringomyelia, cysts, and tumors—a preliminary report. *Surg Neurol*. 1982;18(6):395–399. doi:10.1016/0090-3019(82)90169-0

9. Rubin JM, Dohrmann GJ. The spine and spinal cord during neurosurgical operations: real-time ultrasonography. *Radiology*. 1985;155(1):197–200. doi:10.1148/radiology.155.1.3883416

10. Chryssikos T, Stokum JA, Ahmed A-K, et al. Surgical decompression of traumatic cervical spinal cord injury: a pilot study comparing real-time intraoperative ultrasound after laminectomy with postoperative MRI and CT myelography. *Neurosurgery*. 2023;92(2):353–362. doi:10.1227/neu.0000000000002207

11. Imamura H, Iwasaki Y, Hida K, Cho KH, Abe H. Intraoperative spinal sonography in the cervical anterior approach. *Neurol Med Chir (Tokyo)*. 1995;35(3):144–147. doi:10.2176/nmc.35.144

12. Chang K-V, Wu W-T, Özçakar L. Ultrasound-guided interventions of the cervical spine and nerves. *Phys Med Rehabil Clin N Am*. 2018;29(1):93–103. doi:10.1016/j.pmr.2017.08.008

13. Moreno B, Barbosa J. Ultrasound-guided procedures in the cervical spine. *Cureus*. 2021;13(12):e20361. doi:10.7759/cureus.20361

14. Brooks A, Davies B, Smethurst M, Connolly J. Prospective evaluation of non-radiologist performed emergency abdominal ultrasound for haemoperitoneum. *Emerg Med J*. 2004;21(5):e5. doi:10.1136/emj.2003.006932

15. Bhoi S, Sinha TP, Ramchandani R, Kurrey L, Galwankar S. To determine the accuracy of focused assessment with sonography for trauma done by nonradiologists and its comparative analysis with

radiologists in emergency department of a level 1 trauma center of India. *J Emerg Trauma Shock*. 2013;6(1):42–46. doi:10.4103/0974-2700.106324

16. Lee C, Balk D, Schafer J, et al. Accuracy of focused assessment with sonography for trauma (FAST) in disaster settings: a meta-analysis and systematic review. *Disaster Med Public Health Prep*. 2019;13(5–6):1059–1064. doi:10.1017/dmp.2019.23

17. Zanolotti M, Coppa A, Nazerian P, et al. Chest abdominal-focused assessment sonography for trauma during the primary survey in the emergency department: the CA-FAST protocol. *Eur J Trauma Emerg Surg*. 2018;44(6):805–810. doi:10.1007/s00068-015-0620-y

**Funding:** The authors received no financial support for the research, authorship, and/or publication of this article.

**Declaration of Conflicting Interests:** The authors report no conflicts of interest in this work.

**Corresponding Author:** O. Kenechi Nwawka, Department of Radiology and Imaging, Hospital for Special Surgery, 535 E 70th St, New York, NY 10021, USA; nwawkao@hss.edu

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2023 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.