

INTERNATIONAL
JOURNAL
of
SPINE
SURGERY

Demographic Trends in the Use of Intraoperative Neuromonitoring for Scoliosis Surgery in the United States

REMI M. AJIBOYE, HOWARD Y. PARK, JEREMIAH R. COHEN, EVAN E. VELLIOS, ELIZABETH L. LORD, ADEDAYO O. ASHANA, ZORICA BUSER and JEFFREY C. WANG

Int J Spine Surg 2018, 12 (3) 393-398

doi: <https://doi.org/10.14444/5046>

<https://www.ijssurgery.com/content/12/3/393>

This information is current as of May 17, 2025.

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://ijssurgery.com/alerts>

Demographic Trends in the Use of Intraoperative Neuromonitoring for Scoliosis Surgery in the United States

REMI M. AJIBOYE, MD,¹ HOWARD Y. PARK, MD,¹ JEREMIAH R. COHEN, BS,¹ EVAN E. VELLIOS, MD,¹ ELIZABETH L. LORD, MD,¹ ADEDAYO O. ASHANA, MD,¹ ZORICA BUSER, PHD,² JEFFREY C. WANG, MD²

¹UCLA Medical Center, Department of Orthopaedic Surgery, Los Angeles, California, ²Keck Medicine of USC, Department of Orthopaedic Surgery, Los Angeles, California

ABSTRACT

Background: Intraoperative neuromonitoring (ION), such as motor-evoked potential (MEP), somatosensory evoked potentials (SSEP), and electromyography (EMG), is used to detect impending neurological injuries during spinal surgery. To date, little is known about the trends in the use of ION for scoliosis surgery in the United States.

Methods: A retrospective review was performed using the PearlDiver Database to identify patients that had scoliosis surgery with and without ION from years 2005 to 2011. Demographic information (such as age, gender, region within the United States) and clinical information (such as type of ION and rates of neurological injury) were assessed.

Results: There were 3618 patients who had scoliosis surgery during the study period. Intraoperative neuromonitoring was used in 1361 (37.6%) of these cases. The number of cases in which ION was used increased from 27% in 2005 to 46.9% in 2011 ($P < .0001$). Multimodal ION was used more commonly than unimodal ION (64.6% versus 35.4%). The most commonly used modality was combined SSEP and EMG, while the least used modality was MEP only. Neurological injuries occurred in 1.8 and 2.0% of patients that had surgery with and without ION, respectively ($P = .561$). Intraoperative neuromonitoring was used most commonly in patients <65 years of age and in the Northeastern part of the United States (age $P = .006$, region $P < .0001$).

Conclusions: The use of ION for scoliosis surgery gradually increased annually from 2005 to 2011. Age and regional differences were noted with neuromonitoring being most commonly used for scoliosis surgery in nonelderly patients and in the Northeastern part of the United States. No differences were noted in the risk of neurological injury in patients that had surgery with and without ION. Although the findings from this study may seem to suggest that ION may not influence the risk of neurologic injury, this result must be interpreted with caution as inherently riskier surgeries may utilize ION more, leading to an actual reduction in injuries more dramatic than observed in this study.

Other & Special Categories

Keywords: scoliosis, neuromonitoring, motor-evoked potential, somatosensory evoked potential, electromyography

INTRODUCTION

Neurological injuries are known complications of spine surgery. In spinal deformity surgery, the risk of neurological injury is estimated to be from 0.5 to 3%.^{1–7} These injuries are thought to occur from implant-related damages, correction maneuvers, or ischemia.⁸ To decrease the risk of these adverse events, intraoperative neuromonitoring (ION), such as motor-evoked potential (MEP), somatosensory evoked potential (SSEP), and electromyography (EMG), is used to detect impending injury of neural elements. Somatosensory evoked potentials have been used clinically since 1977 and work by monitoring the ascending sensory afferent pathways in the spinal cord.⁹ Motor-evoked potentials work by monitoring peripheral muscle activity from direct stimulation of the motor cortex, while (triggered or

spontaneous) EMGs monitor muscle contractions from nerve root stimulation. Prior to the widespread use of ION, the Stagnara wake-up test served as the only way to assess the functional integrity of the spinal cord intraoperatively.¹⁰ The Stagnara wake-up test is performed by waking a patient up during surgery and checking for gross motor movements. Some surgeons advocate for the adjunctive use of the Stagnara wake-up test when there is no improvement in ION signals despite actions to reverse a suspected intraoperative neurological injury or when reliable ION signals cannot be obtained.¹¹

In 2009, the Scoliosis Research Society (SRS) released an updated position statement stating that ION is the preferred method for early detection of an evolving or impending neurological injury during

deformity surgery.¹² However, the decision to use ION during spinal deformity surgery is often guided by the type of surgery, surgeon choice, and experience, and there is no consensus on the optimal neuromonitoring modality to use. In addition, most of the published studies on the use of ION for scoliosis surgery are from academic centers, and little is known on how neuromonitoring is used in the real world, ie, in academic and nonacademic settings. The goal of this study was to evaluate the trends in the use of neuromonitoring for scoliosis surgery in the United States.

MATERIALS AND METHODS

A retrospective review was performed using the PearlDiver Patient Record Database (<http://www.pearldiver.com>; PearlDiver, Warsaw, Indiana) to search through the patient records within both the Standard Analytical Files (SAF) of Medicare and the United Healthcare (UHC) databases. The PearlDiver database is commercially available and contains de-identified patient data that is Health Insurance Portability and Accountability Act (HIPAA) compliant and allows researchers to construct queries to identify patient groupings that meet specified criteria of interest. The raw datasets are filtered by characteristics such as age group, gender, region of the country, and year. The SAF dataset used in this study spans from 2005 to 2011 and contains more than 40 million patients per year, whereas the UHC set contains 21 million patients with records from 2007 to 2011.

Data Collection

The database was used to identify cases of scoliosis undergoing spinal surgery with neuromonitoring from years 2005 to 2011 using both Current Procedural Terminology (CPT) and International Classification of Diseases, Ninth Revision (ICD-9) codes (see Appendix). Each record provided demographic information (such as age, gender, and region within the United States) and clinical information (such as type of neuromonitoring modality used and rates of neurological injury; see Appendix). Neurologic injury was defined as neurologic weakness within 30 days after the index surgery.

Statistical Analysis

The Stata statistical software version 11.0 (Stata-Corp, College Station, Texas) was used to perform




Figure 1. Percentage of scoliosis surgery performed with neuromonitoring during the study period (2005–2011).

the analyses. The χ^2 test was used to detect any differences in the variables of interest (ie, temporal trends, complications, age, gender, and region). Significance level was set at the $P < .05$.

RESULTS

Neuromonitoring Use in the United States During the Study Period (2005–2011)

During the study period, 3618 patients underwent scoliosis surgery. Overall, neuromonitoring was used in 1361 (37.6%) of these cases. There was a statistically significant steady increase in the use of neuromonitoring for scoliosis surgery from 27% in 2005 to 46.9% in 2011 ($P < .0001$; Figure 1; Table 1).

Type of Neuromonitoring Modality

Out of a total of 1361 patients that had scoliosis surgery with neuromonitoring, multimodal neuromonitoring was used in 64.6% of cases compared to 34.6% with unimodal neuromonitoring. In terms of specific combinations of neuromonitoring, the most commonly used modality was SSEP + EMG, while the least used modality was MEP only (SSEP + EMG = 37.4%, EMG only = 22.9%, SSEP + MEP + EMG = 20.8%, SSEP only = 12.4%, SSEP + MEP = 5.7%, MEP + EMG = 0.8%, and MEP only = 0%; Table 2).

Neurological Injury

Neurological injuries within 30 days from the date of the index surgery occurred in 1.8% (24/1361) and 2.0% (46/2257) of patients that underwent

Table 1. Demographic information of patients from 2005 to 2011.

	Total Number of Scoliosis Surgeries with Neuromonitoring	Total Number of Scoliosis Surgeries	P Value
Year			<.0001
2005	93	345	
2006	123	445	
2007	154	512	
2008	213	563	
2009	258	686	
2010	183	386	
2011	195	416	
Age			.006
<65	255	580	
65–69	372	1002	
70–74	337	931	
75–79	267	741	
80–84	113	352	
>84	30	88	
Gender			.106
Female	935	2542	
Male	421	1062	
Region			<.0001
Midwest	336	919	
Northeast	231	436	
South	640	1516	
West	341	852	
Total*	1361	3618	

*Discrepancies between total value and summation of values in each group are attributed to the transfer of patients between subgroups.

surgery with and without ION, respectively ($P = .561$; Table 3).

Age

Neuromonitoring was used in 44.0% (255/580), 37.1% (372/1002), 36.2% (337/931), 36% (267/741), 32.1% (113/352), and 34% (30/88) of patients in age groups <65, 65–69, 70–74, 75–79, and 80 years and over, respectively ($P = .006$; Table 1).

Gender

Neuromonitoring was used in 36.8% (935/2542) of women compared to 39.6% (421/1062) of men ($P = .106$; Table 1).

Table 2. Types of neuromonitoring modality used for scoliosis surgery.

Type	No.
Unimodal neuromonitoring	
SSEP only	169
MEP only	0
EMG only	313
Multimodal neuromonitoring	
SSEP and MEP	77
SSEP and EMG	509
MEP and EMG	11
SSEP, MEP and EMG	283
Total*	1361

Abbreviations: EMG, electromyography; MEP, motor-evoked potential; SSEP, somatosensory evoked potentials.

*Discrepancies between total value and summation of values in each group are attributed to the transfer of patients between subgroups.

Table 3. Risk of neurological injury after scoliosis surgery with and without neuromonitoring.

	With Neuromonitoring	Without Neuromonitoring	P Value
Risk of neurological injury	24/1361 (1.8%)	46/2257 (2.0%)	.561

Region

Neuromonitoring was used in 53% (231/436) of scoliosis surgery in the Northeastern part of the United States compared to 42.2% (640/1516) in the South, 40.0% (341/852) in the West and 36.6% (336/919) in the Midwest ($P < .0001$; Table 1).

DISCUSSION

The goal of this study was to evaluate the trends in the use of neuromonitoring for scoliosis surgery in the United States. To that end, we found increased utilization rates of neuromonitoring from 2005 to 2011. The vast majority of monitoring was multimodal, and the risk of neurological injury was not significantly altered by its use. Although there were no gender-related differences noted in the utility of neuromonitoring, age and regional differences were noted with neuromonitoring being most commonly used for scoliosis surgery in nonelderly patients and in the Northeastern part of the United States.

Intraoperative neuromonitoring has emerged as a component of the standard of care for scoliosis surgery with data from this study showing that its use has increased from 2005 to 2011. The intuitive reason for the utility of ION is to raise warning against devastating neurologic complications that can be prevented with intervention, such as reducing the degree of distraction, adjusting retractors, removing hardware, and minimizing the length of surgery.¹³ In a retrospective study of 443 194 patients by James et al,¹⁴ the utilization of ION within the United States increased from 1% of all spine procedures in 2007 to 12% in 2011, which are lower than the 37.6% overall utilization rate in scoliosis surgery found in this study. This difference can be accounted for by the inclusion of a wide range of spinal procedures including microdiscectomy in the study by James et al.¹⁴ These procedures traditionally do not utilize ION to the same degree as scoliosis surgery. Furthermore, studies have shown no clear benefit or even recommendation against ION in certain low-risk spinal procedures.^{15,16} In their single institutional study of

4467 neurosurgical procedures performed at Texas Children's Hospital, Vadivelu et al found ION use increased from 2008 to 2011 with surgeon-related factors such as less than 10 years of practice and subspecialty interest in spine positively associated with its use.¹⁷ Although the inclusion criteria of the aforementioned studies may differ, the overall trend of increased ION utilization in the last decade is seemingly well established.

When considering the absolute utilization rate of 37.6% in our study, it may be interpreted as a low especially in comparison to a recent study of 108 419 procedures by members of the SRS Morbidity and Mortality Committee in which ION was used in 65% of cases.¹⁸ There are several factors that account for the differences in utilization rates for ION in both studies. Members of the SRS Morbidity and Mortality Committee are primarily from high-performing academic centers that perform cases of higher complexity. This current study utilizes a national database, which captures practice patterns from both academic and nonacademic centers. In addition, there may be overcoding for the diagnosis of scoliosis in this study, as varying degrees of coronal deformity occur concurrently with lumbar stenosis and other low-risk spinal procedures. Lastly, the lack of availability of ION (especially in nonacademic centers) and the increased cost associated with its use may account for the low overall utilization rate observed in the current study.

With regard to the specific modalities of use, we found that combined SSEP with EMG was the most commonly utilized modality. Unimodal neuromonitoring comprised only 35.3% of monitoring use in our study, which is somewhat comparable to the rate of 22.6% reported by Hamilton et al.¹⁸ Multimodal ION is commonplace, as it can monitor both spinal cord and nerve root function.¹⁹ Multimodal ION has resulted in increases in the sensitivity and specificity for neurologic injury to nearly 100%.¹⁹⁻²²

Neurologic injury within 30 days from the date of index surgery occurred in 1.8 and 2.0% of patients that underwent scoliosis surgery with and without ION, respectively. These rates are within the range of 0.5 to 3% risk of neurological injury reported in the literature for deformity surgeries.¹⁻⁷ Although the findings from this study may seem to suggest that ION may not influence the rate of neurologic injury, this result must be interpreted with caution

as inherently riskier surgeries may utilize neuromonitoring more, leading to an actual reduction in injuries more dramatic than observed in this study. Fu et al reported higher rates of neurologic deficits with ION in pediatric spine cases, which were attributed to the disproportionate use of monitoring in higher risk cases.²³ To this end, no randomized controlled trials have been undertaken to elucidate the true effect of neuromonitoring on neurological injuries following scoliosis surgery.

Our results revealed no difference with regard to gender, but showed a proclivity of ION to be utilized for scoliosis surgery more in nonelderly patients (<65 years of age) and in the Northeastern region of the United States. To our knowledge, this is the first study to review the age- or gender-related differences with respect to ION utilization. James et al examined the regional use of neuromonitoring from 2008 to 2011, and the lowest utilization was noted to be in the Northeastern part of the United States, which conflicts with the results from our study.¹⁴ This difference may stem from the fact that our study exclusively studied scoliosis surgery as opposed to a heterogeneous group of neurosurgical spinal procedures examined by James et al. Within our dataset, the increased utility of ION in the Northeastern region of the United States may be influenced by medicolegal concerns and malpractice premiums in these areas.²⁴ According to a recent report, the top 5 states with the highest medical malpractice payout per capita are New York, New Jersey, Pennsylvania, Massachusetts, and Rhode Island, all of which are located in the Northeastern region of the United States.²⁵ This finding highlights the fact that litigation and malpractice claims in various parts of the United States may have an influence on physicians' pattern of practice.

Limitations

There are some limitations inherent to using an administrative database for research. Inaccuracies in the coding of diagnoses and procedures may have influenced the results of this study. In addition, important detailed clinical information, such as operative time, disease severity, complexity of surgery, intraoperative events, neuromonitoring sensitivity, specificity, false positive and negatives, and information on type and severity of neurological complications, is not recorded in this database. Despite these limitations, this study is valuable because of its large sample size, which makes it

suitable to study national trends, and it captures a heterogeneous sample of the practice patterns of surgeons with regards to ION use for scoliosis surgery in both academic and nonacademic centers. In addition, we believe that the information from this study will help shed light on disparities associated with neuromonitoring usage and/or availability across different regions of the United States, especially in the current political and economic climate whereby health care cost and patient safety are currently being scrutinized.

CONCLUSIONS

The use of neuromonitoring for scoliosis surgery gradually increased annually from 2005 to 2011. No differences were noted in the rates of neurological injury in patients that underwent scoliosis surgery with and without neuromonitoring. Although there were no gender-related differences noted in the utility of neuromonitoring, age and regional differences were noted with neuromonitoring being most commonly used for scoliosis surgery in nonelderly patients and in the Northeastern part of the United States.

REFERENCES

- Eggspuehler A, Sutter MA, Grob D, Jeszenszky D, Dvorak J. Multimodal intraoperative monitoring during surgery of spinal deformities in 217 patients. *Eur Spine J*. 2007;16(Suppl 2):S188–196.
- Diab M, Smith AR, Kuklo TR. Neural complications in the surgical treatment of adolescent idiopathic scoliosis. *Spine (Phila Pa 1976)*. 2007;32(24):2759–2763.
- Kamerlink JR, Errico T, Xavier S, et al. Major intraoperative neurologic monitoring deficits in consecutive pediatric and adult spinal deformity patients at one institution. *Spine (Phila Pa 1976)*. 2010;35(2):240–245.
- Nuwer MR, Dawson EG, Carlson LG, Kanim LE, Sherman JE. Somatosensory evoked potential spinal cord monitoring reduces neurologic deficits after scoliosis surgery: results of a large multicenter survey. *Electroencephalogr Clin Neurophysiol*. 1995;96(1):6–11.
- Dawson EG, Sherman JE, Kanim LE, Nuwer MR. Spinal cord monitoring. Results of the Scoliosis Research Society and the European Spinal Deformity Society survey. *Spine (Phila Pa 1976)*. 1991;16(8 Suppl):S361–364.
- Forbes HJ, Allen PW, Waller CS, et al. Spinal cord monitoring in scoliosis surgery. Experience with 1168 cases. *J Bone Joint Surg Br*. 1991;73(3):487–491.
- Zhuang Q, Wang S, Zhang J, et al. How to make the best use of intraoperative motor evoked potential monitoring? Experience in 1162 consecutive spinal deformity surgical procedures. *Spine (Phila Pa 1976)*. 2014;39(24):E1425–1432.
- Malhotra NR, Shaffrey CI. Intraoperative electrophysiological monitoring in spine surgery. *Spine (Phila Pa 1976)*. 2010;35(25):2167–2179.
- Nash CL, Jr., Lorig RA, Schatzinger LA, Brown RH. Spinal cord monitoring during operative treatment of the spine. *Clin Orthop Relat Res*. 1977;(126):100–105.
- Vauzelle C, Stagnara P, Jouvinroux P. Functional monitoring of spinal cord activity during spinal surgery. *Clin Orthop Relat Res*. 1973;(93):173–178.
- Leong JJ, Curtis M, Carter E, Cowan J, Lehovsky J. Risk of neurological injuries in spinal deformity surgery. *Spine (Phila Pa 1976)*. 2016;41(12):1022–1027.
- Scoliosis Research Society. Position statements: neuromonitoring information statement. January 2009; <https://www.srs.org/about-srs/quality-and-safety/position-statements/neuromonitoring-information-statement>. Accessed July 1, 2016.
- Nuwer MR, Emerson RG, Galloway G, et al. Evidence-based guideline update: intraoperative spinal monitoring with somatosensory and transcranial electrical motor evoked potentials*. *J Clin Neurophysiol*. 2012;29(1):101–108.
- James WS, Rughani AI, Dumont TM. A socioeconomic analysis of intraoperative neurophysiological monitoring during spine surgery: national use, regional variation, and patient outcomes. *Neurosurg Focus*. 2014;37(5):E10.
- Resnick DK, Choudhri TF, Dailey AT, et al. Guidelines for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 15: electrophysiological monitoring and lumbar fusion. *J Neurosurg Spine*. 2005;2(6):725–732.
- Gundanna M, Eskenazi M, Bendo J, Spivak J, Moskovich R. Somatosensory evoked potential monitoring of lumbar pedicle screw placement for in situ posterior spinal fusion. *Spine J*. 2003;3(5):370–376.
- Vadivelu S, Sivaganesan A, Patel AJ, et al. Practice trends in the utilization of intraoperative neurophysiological monitoring in pediatric neurosurgery as a function of complication rate, and patient-, surgeon-, and procedure-related factors. *World Neurosurg*. 2014;81(3–4):617–623.
- Hamilton DK, Smith JS, Sansur CA, et al. Rates of new neurological deficit associated with spine surgery based on 108,419 procedures: a report of the scoliosis research society morbidity and mortality committee. *Spine (Phila Pa 1976)*. 2011;36(15):1218–1228.
- Pelosi L, Lamb J, Grevitt M, Mehdian SM, Webb JK, Blumhardt LD. Combined monitoring of motor and somatosensory evoked potentials in orthopaedic spinal surgery. *Clin Neurophysiol*. 2002;113(7):1082–1091.
- Sutter MA, Eggspuehler A, Grob D, Porchet F, Jeszenszky D, Dvorak J. Multimodal intraoperative monitoring (MIOM) during 409 lumbosacral surgical procedures in 409 patients. *Eur Spine J*. 2007;16(Suppl 2):S221–S228.
- Quraishi NA, Lewis SJ, Kelleher MO, Sarjeant R, Rampersaud YR, Fehlings MG. Intraoperative multimodality monitoring in adult spinal deformity: analysis of a prospective series of one hundred two cases with independent evaluation. *Spine (Phila Pa 1976)*. 2009;34(14):1504–1512.
- Thuet ED, Winscher JC, Padberg AM, et al. Validity and reliability of intraoperative monitoring in pediatric spinal deformity surgery: a 23-year experience of 3436 surgical cases. *Spine (Phila Pa 1976)*. 2010;35(20):1880–1886.
- Fu KM, Smith JS, Polly DW, et al. Morbidity and mortality associated with spinal surgery in children: a review of

the Scoliosis Research Society morbidity and mortality database. *J Neurosurg Pediatr.* 2011;7(1):37–41.

24. The Medical Liability Monitor, Annual Survey, Annual Rate Survey. 2013.

25. Gower J. *Medical Malpractice Payout Analysis.* Carbondale, IL: Diederich Healthcare; 2015.

Disclosures and COI: No funds were received in support of this work. No relevant financial disclosures are associated with this work.

Corresponding Author: Remi M. Ajiboye,

MD, UCLA Department of Orthopaedic Surgery, 10833 LeConte Avenue, 76-119 CHS, Los Angeles, CA 90095-6902. Phone: (510) 828-7906; Fax: (310) 825-1311; Email: Remi.Ajiboye@gmail.com.

Published 15 August 2018

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2018 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.

Appendix. Current procedural terminology (CPT) and International Classification of Diseases, Ninth Revision (ICD-9) codes used to identify cases of scoliosis undergoing spinal surgery with neuromonitoring.

Code	Description
CPT: Scoliosis Surgery	
22800	Arthrodesis posterior for spinal deformity with or without cast up to 6 vertebral segments
22802	Posterior 7 to 12 vertebral segments
22804	Posterior 13 or more vertebral segments
22808	Arthrodesis, anterior, for spinal deformity, with or without cast; 2 to 3 vertebral segments
22810	Arthrodesis, anterior, for spinal deformity, with or without cast; 4 to 7 vertebral segments
22812	Arthrodesis, anterior, for spinal deformity, with or without cast; 8 or more vertebral segments
22840	Posterior non-segmental instrumentation (eg, Harrington rod technique, pedicle fixation across one interspace atlantoaxial, transarticular screw, fixation, sublaminar wiring at C1, facet screw fixation)
22841	Wiring
22842	Posterior segmental instrumentation (eg, pedicle fixation, dual rods with multiple hooks and sublaminar wires); 3 to 6 vertebral segments)
22843	Posterior segmental instrumentation (eg, pedicle fixation, dual rods with multiple hooks and sublaminar wires); 7 to 12 vertebral segments)
22844	Posterior segmental instrumentation (eg, pedicle fixation, dual rods with multiple hooks and sublaminar wires); 13 or more vertebral segments)
22845	Anterior instrumentation; 2 to 3 vertebral segments
22846	Anterior instrumentation; 4 to 7 vertebral segments
22847	Anterior instrumentation; 8 or more vertebral segments
22848	Pelvic fixation (attachment of caudal end of instrumentation to pelvic bony structures) other than sacrum
ICD-9: Scoliosis	
737.30	Scoliosis (and kyphoscoliosis), idiopathic
737.31	Resolving infantile idiopathic scoliosis
737.32	Progressive infantile idiopathic scoliosis
737.33	Scoliosis due to radiation
737.34	Thoracogenic scoliosis
737.39	Other kyphoscoliosis and scoliosis
737.43	Scoliosis associated with other conditions
754.2	Congenital scoliosis
CPT: Neuromonitoring	
95925, 95926, 95927	Somatosensory-evoked potential
95928, 95929	Motor-evoked potential
95860, 95861, 95862, 95863, 95864, 95869, 95870, 95872	Electromyography
ICD-9: Neurologic Complications	
997.0	Nervous system complications
997.00	Nervous system complication, unspecified
997.01	Central nervous system complication
997.09	Other nervous system complications