

Perioperative Change in Cervical Lordosis and Health-Related Quality-of-Life Outcomes

Jose A. Canseco, Brian A. Karamian, Parthik D. Patel, Michael Markowitz, Joseph K. Lee, Mark F. Kurd, D. Greg Anderson, Jeffrey A. Rihn, Alan S. Hilibrand, Christopher K. Kepler, Alexander R. Vaccaro and Gregory D. Schroeder

Int J Spine Surg 2022, 16 (6) 960-968

doi: <https://doi.org/10.14444/8325>

<http://ijssurgery.com/content/16/6/960>

This information is current as of April 16, 2024.

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://ijssurgery.com/alerts>

Perioperative Change in Cervical Lordosis and Health-Related Quality-of-Life Outcomes

JOSE A. CANSECO, MD, PhD¹; BRIAN A. KARAMIAN, MD¹; PARTHIK D. PATEL, MD¹; MICHAEL MARKOWITZ, DO¹; JOSEPH K. LEE, MD¹; MARK F. KURD, MD¹; D. GREG ANDERSON, MD¹; JEFFREY A. RIHN, MD¹; ALAN S. HILIBRAND, MD¹; CHRISTOPHER K. KEPLER, MD, MBA¹; ALEXANDER R. VACCARO, MD, PhD, MBA¹; AND GREGORY D. SCHROEDER, MD¹

¹Rothman Orthopaedic Institute at Thomas Jefferson University, Philadelphia, PA, USA

ABSTRACT

Background: Surgeons have scrutinized spinal alignment and its impact on improving clinical outcomes following anterior cervical discectomy and fusion (ACDF). The primary analysis of this study examines the relationship between change in perioperative cervical lordosis (CL) and health-related quality-of-life (HRQOL) outcomes after ACDF. Secondary analysis evaluates the effects of fusion construct length on outcomes in patients grouped by preoperative cervical alignment.

Methods: A retrospective cohort study was performed on an institutional database including patients who underwent 1- to 3-level ACDF. C2-C7 CL was measured preoperatively and at final follow-up. For primary analysis, patients were classified based on their perioperative cervical lordotic correction: (1) kyphotic, (2) maintained, and (3) restored. For secondary analysis, patients were categorized based on their preoperative C2-C7 CL: (1) kyphotic, (2) neutral, and (3) lordotic. Demographics and perioperative change in patient-reported outcome measures were compared between groups.

Results: A total of 308 patients were included. A significant difference was noted among maintained, restored, and kyphotic groups in terms of delta physical compositeshort form-12 score (Δ PCS-12) (9.0 vs 10.3 vs 1.5; $P = 0.04$) and delta visual analog scale score (Δ VAS) for arm pain (−0.9 vs −3.8 vs −0.6; $P = 0.03$). Regression analysis revealed significantly greater improvement of PCS-12 (β : 8.6; $P = 0.03$) and VAS arm (β : −2.0; $P = 0.03$) scores in restored patients compared with kyphotic patients. The length of fusion construct in patients grouped by preoperative cervical alignment had no significant impact on the clinical outcomes on regression analysis.

Conclusions: Significantly greater PCS-12 and VAS arm improvement were seen in patients whose cervical sagittal alignment was restored to neutral/lordotic compared with those who remained kyphotic. Multivariate analysis demonstrated no association between construct length and perioperative outcomes.

Clinical Relevance: The results of this study highlight the importance of sagittal alignment and restoration of CL after short-segment ACDF. Irrespective of preoperative sagittal alignment, the length of ACDF fusion construct does not have a significant impact on clinical outcomes.

Level of Evidence: 3.

Cervical spine

Keywords: cervical lordosis, kyphosis, patient-reported clinical outcomes (PROMs), anterior cervical discectomy and fusion (ACDF), alignment, HRQOL, construct length

INTRODUCTION

Anterior cervical discectomy and fusion (ACDF) is one of the most commonly performed spine procedures, with a nonlinear increase of 5.7% (120,617–127,500) in ACDF cases from 2006 to 2013.¹ Prior studies have overwhelmingly shown the efficacy of ACDF in improving clinical outcomes and radiographic fusion rates for various pathologies, including symptomatic cervical spondylosis, moderate-to-severe disc herniations, and cervical deformity.^{2–5}

Debusset’s “cone of economy” theory describes a stable region of spine alignment over which the patient expends minimal muscle energy in order to maintain

an upright balance.⁶ While the majority of current literature describes the importance of sagittal lumbosacral alignment on clinical outcomes, the effect of Debusset’s theory on the cervical spine has ignited recent academic interest. A previous investigation of patients with adult spinal deformity demonstrated that mild positive cervical sagittal imbalance resulted in patients reporting significantly worse health-related quality-of-life (HRQOL) outcomes.⁷ Furthermore, a retrospective study in 235 asymptomatic volunteers found a 37% prevalence rate of cervical kyphosis, which was associated with significantly lower HRQOL outcome scores compared with volunteers with cervical lordosis (CL).⁸

Pre- and postoperative cervical alignment is important in the modern health care era with increasing emphasis on value-based care dependent on improved patient outcomes. The primary aim of this study was to explore the effects of perioperative correction of CL on HRQOL outcomes. The secondary aim was to examine the relationship between fusion construct length and HRQOL outcomes in patients grouped by preoperative cervical alignment.

METHODS

Upon obtaining Institutional Review Board (No. 19D.508) approval, a retrospective cohort study was conducted in patients who underwent a 1- to 3-level ACDF between January 2013 and December 2017 at a single academic medical center. A waiver was granted for patient-informed consent as a minimal risk research study. All procedures were performed by 1 of 7 fellowship-trained spine surgeons. Patients were identified via a standardized query language search using the following Common Procedural Terminology codes: 22551, 22552, 22853, 22859, and 22845. Study inclusion criteria were as follows: (1) patients aged ≥ 18 years and (2) patients who underwent a primary ACDF procedure. Patients were excluded if they (1) were younger than 18 years; (2) had surgical indications including infection, malignancy, or trauma; (3) underwent a related revision surgery; or (4) had combined anterior/posterior cervical fusion procedures.

Patient Demographics, Surgical Characteristics, and HRQOL Outcomes

Patient demographic data and surgical case characteristics were obtained via standardized query language search and manual chart review. Demographic data of interest included age, sex, body mass index (BMI), smoking status, and Charlson Comorbidity Index (CCI). Surgical case characteristics of interest included preoperative diagnosis, number of levels fused, and length of follow-up.

HRQOL outcomes were collected in person during clinic appointments using an electronic tablet. HRQOL data were retrieved from the OBERD software system (Columbia, MO, USA) in the form of Short Form-12 Physical Composite Score (PCS-12) and Mental Composite Score (MCS-12), Neck Disability Index (NDI), and visual analog scale (VAS) arm and back scores.

Preoperative CL, Length of Fusion Construct, and HRQOL Outcomes

Radiographic parameters were measured on a Sectra Workstation IDS7 Version 21.1 (Sectra AB; Linköping, Sweden) by a single observer. C2-C7 CL measurement was performed using standing lateral cervical radiographs before the operation and at last postoperative follow-up. C2-C7 cervical sagittal alignment was measured as the angle between the inferior endplate of C2 and inferior endplate of C7. Given prior studies demonstrating C2-C7 CL $>10^\circ$ for asymptomatic individuals, patients were classified into 1 of 3 groups based on the preoperative C2-C7 cervical sagittal alignment: kyphotic, neutral, and lordotic.⁹⁻¹¹ Patients were classified into each group based on the following measurements:

1. Kyphotic: C2-C7 CL $<0^\circ$
2. Neutral: $0^\circ \leq$ C2-C7 CL $<10^\circ$
3. Lordotic: C2-C7 CL $\geq 10^\circ$

HRQOL outcomes within each of the 3 aforementioned groups were compared based on the number of levels fused.

Perioperative Change of CL and HRQOL Outcomes

Patients were classified into 1 of 3 groups based on the change in C2-C7 cervical sagittal alignment from the preoperative to postoperative time state: “kyphotic,” “maintained,” or “restored” groups. Patients were classified into each group based on the following criteria:

1. Kyphotic: Patients with a preoperative kyphotic C2-C7 CL to postoperative kyphotic C2-C7 CL.
2. Maintained: Patients with a preoperative neutral/lordotic C2-C7 CL to postoperative neutral/lordotic C2-C7 CL.
3. Restored: Patients with a preoperative kyphotic C2-C7 CL to postoperative neutral/lordotic C2-C7 CL.

No patients within the cohort were found to have a C2-C7 CL change from preoperative straight/lordotic to postoperative kyphotic. HRQOL outcomes were compared between groups.

Statistical Analysis

Standard descriptive statistics, including proportions, means/medians, and 95% CIs/interquartile ranges, were reported for patient demographic data, follow-up, functional outcomes, and radiographic outcomes

(subsidence and alignment data). Normally distributed data were compared using parametric tests and reported as means and 95% CIs, while non-normally distributed (skewed) data were compared using nonparametric tests and reported as medians with interquartile ranges. Sample means between the 3 groups were compared using a parametric analysis of variance test or a non-parametric Kruskal-Wallis test. Categorical data were compared using Pearson’s χ^2 test. Multivariate linear regression analysis was performed to determine the effect of the correction of C2-C7 CL, as well as fusion construct length in groups based on preoperative CL, on HRQOL outcome after controlling for age, BMI, CCI, and perioperative diagnosis. For all analyses, *P* values less than 0.05 were considered statistically significant. All statistical analyses were performed using SPSS Statistics version 26 (IBM Corporation, Armonk, NY).

RESULTS

Baseline Characteristics

A total of 308 patients were included in the final analysis. The median age for the cohort was 57.0 (49.0, 64.0) years with the majority of patients being men (52.9%). Of the total patient population, 204 (66.2%) patients reported never having smoked, while 72 (23.4%) patients reported being former smokers, and 32 (10.4%) reported being current smokers. Univariate analysis comparing maintained, restored, and kyphotic group patients showed a significant difference in sex (*P*

= 0.02), with a higher proportion of kyphotic patients being women (66.7% vs 33.3%) and a higher proportion of restored patients being men (61.4% vs 38.6%) (Table 1). Furthermore, there was a significant difference in the 3 groups in terms of preoperative (8.9 vs -5.6 vs -7.2; *P* < 0.001) and postoperative (12.3 vs 8.2 vs -6.2; *P* < 0.001) degrees of C2-C7 CL (maintained, restored, and kyphotic groups, respectively). Example pre- and postoperative x-ray images for each group are presented in the Figure. No significant differences were present in age, BMI, smoking status, CCI, preoperative diagnosis, levels fused, or follow-up duration between groups.

Perioperative Change of CL and HRQOL Outcomes

A significant difference was appreciated among maintained, restored, and kyphotic patients in PCS-12 postoperative (41.6 vs 45.8 vs 36.6, respectively, *P* = 0.02) and delta (9.0 vs 10.3 vs 1.5, respectively, *P* = 0.04) scores, with the restored cohort reporting the highest postoperative scores and delta improvement (Table 2). Similarly, there was a significant difference in VAS arm preoperative (3.3 vs 5.3 vs 2.8, respectively, *P* = 0.02) and delta (-0.9 vs -3.8 vs -0.6, respectively, *P* = 0.03) scores, with patients in the restored cohort reporting the highest preoperative arm pain scores but the greatest perioperative improvement. Regression analysis showed significantly greater improvement

Table 1. Baseline characteristics.

Characteristic	Maintained (n = 177)	Restored (n = 101)	Kyphotic (n = 30)	<i>P</i> Value ^{a,b}
Age	57.0 (59.5, 65.0)	57.0 (49.0, 63.0)	52.5 (45.5, 68.7)	0.18
Sex				<u>0.02</u>
Female	86 (48.6%)	39 (38.6%)	20 (66.7%)	
Male	91 (51.4%)	62 (61.4%)	10 (33.3%)	
Body mass index	28.7 (25.6, 32.6)	29.5 (27.1, 34.7)	27.7 (24.9, 31.4)	0.07
Smoking status				0.59
Never smoker	117 (66.1%)	64 (63.4%)	23 (76.7%)	
Former smoker	39 (22.0%)	27 (26.7%)	6 (20.0%)	
Current smoker	21 (11.9%)	10 (9.9%)	1 (3.3%)	
Charlson Comorbidity Index	2.0 (1.0, 3.0)	2.0 (1.0, 3.0)	1.0 (0.7, 2.0)	0.17
Preoperative diagnosis				0.52
Radiculopathy	90 (50.8%)	49 (48.5%)	19 (63.4%)	
Myelopathy	73 (41.2%)	46 (45.5%)	10 (33.3%)	
Myeloradiculopathy	14 (7.9%)	6 (5.9%)	1 (3.3%)	
Levels fused				0.28
1	58 (32.8%)	35 (34.7%)	8 (26.7%)	
2	58 (32.8%)	42 (41.6%)	10 (33.3%)	
3	61 (34.5%)	24 (23.8%)	12 (40.0%)	
Preoperative C2-C7 Cobb angle	8.9 (7.5, 9.7)	-5.6 (-6.5, -4.8)	-7.2 (-9.0, -5.4)	<u><0.001</u>
Postoperative C2-C7 Cobb angle	12.3 (13.3, 11.2)	8.2 (4.6, 13.8)	-6.2 (-8.5, -5.2)	<u><0.001</u>
Follow-up	14.6 (12.1, 18.3)	14.8 (12.6, 19.5)	16.2 (12.0, 21.1)	0.28

Note: Significance achieved at *P* < 0.05.

^aIndependent samples *t* test or Mann-Whitney *U* test for continuous variables.

^bPearson χ^2 test for categorical variables.




Figure. Example pre- and postoperative lateral x-ray images for each group.

in restored patients compared with kyphotic patients in PCS-12 (β : 8.6; $P = 0.03$) and VAS arm (β : -2.0; $P = 0.03$) scores (Table 3). No further significant differences were observed between groups in any of the other HRQOL outcomes. Patients within each group showed a significant improvement from the preoperative to postoperative period, with the exception of Short

Form-12 Mental Composite Scores of the restored ($P = 0.22$) and kyphotic ($P = 0.07$) groups.

Preoperative CL, Length of Fusion Construct, and HRQOL Outcomes

There were 130 patients with preoperative cervical kyphosis (kyphotic), 114 patients with preoperative neutral cervical alignment (neutral), and 64 patients with preoperative CL (lordotic).

Patients in the preoperative kyphotic group demonstrated a significant difference in VAS neck postoperative scores among patients who underwent a 1-, 2-, and 3-level ACDF, with 3-level ACDF patients reporting the worst scores (3.0 vs 1.5 vs 1.6, respectively, $P = 0.01$) (Table 4). However, univariate and regression analysis failed to show a significant difference in perioperative improvement in VAS neck postoperative scores among the 3 groups. No other significant differences between fusion construct lengths were observed in any of the other 4 HRQOL outcomes.

Patients in the preoperative neutral group demonstrated a significant difference in VAS arm preoperative (4.7 vs 6.1 vs 2.5, $P < 0.001$) and delta (-2.5 vs -4.2 vs -0.9, $P = 0.003$) scores among patients who underwent a 1-, 2-, and 3-level ACDF, respectively. Similar differences were seen in VAS neck postoperative (0.9 vs 1.7 vs 2.9) and delta (-3.9

Table 2. Health-related quality-of-life outcomes stratified by postoperative correction of cervical lordosis.

Outcome Measure	Maintained (n = 177)	Restored (n = 101)	Kyphotic (n = 30)	P Value ^a
Physical component of short form-12				
Preoperative	32.6 (30.2, 35.0)	35.5 (32.0, 39.0)	35.0 (28.1, 41.9)	0.32
Postoperative	41.6 (38.4, 44.8)	45.8 (41.9, 49.6)	36.6 (29.1, 44.1)	0.02
Delta	9.0 (5.4, 12.6)	10.3 (7.0, 13.5)	1.5 (-7.8, 10.8)	0.04
P value	<0.001	<0.001	0.01	
Mental component of short form-12				
Preoperative	49.0 (45.3, 52.7)	46.7 (43.0, 50.3)	49.1 (41.7, 56.5)	0.86
Postoperative	50.8 (47.2, 54.3)	45.2 (38.8, 51.6)	51.4 (43.5, 59.3)	0.73
Delta	1.8 (-2.8, 6.3)	-1.4 (-7.7, 4.9)	2.3 (-5.8, 10.3)	0.68
P value	0.04	0.22	0.07	
Neck Disability Index				
Preoperative	36.8 (31.6, 42.0)	43.0 (34.7, 51.4)	35.6 (28.8, 42.4)	0.65
Postoperative	23.8 (17.6, 30.2)	28.0 (18.2, 37.8)	17.4 (8.8, 26.0)	0.65
Delta	-13.0 (-19.8, -6.1)	-14.8 (-27.0, -2.6)	-18.2 (-25.8, -10.5)	0.89
P value	<0.001	<0.001	0.002	
VAS arm				
Preoperative	3.3 (2.1, 4.4)	5.3 (3.8, 6.8)	2.8 (0.9, 4.9)	0.02
Postoperative	2.3 (1.3, 3.3)	1.5 (0.6, 2.4)	2.3 (0.6, 4.1)	0.93
Delta	-0.9 (-2.0, 0.1)	-3.8 (-5.1, -2.6)	-0.6 (-2.8, 1.5)	0.03
P value	<0.001	<0.001	0.01	
VAS neck				
Preoperative	5.1 (4.1, 6.1)	5.3 (4.0, 6.6)	6.4 (5.4, 7.3)	0.31
Postoperative	2.2 (1.3, 3.1)	2.0 (0.7, 3.2)	3.5 (1.0, 6.1)	0.23
Delta	-2.9 (-4.3, -1.5)	-3.3 (-5.0, -1.7)	-2.8 (-5.2, -0.4)	0.26
P value	<0.001	<0.001	0.002	

Abbreviation: VAS, visual analog scale.

Note: Significance achieved at $P < 0.05$.

^aMultiple linear regression analysis was done between groups controlling for age, body mass index, Charlson Comorbidity Index, and preoperative diagnosis.

Table 3. Regression analysis comparing correction of cervical lordosis and health-related quality-of-life outcomes.

Outcome Measure	β Coefficient	95% CI	P Value ^a
Physical component of short form-12			
Maintained : restored	-1.5	(-5.8, 2.7)	0.19
Restored : kyphotic	8.6	(3.9, 13.3)	0.03
Maintained : kyphotic	4.2	(-0.7, 9.1)	0.17
Mental component of short form-12			
Maintained : restored	1.1	(-2.9, 5.0)	0.50
Restored : kyphotic	-3.2	(-9.7, 3.4)	0.54
Maintained : kyphotic	-2.2	(-8.5, 3.9)	0.76
Neck Disability Index			
Maintained : restored	1.7	(-8.6, 5.2)	0.71
Restored : kyphotic	5.6	(-5.8, 17.2)	0.82
Maintained : kyphotic	7.6	(-3.2, 18.5)	0.62
VAS arm			
Maintained : restored	1.9	(-0.1, 2.9)	0.27
Restored : kyphotic	-2.0	(-3.7, -0.4)	0.03
Maintained : kyphotic	-1.2	(-2.7, 0.4)	0.11
VAS neck			
Maintained : restored	0.3	(-1.4, 1.9)	0.22
Restored : kyphotic	-0.5	(-1.5, 0.5)	0.12
Maintained : kyphotic	-0.8	(-2.3, 0.8)	0.33

Abbreviation: VAS, visual analog scale.

Note: Significance achieved at $P < 0.05$.

^aRegression analysis controlled for age, body mass index, Charlson Comorbidity Index, and preoperative diagnosis.

vs -5.2 vs -2.5) scores, as well as PCS-12 postoperative (44.8 vs 39.1 vs 38.8, $P = 0.02$) (Table 5). However, regression analysis failed to demonstrate a significant relationship between length of fusion construct and HRQOL outcomes.

Patients in the preoperative lordotic group demonstrated a significant difference in VAS arm preoperative (6.4 vs 6.7 vs 2.9, $P = 0.01$) and delta (-4.2 vs

-5.5 vs -1.3, $P = 0.01$) scores among patients who underwent a 1-, 2-, and 3-level ACDF, respectively (Table 6). Similarly, a significant difference was demonstrated in VAS neck postoperative (0.6 vs 1.0 vs 2.2, $P = 0.04$) scores. However, regression analysis failed to demonstrate a significant relationship between the length of fusion construct and HRQOL outcomes.

Table 4. Health-related quality-of-life outcomes in patients with preoperative cervical lordosis less than 0° (kyphotic).

Outcome Measure	1-Level (n = 43)	2-Level (n = 51)	3-Level (n = 36)	P Value	Regression Analysis ^a
Physical component of short form-12					
Preoperative	34.4(31.1, 37.8)	36.0 (33.0, 38.9)	34.4 (31.4, 37.3)	0.87	β : 0.03 (-1.96, 2.02) $P = 0.98$
Postoperative	42.4 (39.2, 45.7)	42.4 (39.3, 45.5)	42.4 (39.6, 45.2)	0.95	
Delta	7.8 (5.5, 10.0)	7.9 (5.5, 10.3)	7.5 (4.0, 11.0)	0.99	
P value	<0.001	<0.001	<0.001		
Mental component of short form-12					
Preoperative	50.5 (47.2, 53.7)	50.8 (47.8, 53.8)	48.1 (44.3, 51.9)	0.52	β : 1.06 (-2.04, 4.16) $P = 0.50$
Postoperative	51.9 (48.5, 55.4)	49.9 (46.3, 53.5)	52.6 (49.0, 56.3)	0.53	
Delta	1.5 (-1.9, 5.0)	0.8 (-3.6, 5.1)	3.9 (-1.3, 9.0)	0.65	
P value	0.35	0.55	0.08		
Neck Disability Index					
Preoperative	39.7 (32.8, 46.5)	33.4 (27.7, 39.1)	38.4 (33.1, 43.8)	0.37	β : 0.76 (-7.89, 9.41) $P = 0.86$
Postoperative	35.8 (15.6, 56.0)	22.3 (15.3, 29.4)	23.3 (16.2, 30.3)	0.39	
Delta	-15.5 (-29.1, -2.0)	-16.7 (-28.6, -4.9)	-15.6 (-27.1, -4.0)	0.52	
P value	0.01	0.01	0.001		
VAS arm					
Preoperative	4.6 (3.7, 5.5)	5.0 (4.1, 5.9)	4.2 (2.6, 5.8)	0.62	β : 0.37 (-0.40, 1.14) $P = 0.34$
Postoperative	2.4 (1.5, 3.2)	2.5 (1.8, 3.3)	1.8 (1.0, 2.6)	0.33	
Delta	-3.1 (-3.9, -2.4)	-3.1 (-4.1, -2.2)	-2.4 (-4.2, -0.7)	0.60	
P value	<0.001	<0.001	0.01		
VAS neck					
Preoperative	4.5 (3.5, 5.4)	5.0 (3.9, 6.1)	5.8 (4.7, 7.0)	0.29	β : 0.31 (-0.61, 1.23) $P = 0.50$
Postoperative	1.5 (0.7, 2.2)	1.6 (0.8, 2.4)	3.0 (1.8, 4.2)	0.01	
Delta	-3.4 (-4.5, -2.2)	-3.1 (-4.2, -2.0)	-2.7 (-4.4, -0.9)	0.76	
P value	<0.001	<0.001	0.01		

Abbreviation: VAS, visual analog scale.

Note: Significance achieved at $P < 0.05$.

^aRegression analysis controlled for age, body mass index, Charlson Comorbidity Index, and preoperative diagnosis.

Table 5. Health-related quality-of-life outcomes in patients with preoperative cervical lordosis between 0° and 10° (neutral).

Outcome Measure	1-Level (n = 40)	2-Level (n = 42)	3-Level (n = 32)	P Value	Regression Analysis ^a
Physical component of short form-12					
Preoperative	34.5 (30.6, 38.4)	32.7 (30.4, 35.0)	35.2 (32.2, 38.1)	0.43	β : -2.91 (-6.26, 0.43) $P = 0.09$
Postoperative	44.8 (41.0, 48.6)	39.1 (35.9, 42.2)	38.8 (35.7, 42.0)	0.02	
Delta	11.3 (5.8, 16.7)	6.8 (3.2, 10.5)	5.1 (1.3, 8.9)	0.16	
P value	<0.001	0.001	0.02		
Mental component of short form-12					
Preoperative	49.3 (44.4, 54.1)	52.6 (48.7, 56.5)	49.8 (45.7, 53.9)	0.27	β : -0.02 (-3.86, 3.83) $P = 0.99$
Postoperative	52.3 (48.1, 56.4)	52.2 (49.0, 55.4)	50.0 (45.7, 54.3)	0.40	
Delta	2.3 (-3.6, 8.3)	0.2 (-3.1, 3.4)	2.4 (-4.1, 9.1)	0.84	
P value	0.49	0.85	0.76		
Neck Disability Index					
Preoperative	44.2 (36.6, 51.8)	43.4 (36.5, 50.3)	34.7 (29.4, 40.0)	0.11	β : 3.40 (-2.81, 9.60) $P = 0.27$
Postoperative	26.5 (6.1, 47.0)	52.4 (15.2, 79.3)	22.6 (17.0, 28.2)	0.09	
Delta	-23.6 (-34.9, -12.3)	-14.9 (-55.5, 25.8)	-14.2 (-20.1, -8.2)	0.35	
P value	0.027	0.29	<0.001		
VAS arm					
Preoperative	4.7 (3.7, 5.7)	6.1 (5.2, 7.1)	2.5 (1.1, 3.9)	<0.001	β : 0.50 (-0.47, 1.47) $P = 0.31$
Postoperative	2.4 (1.4, 3.5)	3.2 (2.0, 4.4)	2.4 (1.4, 3.5)	0.56	
Delta	-2.5 (-3.6, -1.4)	-4.2 (-5.4, -2.9)	-0.9 (-2.4, 0.5)	0.003	
P value	0.001	<0.001	0.55		
VAS neck					
Preoperative	4.8 (3.7, 5.9)	6.5 (5.4, 7.4)	4.8 (3.3, 6.2)	0.10	β : 0.43 (-0.59, 1.45) $P = 0.40$
Postoperative	0.9 (0.3, 1.6)	1.7 (0.7, 2.7)	2.9 (1.8, 4.0)	0.001	
Delta	-3.9 (-5.1, -2.8)	-5.2 (-6.5, -3.9)	-2.5 (-4.2, -0.7)	0.03	
P value	<0.001	<0.001	0.07		

Abbreviation: VAS, visual analog scale.

Note: Significance achieved at $P < 0.05$.

^aRegression analysis controlled for age, body mass index, Charlson Comorbidity Index, and preoperative diagnosis.

DISCUSSION

ACDF is an increasingly common procedure performed in the United States.^{12,13} Cervical sagittal alignment and the importance of restoration of proper cervical alignment have gained significant interest recently.^{14–16} Lordosis of the cervical spine provides patients with proper motor function and balance; however, degenerative changes over time can result in loss of CL.^{17,18} As a result, patients can experience myelopathy or radiculopathy leading to disability and inability to perform activities of daily living. The present study investigated the effects of perioperative change of CL on HRQOL outcomes, demonstrating significantly greater improvement in outcomes for patients whose cervical sagittal alignment was restored to neutral/lordotic vs compared with those who remained kyphotic.

Our study showed significantly greater PCS-12 and VAS arm improvement in patients whose cervical sagittal alignment was restored to neutral/lordotic vs those who remained kyphotic postoperatively. In contrast, a previous study examining the effects of change in perioperative CL after ACDF demonstrated no significant difference in VAS arm or neck pain scores among similar maintained, restored, and kyphotic study groups.¹⁹ The authors did report significantly greater improvement

between the restored/maintained groups and restored/kyphotic groups, although the improvement was noted in a different HRQOL outcome parameter (NDI) than that of the present study. Our results include a minimum 1-year follow-up, allowing for longer assessment of cervical sagittal alignment postoperatively compared with the prior study. This is important, considering the observed change in CL that occurs with longer follow-up times.^{20–22} In addition, the larger patient sample of our cohort compared with the aforementioned study (308 vs 104) provides superior power for statistical analysis.¹⁹ Our data suggest that correction of cervical sagittal alignment does contribute to improved clinical outcomes. This finding is corroborated by prior studies, including a prospective, randomized study of 122 patients demonstrating that patients with maintained or improved C2-C7 CL had significantly greater clinical improvement compared with patients with decreased CL postoperatively.^{23–25}

When evaluating the effect of fusion construct length on HRQOL outcomes in patients grouped by their preoperative cervical alignment, multivariate regression demonstrated no significant associations. Other reports have confirmed these findings.^{26,27} In a retrospective cohort study comparing 1-, 2-, and multilevel ACDF

Table 6. Health-related quality-of-life outcomes in patients with preoperative cervical lordosis greater than 10° (lordotic).

Outcome Measure	1-Level (n = 18)	2-Level (n = 17)	3-Level (n = 29)	P Value	Regression Analysis ^a
Physical component of short form-12					
Preoperative	33.6 (28.7, 38.5)	31.5 (28.3, 34.8)	30.1 (25.2, 34.9)	0.78	β: 2.09 (-2.56, 6.75) P = 0.37
Postoperative	37.0 (30.5, 43.6)	40.9 (35.8, 45.9)	36.9 (32.3, 41.5)	0.47	
Delta	3.6 (-2.2, 9.4)	8.7 (2.8, 14.6)	6.0 (-0.5, 12.5)	0.57	
P value	0.20	0.01	0.01		
Mental component of short form-12					
Preoperative	48.1 (42.2, 54.1)	49.3 (44.6, 54.0)	47.1 (40.5, 53.6)	0.96	β: -1.14 (-6.15, 3.87) P = 0.65
Postoperative	53.0 (47.5, 58.5)	56.4 (51.9, 60.9)	50.7 (46.3, 55.1)	0.15	
Delta	3.6 (-5.0, 12.2)	6.4 (0.3, 12.6)	4.1 (-1.7, 9.8)	0.64	
P value	0.40	0.04	0.31		
Neck Disability Index					
Preoperative	41.1 (28.3, 53.9)	34.4 (26.0, 42.7)	34.2 (27.5, 40.9)	0.64	β: 5.14 (-12.21, 22.49) P = 0.53
Postoperative	18.5 (-24.1, 61.1)	19.0 (0.4, 45.7)	26.2 (16.6, 35.7)	0.62	
Delta	-21.3 (-49.6, 6.9)	-19.0 (-41.0, -0.2)	-7.9 (-25.6, 9.8)	0.70	
P value	0.11	0.18	0.02		
VAS arm					
Preoperative	6.4 (4.8, 7.6)	6.7 (5.6, 7.7)	2.9 (1.1, 4.9)	0.01	β: 0.99 (-0.44, 2.42) P = 0.17
Postoperative	2.7 (1.3, 4.0)	2.2 (0.6, 3.8)	1.7 (0.5, 2.9)	0.20	
Delta	-4.2 (-5.8, -2.6)	-5.5 (-7.4, -3.7)	-1.3 (-3.6, 0.7)	0.01	
P value	0.003	0.02	0.58		
VAS neck					
Preoperative	6.1 (4.5, 7.8)	5.7 (3.7, 7.8)	4.8 (3.1, 6.5)	0.29	β: 0.79 (-0.78, 2.36) P = 0.31
Postoperative	0.6 (-0.1, 1.3)	1.0 (-0.1, 2.1)	2.2 (1.1, 3.4)	0.04	
Delta	-5.5 (-7.1, -3.8)	-5.7 (-7.8, -3.5)	-2.6 (-5.1, -0.1)	0.07	
P value	0.003	0.03	0.02		

Abbreviation: VAS, Visual Analog Scale.

Note: Significance achieved at P < 0.05.

^aRegression analysis controlled for age, body mass index, Charlson Comorbidity Index, and preoperative diagnosis.

fusions, no significant difference in improvement of VAS arm, VAS neck, or NDI scores was found.²⁷ Similarly, another retrospective study comparing patients who underwent 1-, 2-, and 3-level ACDF procedures in terms of HRQOL outcomes reported no significant difference among the fusion groups.²⁶

While our study shows improvements in certain clinical outcomes when considering perioperative change in CL, it is important that providers consider the interplay that exists between CL and other cervical spine alignment parameters. An adequate balance between cervical alignment parameters, such as CL, T1 slope (measured as the angle between the superior T1 endplate and horizontal), and cervical sagittal vertical axis (measured as the horizontal distance from the C2 plumb line to posterior-superior aspect of C7), is essential to maintaining a stable cervical spine and preserving horizontal gaze.^{28,29} Previous studies have established the relationship between these parameters, with cervical sagittal vertical axis inversely correlated with CL and directly correlated with T1 slope.^{21,30} Recently, Goldschmidt et al established a model to use preoperative CL to predict postoperative cervical sagittal vertical axis.³¹

$$cSVA = CH \times \tan \left(\frac{\pi}{180} \times \left(TIS - \frac{CL}{2} \right) \right)$$

where cSVA is the cervical sagittal vertical axis, CH is the cervical height (measured as the vertical distance from the anterior aspect of T1 to upper endplate of C2), T1S is the T1 slope, and CL is the cervical lordosis. This equation ultimately allows surgeons to understand the compensation of C2-C7 CL in relation to T1 slope, as well as predict postoperative cervical sagittal vertical axis to help tailor cervical interbody grafts using patient-specific alignment parameters. Similarly, another retrospective study evaluated 70 preoperative kyphotic patients in an attempt to predict the ideal postoperative CL associated with improved clinical outcomes.³² They reported a ratio of postoperative CL to C7 slope (measured as the angle between the superior C7 endplate and horizontal) greater than 0.7 correlated with better postoperative outcomes. These relationships exemplify the ability to use preoperative cervical alignment parameters to aid in surgical planning and help set patient expectations.

While the current study provides evidence regarding the importance of cervical sagittal correction, it is not without limitations. First, due to the retrospective nature of the study, available HRQOL outcome data and radiographic imaging limited the sample size, thus resulting in a relatively smaller cohort of kyphotic group patients and underpowered subanalysis of the impact of preoperative CL on HRQOL outcomes. Second, variations

in operative techniques and postoperative management exist among surgeons, and confounding is a possible source of bias. Multivariate regression analysis, controlling for patient demographic and surgical case characteristics, was employed to help mitigate this bias. Finally, while the present study focused on the impact of preoperative and perioperative change in cervical sagittal alignment, further information regarding lumbosacral sagittal alignment and its impact on cervical deformity, HRQOL outcomes, and clinical outcomes should be considered in future, high-quality studies.

CONCLUSION

Overall, our study shows restoring CL after ACDF results in significant improvements in PCS-12 and VAS arm scores. The length of fusion construct in patients grouped by preoperative cervical alignment had no significant impact on outcomes. Preoperative planning is an essential part of patient care when considering operative management, and this study provides evidence for the effectiveness of ACDF procedures in patients with varying degrees of cervical alignment.

REFERENCES

1. Saifi C, Fein AW, Cazzulino A, et al. Trends in resource utilization and rate of cervical disc arthroplasty and anterior cervical discectomy and fusion throughout the united states from 2006 to 2013. *Spine J*. 2018;18(6):1022–1029. doi:10.1016/j.spinee.2017.10.072
2. McAnany SJ, Baird EO, Overley SC, Kim JS, Qureshi SA, Anderson PA. A meta-analysis of the clinical and fusion results following treatment of symptomatic cervical pseudarthrosis. *Global Spine J*. 2015;5(2):148–155. doi:10.1055/s-0035-1544176
3. Andresen AK, Paulsen RT, Busch F, Isenberg-Jørgensen A, Carreon LY, MØ A. Patient-reported outcomes and patient-reported satisfaction after surgical treatment for cervical radiculopathy. *Global Spine J*. 2018;8(7):703–708. doi:10.1177/2192568218765398
4. Niu CC, Liao JC, Chen WJ, Chen LH. Outcomes of interbody fusion cages used in 1 and 2-levels anterior cervical discectomy and fusion: titanium cages versus polyetheretherketone (PEEK) cages. *J Spinal Disord Tech*. 2010;23(5):310–316. doi:10.1097/BSD.0b013e3181af3a84
5. Matz PG, Holly LT, Groff MW, et al. Indications for anterior cervical decompression for the treatment of cervical degenerative radiculopathy. *J Neurosurg Spine*. 2009;11(2):174–182. doi:10.3171/2009.3.SPINE08720
6. Schwab F, Patel A, Ungar B, Farcy J-P, Lafage V. Adult spinal deformity-postoperative standing imbalance. *Spine*. 2010;35(25):2224–2231. doi:10.1097/BRS.0b013e3181ee6bd4
7. Glassman SD, Bridwell K, Dimar JR, Horton W, Berven S, Schwab F. The impact of positive sagittal balance in adult spinal deformity. *Spine (Phila Pa 1976)*. 2005;30(18):2024–2029. doi:10.1097/01.brs.0000179086.30449.96
8. Ao S, Liu Y, Wang Y, Zhang H, Leng H. Cervical kyphosis in asymptomatic populations: incidence, risk factors, and its relationship with health-related quality of life. *J Orthop Surg Res*. 2019;14(1):322. doi:10.1186/s13018-019-1351-2
9. Iyer S, Lenke LG, Nemani VM, et al. Variations in occipitocervical and cervicothoracic alignment parameters based on age. *Spine*. 2016;41(23):1837–1844. doi:10.1097/BRS.0000000000001644
10. Park MS, Kelly MP, Lee D-H, Min W-K, Rahman RK, Riew KD. Sagittal alignment as a predictor of clinical adjacent segment pathology requiring surgery after anterior cervical arthrodesis. *Spine J*. 2014;14(7):1228–1234. doi:10.1016/j.spinee.2013.09.043
11. Theologis AA, Iyer S, Lenke LG, Sides BA, Kim HJ, Kelly MP. Cervical and cervicothoracic sagittal alignment according to roussouly thoracolumbar subtypes. *Spine (Phila Pa 1976)*. 2019;44(11):E634–E639. doi:10.1097/BRS.0000000000002941
12. Cole T, Veeravagu A, Zhang M, Ratliff JK. Surgeon procedure volume and complication rates in anterior cervical discectomy and fusions. *Clin Spine Surg*. 2017;30(5):E633–E639. doi:10.1097/BSD.0000000000000238
13. Marawar S, Girardi FP, Sama AA, et al. National trends in anterior cervical fusion procedures. *Spine (Phila Pa 1976)*. 2010;35(15):1454–1459. doi:10.1097/BRS.0b013e3181bef3cb
14. Joubari MF, Le Huec JC, Ranjbar Hameghavandi MH, et al. Comparison of cervical sagittal parameters among patients with neck pain and healthy controls: a comparative cross-sectional study. *Eur Spine J*. 2019;28(10):2319–2324. doi:10.1007/s00586-019-06117-8
15. Neuman BJ, Harris A, Jain A, et al. Reciprocal changes in cervical alignment after thoracolumbar arthrodesis for adult spinal deformity. *Spine (Phila Pa 1976)*. 2019;44(22):E1311–E1316. doi:10.1097/BRS.0000000000003159
16. Shaw KA, Murphy JS. Compensatory or pathologic? Cervical spine sagittal alignment in adolescent idiopathic scoliosis. *J Spine Surg*. 2019;5(1):174–175. doi:10.21037/jss.2019.01.01
17. Gao K, Zhang J, Lai J, et al. Correlation between cervical lordosis and cervical disc herniation in young patients with neck pain. *Medicine (Baltimore)*. 2019;98(31). doi:10.1097/MD.00000000000016545
18. Lippa L, Lippa L, Cacciola F. Loss of cervical lordosis: what is the prognosis? *J Craniovertebr Junction Spine*. 2017;8(1):9–14. doi:10.4103/0974-8237.199877
19. Hu X, Ohnmeiss DD, Zigler JE, Guyer RD, Lieberman IH. Restoration of cervical alignment is associated with improved clinical outcome after one and two level anterior cervical discectomy and fusion. *Int J Spine Surg*. 2015;9:61. doi:10.14444/2061
20. Been E, Shefi S, Soudack M. Cervical lordosis: the effect of age and gender. *Spine J*. 2017;17(6):880–888. doi:10.1016/j.spinee.2017.02.007
21. Iorio J, Lafage V, Lafage R, et al. The effect of aging on cervical parameters in a normative north american population. *Global Spine J*. 2018;8(7):709–715. doi:10.1177/2192568218765400
22. Guo G-M, Li J, Diao Q-X, et al. Cervical lordosis in asymptomatic individuals: a meta-analysis. *J Orthop Surg Res*. 2018;13(1):147. doi:10.1186/s13018-018-0854-6
23. Kapetanakis S, Thomaidis T, Charitoudis G, Pavlidis P, Theodosiadis P, Gkasdaris G. Single anterior cervical discectomy and fusion (ACDF) using self-locking stand-alone polyetheretherketone (PEEK) cage: evaluation of pain and health-related quality of life. *J Spine Surg*. 2017;3(3):312–322. doi:10.21037/jss.2017.06.21
24. Chang H, Baek D-H, Choi B-W. The relationship between increased intervertebral disc height and development of postoperative axial neck pain after anterior cervical fusion. *J Korean Neurosurg Soc*. 2014;55(6):343–347. doi:10.3340/jkns.2014.55.6.343

25. Villavicencio AT, Babuska JM, Ashton A, et al. Prospective, randomized, double-blind clinical study evaluating the correlation of clinical outcomes and cervical sagittal alignment. *Neurosurgery*. 2011;68(5):1309–1316. doi:10.1227/NEU.0b013e31820b51f3

26. Lee S-H, Lee JC, Tauchi R, Daniel Riew K. Influence of the number of cervical fusion levels on cervical spine motion and health-related quality of life. *Spine (Phila Pa 1976)*. 2016;41(8):E474–80. doi:10.1097/BRS.0000000000001299

27. Basques BA, Louie PK, Mormol J, et al. Multi- versus single-level anterior cervical discectomy and fusion: comparing sagittal alignment, early adjacent segment degeneration, and clinical outcomes. *Eur Spine J*. 2018;27(11):2745–2753. doi:10.1007/s00586-018-5677-y

28. Diebo BG, Challier V, Henry JK, et al. Predicting cervical alignment required to maintain horizontal gaze based on global spinal alignment. *Spine (Phila Pa 1976)*. 2016;41(23):1795–1800. doi:10.1097/BRS.0000000000001698

29. Staub BN, Lafage R, Kim HJ, et al. Cervical mismatch: the normative value of T1 slope minus cervical lordosis and its ability to predict ideal cervical lordosis. *J Neurosurg Spine*. 2019;30(1):31–37. doi:10.3171/2018.5.SPINE171232

30. Hyun S-J, Kim K-J, Jahng T-A, Kim H-J. Clinical impact of T1 slope minus cervical lordosis after multilevel posterior cervical fusion surgery: a minimum 2-year follow up data. *Spine (Phila Pa 1976)*. 2017;42(24):1859–1864. doi:10.1097/BRS.0000000000002250

31. Goldschmidt E, Angriman F, Agarwal N, et al. A new piece of the puzzle to understand cervical sagittal alignment: utilizing a

novel angle δ to describe the relationship among T1 vertebral body slope, cervical lordosis, and cervical sagittal alignment. *Neurosurgery*. 2020;86(3):446–451. doi:10.1093/neuros/nyz088

32. Ajello M, Marengo N, Pilloni G, et al. Is it possible to evaluate the ideal cervical alignment for each patient needing surgery? An easy rule to determine the appropriate cervical lordosis in preoperative planning. *World Neurosurg*. 2017;97:471–478. doi:10.1016/j.wneu.2016.09.110

Funding: The authors received no financial support for the research, authorship, and/or publication of this article.

Declaration of Conflicting Interests: The authors report no conflicts of interest in this work.

Corresponding Author: Brian A. Karamian, Rothman Orthopaedic Institute at Thomas Jefferson University, 925 Chestnut St, 5th floor, Philadelphia, PA 19107, USA; brian.karamian@rothmanortho.com

Published 14 July 2022

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2022 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.