

External Validation of the National Surgical Quality Improvement Program Calculator Utilizing a Single Institutional Experience for Adult Spinal Deformity Corrective Surgery

Sara Naessig, Katherine Pierce, Waleed Ahmad, Lara Passfall, Oscar Krol, Nicholas A. Kummer, Tyler Williamson, Bailey Imbo, Peter Tretiakov, Kevin Moattari, Rachel Joujon-Roche, Jack Zhong, Eaman Balouch, Brooke O'Connell, Constance Maglaras, Bassel Diebo, Renaud Lafage, Virginie Lafage, Shaleen Vira, Steven Hale, Michael Gerling, Themistocles Protopsaltis, Aaron Buckland and Peter G. Passias

Int J Spine Surg published online 15 February 2023
<https://www.ijssurgery.com/content/early/2023/02/15/8419>

This information is current as of May 9, 2025.

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://ijssurgery.com/alerts>

External Validation of the National Surgical Quality Improvement Program Calculator Utilizing a Single Institutional Experience for Adult Spinal Deformity Corrective Surgery

SARA NAESSIG, BS¹; KATHERINE PIERCE, BS¹; WALEED AHMAD, MS¹; LARA PASSFALL, BS¹; OSCAR KROL, BA¹; NICHOLAS A. KUMMER, BS¹; TYLER WILLIAMSON, MS¹; BAILEY IMBO, BA¹; PETER TRETIKOV, BS¹; KEVIN MOATTARI, BS¹; RACHEL JOUJON-ROCHE, BS¹; JACK ZHONG, BS¹; EAMAN BALOUCH, MD, PhD¹; BROOKE O'CONNELL, MS¹; CONSTANCE MAGLARAS, PhD¹; BASSEL DIEBO, MD²; RENAUD LAFAGE, MS³; VIRGINIE LAFAGE, PhD³; SHALEEN VIRA, MD¹; STEVEN HALE, MD¹; MICHAEL GERLING, MD¹; THEMISTOCLES PROTOPSALTIS, MD¹; AARON BUCKLAND, MD¹; AND PETER G. PASSIAS, MD¹

¹Department of Orthopaedic Surgery, NYU Langone Orthopedic Hospital, New York, NY, USA; ²Department of Orthopedic Surgery, SUNY Downstate, New York, NY, USA; ³Department of Orthopaedics, Lenox Hill Hospital, Northwell Health, New York, NY, USA

ABSTRACT

Background: Identify the external applicability of the American College of Surgeons' National Surgical Quality Improvement Program (NSQIP) risk calculator in the setting of adult spinal deformity (ASD) and subsets of patients based on deformity and frailty status.

Methods: ASD patients were isolated in our single-center database and analyzed for the shared predictive variables displayed in the NSQIP calculator. Patients were stratified by frailty (not frail <0.03, frail 0.3–0.5, severely frail >0.5), deformity [T1 pelvic angle (TPA) > 30, pelvic incidence minus lumbar lordosis (PI-LL) > 20], and reoperation status. Brier scores were calculated for each variable to validate the calculator's predictability in a single center's database (Quality). External validity of the calculator in our ASD patients was assessed via Hosmer-Lemeshow test, which identified whether the differences between observed and expected proportions are significant.

Results: A total of 1606 ASD patients were isolated from the Quality database (48.7 years, 63.8% women, 25.8 kg/m²); 33.4% received decompressions, and 100% received a fusion. For each subset of ASD patients, the calculator predicted lower outcome rates than what was identified in the Quality database. The calculator showed poor predictability for frail, deformed, and reoperation patients for the category "any complication" because they had Brier scores closer to 1. External validity of the calculator in each stratified patient group identified that the calculator was not valid, displaying *P* values >0.05.

Conclusion: The NSQIP calculator was not a valid calculator in our single institutional database. It is unable to comment on surgical complications such as return to operating room, surgical site infection, urinary tract infection, and cardiac complications that are typically associated with poor patient outcomes. Physicians should not base their surgical plan solely on the NSQIP calculator but should consider multiple preoperative risk assessment tools.

Level of Evidence: 3.

Lumbar Spine

Keywords: ASD, spine, deformity, spine surgery, thoracolumbar, NSQIP, calculator

INTRODUCTION

The current health care environment is increasingly emphasizing the need for proper risk stratification that can not only be applied to a wide range of specialties but can also be utilized for specific procedures given a patient's preoperative disposition. There are many such programs that aim to link patient outcomes from surgery to provider reimbursement, some of which are organized by the Centers for Medicare and Medicaid Services, such as pay for performance and

physician quality reporting system. Such programs are constantly being integrated into clinical practice to minimize patient outcomes as well as decrease hospital costs.^{1,2} The intention of these programs is to be able to create a risk stratification model that facilitates appropriate risk-adjusted profiles for individual patients preoperatively with a certain predictability of potential surgical complications. Such risk assessment tools that are customizable to the patient have been shown to be more powerful than generic predictive models.³

The novel American College of Surgeons' National Surgical Quality Improvement Program (NSQIP) risk calculator was created using data from more than 500 hospitals to aid preoperative risk stratification of patients undergoing major surgery. This calculator is accessible to the public online (<https://riskcalculator.facs.org/RiskCalculator/>), is inconclusive of all surgical specialties, and has been previously validated.⁴ More specifically, the calculator uses 21 patient-specific variables as well as a current procedural terminology (CPT) codes for the patient's specific procedure in order to generate a predicted risk for the 11 complication categories. The National Quality Forum has previously advocated that this is a viable tool to assess individual risk for numerous specialties.⁵ Utilization of the calculator in a spine cohort has been previously studied, with Veeravague et al finding that the calculator consistently underestimated complication occurrence.⁶ McCarthy et al found that in cervical patients undergoing fusions, the calculator was only predictive of overall complication occurrence and discharge status as it was unable to accurately predict complications on a more granular basis.⁷

Despite the increasing research of the calculator's predictability in various surgical specialties, there has yet to be a study that utilizes the calculator in an adult spinal deformity (ASD) cohort. The current study aimed to validate the calculator's applicability in a single institution ASD cohort for all of the shared outcomes predicted by the risk stratification tool.

MATERIALS AND METHODS

Study Design and Data Sources

This study is a single-center prospectively collected retrospectively analyzed validation cohort study. The single-center database (Quality) contains spine patients presenting to a single academic institution from September 2011 to June 2018. Institutional Review Board approval was obtained. Inclusion criteria consisted of age >18 years, operative treatment for ASD, with available radiographic, surgical, and health-related quality of life data. ASD was defined as scoliosis $\geq 20^\circ$, sagittal vertical axis ≥ 5 cm, pelvic tilt $\geq 25^\circ$, or thoracic kyphosis $\geq 60^\circ$ and undergoing ≥ 4 -level fusions.

ASD patients from NSQIP were analyzed from 2005 to 2016. The NSQIP database is an initiative developed by the Veterans Health Administration to track the risk-adjusted outcomes of surgical patients. NSQIP collects and tracks patient demographics, preoperative risk factors, CPT coding, International Classification of

Table 1. NSQIP predicted variables.

NSQIP Calculator
Predicted Variables

1. Serious complication
2. Any complication^a
3. Pneumonia
4. Cardiac complication^a
5. Surgical site infection^a
6. Urinary tract infection^a
7. Venous thromboembolism
8. Renal failure
9. Readmission
10. Return to OR^a
11. Death^a
12. Discharge to nursing or rehabilitation facility
13. Length of stay^a

Abbreviations: NSQIP, National Surgical Quality Improvement Program; OR, operating room.

^aIndicates variables were shared between the 2 cohorts.

Disease 9th Edition coding, surgical information, and 30-day perioperative outcomes from randomly assigned patients at participating hospitals. Online Supplemental Appendix A displays the CPT/International Classification of Disease 9th Edition codes used to define our ASD cohort.

Using NSQIP Calculator

There are a total of 13 postoperative variables that are predicted by the NSQIP calculator as listed in Table 1. However, between our single institution Quality database and the NSQIP database, there were a total of 7 postoperative variables shared and labeled in Table 1. In order to utilize the NSQIP calculator in our single institution, we collected the baseline demographic data such as age, sex, functional status, emergency case, American Society of Anesthesiologists class, steroid uses, ascites prior, system sepsis prior, ventilator-dependent, disseminated cancer, diabetes, hypertension, congestive heart failure, dyspnea, smoker status, history of severe chronic obstructive pulmonary disease, dialysis, acute renal failure, and body mass index (BMI).

Statistical Analysis

ASD patients were isolated in NSQIP according to their CPT code (Online Supplemental Appendix A). These patients were then stratified according to the frailty status as developed by Miller et al based on the standard procedure published by Searle et al (not frail <0.03, frail 0.3–0.5, severely frail >0.5),^{8,9} deformity graded by T1 pelvic angle (TPA) >30° and pelvic incidence minus lumbar lordosis (PI-LL) >20,¹⁰ and

Table 2. Distribution of patients in the single center database (Quality) and the National Surgical Quality Improvement Program.

Patient Characteristics	Quality (n = 1606)	NSQIP (n = 58,790)
	Distribution	Distribution
Age, y	48.6 ± 23.6	57.3 ± 13.1
Sex, women	63.8%	51.8%
Body mass index	25.8 ± 6.8	30.5
Ascites	1.1%	<0.01%
Sepsis	1.0%	1.2%
Disseminated cancer	3.1%	2.0%
Diabetes	5.2%	16.5%
Hypertension	15.1%	50.8%
COPD	2.8%	4.6%
Dialysis	1.2%	0.4%

Abbreviation: COPD, chronic obstructive pulmonary disease.

Note: Data presented as % or mean ± SD.

Table 3. Patient actual vs the National Surgical Quality Improvement Program predicted outcomes.

Patient Actual Outcomes	Quality (n = 1606)		
	Actual	Predicted	Brier Score
Any complication	16.5%	11.54%	0.01
Cardiac complication	1.9%	0.34%	2.4336×10^{-4}
Surgical site infection	1.0%	2.0%	1.08×10^{-4}
Urinary tract infection	1.7%	1.6%	1×10^{-8}
Return to operating room	2.4%	3.9%	2.25×10^{-4}
Death	0.0%	0.28%	7.84×10^{-6}
Length of stay, d, mean	6.3	3.5	

reoperation status. Individual scores were calculated for each of the above groupings for all the analyzed CPT codes and averaged to create the calculators “predicted” value. Patients in the Quality database were then analyzed for the rate of each of the 7 outcomes listed in Table 1. Brier scores were then calculated for each variable in order to validate the calculator’s predictability in Quality. The Brier score is a quadratic scoring rule to measure the distance between observed and predicted risk. It is calculated as the sum of squared differences between the binary outcome (Y) and the predicted risk (p): $(Y - p)^2$. Having a score closer to 1 and >0.05 means the NSQIP calculator is a poor predictive tool for that specific outcome. A score closer to 0 means the NSQIP calculator was a predictive tool for that factor.

Table 4. Calculator’s predictability for frail patients.

Patient Outcomes	Frail (n = 126)			Not Frail (n = 1480)		
	Actual	Predicted	Brier Score	Actual	Predicted	Brier Score
Any complication	71.4%	17.08%	0.37773316	9.4%	11.54%	0.00045796
Cardiac complication	12.7%	0.42%	0.01527696	0.9%	0.34%	0.00003136
Surgical site infection	7.1%	2.7%	0.00126736	0.5%	2.04%	0.00023716
Urinary tract infection	9.5%	2.27%	0.00609961	1.1%	1.69%	0.00003481
Return to operating room	14.3%	5.2%	0.0016	1.4%	3.9%	0.000625
Death	0.0%	0.32%	0.00000784	0.0%	0.28%	0.00000784
Length of stay, d, mean	6.3	3.5		6.3	3.5	

External Validation of NSQIP Calculator

The Hosmer-Lemeshow test was performed to determine whether the differences between observed and expected proportions are significant. A large P value indicates that the difference between the number of observed and expected values is insignificant, and the model is therefore considered valid. If the P value is smaller than the specified level of significance ($P < 0.05$), the difference between the number of observed and expected values is statistically significant, and the model is therefore considered not valid.

RESULTS

Cohort Overview

A total of 1606 ASD patients were isolated from the Quality database (48.7 years, 63.8% women, 25.8 kg/m²). 33.4% received decompressions, and 100% received a fusion. 15.1% of the Quality patients had past medical history of hypertension, 3.1% malignant cancer, 5.2% diabetes, 2.6% connective tissue disease, and 2.8% chronic pulmonary disease (Table 2). All of the patients included were without metastatic spine disease and recovered from their solid organ malignancy.

Outcomes Between Quality and NSQIP Patients

The average ASD outcome predicted by the NSQIP risk calculator predicted lower rates for NSQIP patients for return to operating room (0.8% vs 2.4%), length of stay (3.5 vs 6.5 days), total complication rate (11.5% vs 16.5%), and cardiac complications (0.34% vs 1.9%) than Quality patients. The single institution did have lower urinary tract infection and surgical site infection outcomes (1.7% vs 2.85%; 1% vs 1.8%, respectively). The calculated Brier scores identified the calculator’s predictability for each factor is displayed in Table 3. As identified by scores <0.05 , all of the variables had great predictability when used in a single institution cohort.

Table 5. The National Surgical Quality Improvement Program calculator predictability by deformed TPA.

Patient Outcomes	High TPA (n = 40)			Low TPA		
	Actual	Predicted	Brier Score	Actual	Predicted	Brier Score
Any complication	42.5%	8.6%	0.114921	24.6%	11.54%	0.01705636
Cardiac complication	10.0%	0.5%	0.009025	10.0%	0.34%	0.00933156
Surgical site infection	2.5%	1.8%	0.000049	3.3%	2.0%	0.00015876
Urinary tract infection	10.0%	1.4%	0.007396	1.6%	1.6%	8.1E-07
Return to operating room	7.5%	3.3%	0.001764	6.6%	3.9%	0.000729
Death	0	0.0%	0.00000784	0	0.28%	0.00000784
Length of stay, d, mean	8.2	4		6.6	3.5	

Abbreviation: TPA, T1 pelvic angle.

NSQIP Calculator in Frail Patients

When analyzed by frailty status, 7.6% of ASD were categorized as frail while 92.4% were not frail. By basic demographics, frail patients were older (65.2 vs 46.9 years), had a larger BMI (32.2 vs 25.2 kg/m²), and had a greater Charlson Comorbidity Index (CCI) (2.9 vs 0.2; all $P < 0.05$). These differences were adjusted in order to properly identify the calculator's predictability. For not frail patients, all the variables were predictive with the NSQIP calculator displaying appropriate Brier scores. However, for frail patients, the calculator did not accurately predict "any complications" displaying the highest Brier score of 0.3 (Table 4).

NSQIP Calculator in Deformed Patients

Patients who had a high TPA (>30) were older (66.7 vs 32.6 years), had a higher BMI (30 vs 24.9), and had a greater CCI (1.9 vs 0.6; all $P < 0.05$) than those who had a low TPA. The same baseline demographic differences were identified for patients with high PI-LL (>20) and low PI-LL: age (63.2 vs 42.6), BMI (30.4 vs 26.3), and CCI (1.9 vs 1.0; all $P < 0.05$). Adjusting for these baseline differences, the calculator displayed the same poor predictability for "any complications" as shown in Table 5 and Table 6 for their TPA and PI-LL deformity, respectively.

NSQIP Calculator in Reoperation Patients

Of the 1606 ASD Quality patients who were isolated, 10.8% required a reoperation. There were no differences in basic demographics among these ASD reoperation patients and therefore did not require adjustment to the NSQIP calculator. As compared with the previously identified predicted values shown in Table 3 and Table 7, the NSQIP calculator accurately predicted cardiac complication, surgical site infection, urinary tract infection, return to operating room, and death. However, it had a Brier score >0.05 for "any complication," indicating poor predictability for this variable.

External Validation of NSQIP Calculator

After performing the Hosmer-Lemeshow test, the NSQIP calculator was not valid in a single institution for ASD patients when stratified by frailty ($\chi^2 = 587.4$; $P = 8.1 \times 10^{-126}$), high TPA ($\chi^2 = 38.9$, $P = 6.9 \times 10^{-8}$), high PI-LL ($\chi^2 = 43.9$, $P = 6.4 \times 10^{-9}$), and reoperations ($\chi^2 = 54.8$; $P = 3.9 \times 10^{-11}$).

DISCUSSION

Perioperative metrics such as the NSQIP risk calculator have become increasingly utilized to assess surgical risk in various fields in order to ensure quality improvement.¹¹⁻¹³ Such preoperative tools enable providers to avoid any potential risk a patient may have by incorporating a pre-treatment plan to minimize these character metrics. Schenker et al found that risk calculators provide surgeons with

Table 6. The National Surgical Quality Improvement Program calculator predictability by deformed PI-LL.

Patient Outcomes	High PI-LL (n = 65)			Low PI-LL		
	Actual	Predicted	Brier Score	Actual	Predicted	Brier Score
Any complication	38.5%	8.8%	0.088209	24.1%	11.54%	0.01577536
Cardiac complication	4.6%	0.057%	0.00207025	3.6%	0.34%	0.00106276
Surgical site infection	3.1%	1.6%	0.000225	1.8%	2.0%	5.76E-06
Urinary tract infection	7.7%	1.83%	0.00344569	1.8%	1.6%	0.00000121
Return to operating room	7.7%	3.44%	0.001849	5.4%	3.9%	0.000225
Death	0	0.28%	0.00000784	0	0.28%	0.00000784
Length of stay, d, mean	7.4	3.5		6.5	3.5	

Abbreviation: PI-LL, pelvic incidence minus lumbar lordosis.

Table 7. The National Surgical Quality Improvement Program predictability of reoperation patients.

Patient Outcomes	Quality Reoperation (n = 168)		
	Actual	Predicted	Brier Score
Any complication	28.6%	11.54%	0.02910436
Cardiac complication	3.0%	0.34%	0.00070756
Surgical site infection	4.2%	2.0%	0.00046656
Urinary tract infection	5.4%	1.6%	0.00137641
Return to operating room	6%	3.9%	0.000441
Death	0%	0.28%	0.00000784
Length of stay, d, mean	7.7	3.5	

improved preoperative morbidity and mortality estimates thus improving the informed consent process. The attention on these assessments is a result of the use complication occurrence as a proxy for the quality of care within public reporting efforts.¹⁴

In the current study, we evaluated the predictive utility of the NSQIP surgical risk calculator in a single institution data for ASD patients in general and various subsets of patients. We identified that the NSQIP calculator has poor predictability for “any complications” in patients with a deformed TPA and PI-LL as well as frail patients and those undergoing a reoperation. Although the calculator has been validated in recent literature,^{15,16} the findings are for a select surgical population and did not apply to our ASD cohort and subgroups since our models failed to be externally validated according to the Hosmer-Lemeshow test.

Currently, the applicability of the NSQIP risk calculator in spine-specific patients is limited. In the studies that have utilized the calculator in spine patients, they have identified that the calculator consistently predicts lower rates than those that area actually observed in the population.^{6,14} This is impart due to the calculator’s inability to accurately assess a patient’s risk profile regardless of the planned procedure. The calculator’s postoperative risk equation is only based on basic demographics and presurgical comorbidities and does not account for a patient’s frailty status, preoperative deformity, and past surgical history. As identified by Cho et al, reoperation of ASD patients inherently predisposes patients to an elevated risk of complications. However, they identified several risk factors to contribute to these outcomes such as fusion length, type of osteotomy, and preoperative radiographic measurements.¹⁷ The use of the modified Frailty Index has also been identified to be related with postoperative complications with higher modified Frailty Index to be associated with an increased risk of 30-day postoperative complications.¹⁸ With the calculator’s lack of taking these risk factors into account, the usability of the risk calculator in a spine-specific ASD population should be combined with other preoperative risk stratification models.

Nonetheless, the role of a preoperative risk assessment tool should not be discouraged in the light of these results. These tools encourage comprehensive preoperative discussion with the patient and create a patient-centered treatment plan. Using the NSQIP calculator in conjunction with other preoperative risk assessment tools can aid in postoperative care adjusting for patient factors at baseline in order to optimize outcomes as previously reported.^{19,20} Given the current health care era being more patient-centric, it is imperative to be cognizant of patient characteristics that may lead to the development of impairments and worse patient satisfactions. Proper use of stratifying spine patients by taking into account the identified factors in this study can ensure appropriate patient care thus minimizing the patient’s financial burden.

This study was not without limitations. First, our single-center data were obtained through retrospective review, which represents inherent limitations and the introduction of biases including the potential for provider selection to confound results. Second, the reoperation rate of our cohort was relatively higher in our study (10.8%) compared with others, which may have contributed to the elevated observed overall complication rate among cohorts. However, despite these limitations, these results provide valuable discussion on the use of such risk stratification tools such as the NSQIP risk calculator in surgical specialties taking into consideration baseline patient characteristics.

CONCLUSIONS

The NSQIP calculator is not a valid calculator in our single institutional database. It is unable to comment on surgical complications, such as return to surgery and cardiac complications that are typically associated with poor patient outcomes.

REFERENCES

1. Khanduja K, Scales DC, Adhikari NKJ. Pay for performance in the intensive care unit—opportunity or threat? *Crit Care Med*. 2009;37(3):852–858. doi:10.1097/CCM.0b013e3181962b0b
2. Cologne KG, Hwang GS, Senagore AJ. Cost of practice in a tertiary/quaternary referral center: is it sustainable? *Tech Coloproctol*. 2014;18(11):1035–1039. doi:10.1007/s10151-014-1175-3
3. Kwok AC, Lipsitz SR, Bader AM, Gawande AA. Are targeted preoperative risk prediction tools more powerful? A test of models for emergency colon surgery in the very elderly. *J Am Coll Surg*. 2011;213(2):220–225. doi:10.1016/j.jamcollsurg.2011.04.025
4. Gray DT, Deyo RA, Kreuter W, et al. Population-based trends in volumes and rates of ambulatory lumbar spine surgery. *Spine (Phila Pa 1976)*. 2006;31(17):1957–1963. doi:10.1097/01.brs.0000229148.63418.c1

5. Lafage V, Schwab F, Vira S, Patel A, Ungar B, Farcy J-P. Spino-pelvic parameters after surgery can be predicted. *Spine*. 2011;36(13):1037–1045. doi:10.1097/BRS.0b013e3181eb9469
6. Veeravagu A, Li A, Swinney C, et al. Predicting complication risk in spine surgery: a prospective analysis of a novel risk assessment tool. *J Neurosurg Spine*. 2017;27(1):81–91. doi:10.3171/2016.12.SPINE16969
7. McCarthy MH, Singh P, Maslak J, et al. Can the American College of surgeons risk calculator predict 30-day complications after cervical spine surgery? *Clinical Spine Surgery*. 2019;32(9):357–362. doi:10.1097/BSD.0000000000000890
8. Miller EK, Neuman BJ, Jain A, et al. An assessment of frailty as a tool for risk stratification in adult spinal deformity surgery. *Neurosurg Focus*. 2017;43(6):E3. doi:10.3171/2017.10.FOCUS17472
9. Searle SD, Mitnitski A, Gahbauer EA, Gill TM, Rockwood K. A standard procedure for creating a frailty index. *BMC Geriatr*. 2008;8:24. doi:10.1186/1471-2318-8-24
10. Schwab F, Farcy J-P, Bridwell K, et al. A clinical impact classification of scoliosis in the adult. *Spine (Phila Pa 1976)*. 2006;31(18):2109–2114. doi:10.1097/01.brs.00000231725.38943.ab
11. Finks JF, Kole KL, Yenumula PR, et al. Predicting risk for serious complications with bariatric surgery: results from the michigan bariatric surgery collaborative. *Ann Surg*. 2011;254(4):633–640. doi:10.1097/SLA.0b013e318230058c
12. Nam RK, Kattan MW, Chin JL, et al. Prospective multi-institutional study evaluating the performance of prostate cancer risk calculators. *J Clin Oncol*. 2011;29(22):2959–2964. doi:10.1200/JCO.2010.32.6371
13. Gupta PK, Gupta H, Sundaram A, et al. Development and validation of a risk calculator for prediction of cardiac risk after surgery. *Circulation*. 2011;124(4):381–387. doi:10.1161/CIRCULATIONAHA.110.015701
14. Wang X, Hu Y, Zhao B, Su Y. Predictive validity of the ACS-NSQIP surgical risk calculator in geriatric patients undergoing lumbar surgery. *Medicine (Baltimore)*. 2017;96(43):e8416. doi:10.1097/MD.00000000000008416
15. Mogal HD, Fino N, Clark C, Shen P. Comparison of observed to predicted outcomes using the ACS NSQIP risk calculator in patients undergoing pancreaticoduodenectomy. *J Surg Oncol*. 2016;114(2):157–162. doi:10.1002/jso.24276
16. Bilimoria KY, Liu Y, Paruch JL, et al. Development and evaluation of the universal ACS NSQIP surgical risk calculator: a decision aid and informed consent tool for patients and surgeons. *J Am Coll Surg*. 2013;217(5):833–842. doi:10.1016/j.jamcollsurg.2013.07.385
17. Cho SK, Bridwell KH, Lenke LG, et al. Major complications in revision adult deformity surgery: risk factors and clinical outcomes with 2- to 7-year follow-up. *Spine (Phila Pa 1976)*. 2012;37(6):489–500. doi:10.1097/BRS.0b013e3182217ab5
18. Ali R, Schwalb JM, Nerenz DR, Antoine HJ, Rubinfeld I. Use of the modified frailty index to predict 30-day morbidity and mortality from spine surgery. *J Neurosurg Spine*. 2016;25(4):537–541. doi:10.3171/2015.10.SPINE14582
19. Hamilton DF, Lane JV, Gaston P, et al. What determines patient satisfaction with surgery? A prospective cohort study

of 4709 patients following total joint replacement. *BMJ Open*. 2013;3(4):1–7. doi:10.1136/bmjopen-2012-002525

20. Osorio JA, Scheer JK, Ames CP. Predictive modeling of complications. *Curr Rev Musculoskelet Med*. 2016;9(3):333–337. doi:10.1007/s12178-016-9354-7

Funding: The authors received no financial support for the research, authorship, and/or publication of this article.

Declaration of Conflicting Interests: The authors report no conflicts of interest in this work.

Disclosures: Renaud Lafage reports stock/stock options from Nemaris. Virginie LaFage reports royalties/licenses from Nuvasive; consulting for Alphatec Spine and Globus Medical; paid presenter or speaker for DuPuy, Stryker, and the Permanente Medical Group; and leadership roles for the Scoliosis Research Society, International Spine Study Group, and European Spine Journal. Michael C. Gerling reports royalties from Integrity Implants; consulting fees from Integrity Implants, RTI Surgical, and Wolf Endoscopic; and a leadership role with AAOS and the Cervical Spine Research Society. Themistocles Stavros Protopsaltis reports royalties from Altus; consulting fees from Globus Medical, Medtronic, Nuvasive, and Stryker; and stock/stock options from Spine Align and Torus Medical. Aaron James Buckland reports consulting fees from Medtronic, Nuvasive, and Stryker. Nuvasive: Paid consultant. Stryker: Paid consultant. Peter G. Passias reports consulting fees from Medtronic, Medtronic, Royal Biologics, SpineWave, and Terumo; paid presenter/speaker from Globus Medical and Zimmer; research support from the Cervical Scoliosis Research Society; leadership role with Spine; and other financial or material support from Allosource, Cerapedics, and Spinevision.

Corresponding Author: Peter G. Passias, Division of Spinal Surgery/Departments of Orthopaedic and Neurosurgery, NYU Medical Center, NY Spine Institute, 301 East 17th St, New York, NY 10003, USA; peter.passias@nyumc.org

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2023 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.