

# Baseline Neck Disability Index and Patient-Reported Outcomes Measurement Information System Physical Function Predict Postoperative Return to Normal in Cervical Spine Surgery

Gregory S. Kazarian, Michael E. Steinhaus, Sravisht Iyer, Drake LeBrun, Robert Cecere, Takashi Hirase, Francis Lovecchio, Todd J. Albert, Darren Lebl, Darren Huang, Harvinder Sandhu, Bernard Rawlins, Frank Schwab, Virginie Lafage and Han Jo Kim

*Int J Spine Surg* published online 9 October 2024  
<https://www.ijssurgery.com/content/early/2024/10/09/8653>

This information is current as of May 9, 2025.

---

**Email Alerts** Receive free email-alerts when new articles cite this article. Sign up at:  
<http://ijssurgery.com/alerts>

# Baseline Neck Disability Index and Patient-Reported Outcomes Measurement Information System Physical Function Predict Postoperative Return to Normal in Cervical Spine Surgery

GREGORY S. KAZARIAN, MD<sup>1</sup>; MICHAEL E. STEINHAUS, MD<sup>1</sup>; SRAVISHT IYER, MD<sup>1</sup>; DRAKE LEBRUN, MD<sup>1</sup>; ROBERT CECERE, BS<sup>1</sup>; TAKASHI HIRASE, MD<sup>1</sup>; FRANCIS LOVECCHIO, MD<sup>1</sup>; TODD J. ALBERT, MD<sup>1</sup>; DARREN LEBL, MD<sup>1</sup>; DARREN HUANG, MD<sup>1</sup>; HARVINDER SANDHU, MD<sup>1</sup>; BERNARD RAWLINS, MD<sup>1</sup>; FRANK SCHWAB, MD<sup>1</sup>; VIRGINIE LAFAGE, PhD<sup>1</sup>; AND HAN JO KIM, MD<sup>1</sup>

<sup>1</sup>Department of Orthopedic Surgery, Hospital for Special Surgery, New York, NY, USA

## ABSTRACT

**Background:** Recent studies assessing the importance of various preoperative factors on postoperative outcomes following spine surgery have uncovered several important variables that influence subjective and objective outcomes following cervical spine surgery, but it is still unclear which patients are most likely to benefit from operative management.

**Purpose:** The objective of this study was to assess whether preoperative patient-reported outcome measures (PROMs) can be used to predict which patients achieve “normal” levels of pain and function after surgery.

**Study Design:** This was a prospective cohort study.

**Patient Sample:** This study included all adult patients undergoing cervical spine surgery by 1 of 7 senior spine surgeons at our institution between 2016 and 2018. Of the 164 patients who were eligible for 6-month follow-up at the time that study data were collected, 139 had available follow-up data and were included in our analysis.

**Outcomes Measures:** Patients completed the Neck Disability Index (NDI) as well as the Patient-Reported Outcomes Measurement Information System (PROMIS) Physical Function (PF) and Pain Interference computer adaptive tests preoperatively and at 6 months postoperatively.

**Methods:** Patients who achieved postoperative patient-acceptable symptom state (PASS) for NDI ( $\leq 17$ ) and the normative mean (50) for PROMIS were identified. The relationship between preoperative PROMs and the probability of achieving PASS and the normative mean was assessed.

**Results:** One hundred thirty-nine patients met inclusion criteria with diagnoses of myelopathy ( $n = 36$ ), radiculopathy ( $n = 48$ ), and myeloradiculopathy ( $n = 49$ ). For NDI, a 1-point worsening in the preoperative score resulted in an OR of achieving PASS of 0.96 ( $P < 0.001$ ) in the overall population. This association held true for patients with radiculopathy (OR 0.96;  $P = 0.022$ ) but not myelopathy (OR 0.98;  $P = 0.35$ ). For PROMIS PF, a 1-point improvement in the preoperative score resulted in an OR of achieving the normative mean of 1.10 ( $P < 0.001$ ). This association held true for patients with radiculopathy (OR 1.14;  $P = 0.033$ ) but did not reach statistical significance for patients with myelopathy (OR 1.03;  $P = 0.515$ ).

**Conclusions:** Preoperative PROMs can predict postoperative benefit for patients undergoing cervical spine surgery, with worse baseline function associated with a lower likelihood of attaining PASS for NDI and the normative mean for PROMIS PF, especially for patients with radiculopathy.

**Clinical Relevance:** Baseline symptoms and function, including myelopathy or radiculopathy-dominant symptoms and preoperative PROMs, may predict postoperative outcomes.

**Level of Evidence:** 3.

Cervical Spine

Keywords: cervical spine, PROMs, myelopathy, radiculopathy, PROMIS, PASS

## INTRODUCTION

Degenerative disease of the cervical spine is 1 of the most common pathologies encountered by spine

surgeons. When conservative treatments like physical therapy and steroid injections fail, surgical treatment is often indicated to alleviate refractory pain, radiculopathy, and myelopathy. Among the various

surgical options used to treat degenerative cervical pathology, anterior cervical discectomy and fusion (ACDF) is the most common, with roughly 132,000 ACDFs being performed annually in the United States between 2006 and 2013,<sup>1</sup> and far more are likely to be performed in the current decade. Surgical treatments for cervical degenerative diseases have a long track record of successful outcomes.<sup>2–9</sup> Despite this track record, postoperative pain, disability, and dissatisfaction remain significant issues following surgical treatment of cervical spine pathology.<sup>10–12</sup>

In recent decades, there has been an increased effort to study and identify preoperative predictors of adverse outcomes following cervical spine surgery. Such studies are crucial not only for uncovering potentially modifiable factors that influence outcomes but also for driving surgical indications by identifying the patient populations that are most likely to achieve a benefit from surgical intervention. One important area of focus is the study of how preoperative symptom severity and disability level impact postoperative outcomes, which to date has shown mixed results. Three prior studies have reported that preoperative disability portends worse outcomes following surgery for myelopathy.<sup>13–15</sup> Conversely, Fehlings et al and the AO North America multicenter study noted no difference in outcomes when stratifying patients by baseline functional status. In contrast to these prior studies, they note that patients with severe myelopathy actually experienced the greatest improvement in modified Japanese Orthopaedic Association (mJOA) scores.<sup>16</sup> Other preoperative factors, including axial neck pain,<sup>17,18</sup> increased preoperative cord signal,<sup>19</sup> ossification of the posterior longitudinal ligament,<sup>20</sup> duration of symptoms,<sup>21,22</sup> advanced age,<sup>23,24</sup> and lower preoperative mJOA score,<sup>24</sup> have also been identified as predictors of poor postoperative outcomes.

Recent studies assessing the importance of various preoperative factors on postoperative outcomes following spine surgery have uncovered several important variables that influence subjective and objective outcomes following surgery. However, additional studies are needed in order to further characterize the patient populations that are most likely to benefit from operative management. While studies have established that patients with poor preoperative function can benefit from surgery, to what extent patients can return to a “normal” level of function remains unknown. In the current study,

we assessed how preoperative patient-reported outcomes measures (PROMs), namely Neck Disability Index (NDI) and Patient-Reported Outcomes Measurement Information System (PROMIS), impact the probability of achieving a satisfactory level of function postoperatively.

## MATERIALS AND METHODS

### Study Design and Patient Population

Institutional Review Board approval was obtained prior to the initiation of the study (number 2015–363). Informed consent was obtained. This study included all adult patients undergoing cervical spine surgery by 1 of 7 senior spine surgeons at our institution between 2016 and 2018. Inclusion criteria were (1) age  $\geq 18$  years, (2) English speaking, and (3) undergoing cervical spine surgery. Exclusion criteria included (1) non-English speaking, (2) undergoing surgical stabilization for cervical instability secondary to trauma, and (3) infection.

### Outcome Measures and Analysis

Patients completed the NDI as well as the PROMIS Physical Function (PF) and Pain Interference (PI) computer adaptive tests preoperatively and at 6 months postoperatively. For the NDI, the threshold of reaching a satisfactory level of function was determined using the patient acceptable symptom state (PASS). Unlike minimal clinically important difference (MCID), which is the minimal change in score that is perceptible by the patient, PASS is defined as the target value on a PROM scale beyond which patients deem themselves to have attained an acceptable outcome. A recent study determined the PASS threshold for NDI as being  $\leq 17$ .<sup>25</sup> For the PROMIS subscales, the normative mean value of 50 was used as the threshold for attaining an acceptable postoperative state. Using the Assessment Center ([www.assessmentcenter.net](http://www.assessmentcenter.net)), preoperative and postoperative NDI and PROMIS PF and PI surveys were administered to the patients enrolled in the current study. Questions were asked in a randomized order to avoid bias associated with question order and survey fatigue.

All statistical analysis for this study was performed using STATA 15.0 (Stata Corp, College Station, TX, USA). Descriptive statistics were used to describe demographic and clinical characteristics.  $\chi^2$  tests were used to identify differences between groups in the proportion of patients achieving the normative mean and PASS. Univariate logistic regressions were used to determine the odds of achieving the binary outcomes under study (attaining PASS vs not attaining pass;

attaining normative mean vs not attaining normative mean) associated with incremental 1-point increases in preoperative NDI and PROMIS scores in the overall population and diagnosis subgroups (myelopathy and radiculopathy). A 2-sided type I error rate of 0.05 was used to determine statistical significance. The proportions of patients achieving PASS and the normative mean were determined, with preoperative outcome scores stratified by quintiles for PROMIS and levels of disability for NDI.<sup>26</sup>

## RESULTS

Of the 164 patients who were eligible for 6-month follow-up at the time that study data were collected, 139 had available follow-up data and were included in our analysis. Six-month follow-up was selected as the minimum, as multiple studies have demonstrated that PROMIS plateau by 3–6 months following spine surgery.<sup>27–30</sup> Of these 139 patients, 83 (59.7%) were men, and the mean age was 56.4 years (range, 24–85). Patients had differing indications for surgery, including isolated radiculopathy ( $n = 48$ , 34.5%), isolated myelopathy ( $n = 36$ , 25.9%), or myeloradiculopathy ( $n = 49$ , 35.3%) (Table 1). The anterior approach was used in 102 patients (73.4%), while the posterior approach was utilized in 37 (26.6%).

### Impact of Preoperative PROMS on Achieving PASS or the Normative Mean

The proportion of patients who met NDI PASS with preoperative NDI scores of 0 to 4 (no disability), 5 to 15 (mild disability), 15 to 24 (moderate disability), 25 to 34 (severe disability), and >34 (complete disability) is illustrated in Figure 1. When stratified by baseline level of disability, the proportion of patients reaching PASS was 83%, 100%, 80%, 57%, and 52% for baseline NDI 0 to 4, 5 to 15, 15 to 24, 25 to 34, and >34, respectively. This difference between groups was statistically significant ( $P = 0.005$ ). In Figures 2 and 3, the proportion of patients who met the normative mean for

PROMIS PF and PI, respectively, is shown as a function of the preoperative PROMIS quintile score. For PROMIS PF, the proportion of patients reaching the normative threshold was 62%, 46%, 30%, 33%, and 26% for baseline PF 47.7 to 66.2, 43.3 to 47.2, 40.1 to 43.1, 34.7 to 40.0, and 23.5 to 34.6, respectively. This difference between groups was statistically significant ( $P = 0.035$ ). For PROMIS PI, the proportion of patients reaching the normative threshold was 69%, 48%, 70%, 64%, and 74%. This difference between groups was not statistically significant.

For NDI, a 1-point increase (worse NDI) in the preoperative score resulted in OR of achieving PASS (NDI  $\leq 17$ ) of 0.96 ( $P < 0.001$ ). This association held true for patients with radiculopathy (OR 0.96;  $P = 0.022$ ) but did not reach statistical significance for patients with myelopathy (OR 0.98;  $P = 0.35$ ). For PROMIS PF, a 1-point increase (improved PROMIS) in the preoperative score resulted in OR of achieving the normative mean (PROMIS  $\geq 50$ ) of 1.10 ( $P < 0.001$ ). This association held true for patients with radiculopathy (OR 1.14;  $P = 0.033$ ) but did not reach statistical significance for patients with myelopathy (OR 1.03;  $P = 0.515$ ).

**Patients Achieving PASS (NDI)**

		PASS Achieved after Surgery	
		Met PASS	Did not meet PASS
NDI Preoperative Score	0 to 4	83%	17%
	5 to 15	100%	0%
	15 to 24	80%	20%
	25 to 34	57%	43%
	>34	52%	48%

**Figure 1.** Proportion of patients meeting NDI PASS shown as a function of preoperative NDI. Differences between these groups were statistically significant ( $P = 0.005$ ). NDI, Neck Disability Index; PASS, patient acceptable symptom state.

**Table 1.** Association between preoperative outcomes and postoperative clinical benefit.

Surgery Indication	PROMIS PF		PROMIS PI		NDI	
	Preoperative	Postoperative Improvement	Preoperative	Postoperative Improvement	Preoperative	Postoperative Improvement
Radiculopathy	39.2 (8.1)	5.2 (8.0)	60.4 (7.8)	−8.9 (10.8)	29.4 (22.0)	13.4 (17.8)
Myelopathy	41.7 (7.4)	9.0 (9.4)	61.9 (6.0)	−10.3 (9.5)	39.2 (2.2)	22.2 (17.6)
Myeloradiculopathy	43.0 (8.4)	5.6 (7.2)	59.9 (7.3)	−6.9 (9.1)	31.6 (19.5)	14.4 (14.5)

Abbreviations: NDI, Neck Disability Index; PROMIS PF, Patient-Reported Outcomes Measurement Information System Physical Function; PROMIS PI, Patient-Reported Outcomes Measurement Information System Pain Interference.

Note: For Neck Disability Index, postoperative improvements reflect a decrease in score, as lower scores reflect lower disability.



**Patients Achieving PROMIS PF  $\geq 50$** 

		PROMIS $\geq 50$ Achieved after Surgery	
		Met normative mean	Did not meet normative mean
PROMIS PF Preoperative Score	Better health 47.7 – 66.2	62%	38%
	43.3 – 47.2	46%	54%
	40.1 – 43.1	30%	70%
	34.7 – 40.0	33%	67%
	Worse health 23.5 – 34.6	26%	74%

**Figure 2.** Proportion of patients attaining the normative means ( $\geq 50$ ) for PROMIS PF shown as a function of preoperative PROMIS PF quintile. Differences between these groups were statistically significant ( $P = 0.035$ ). PROMIS PF, Patient-Reported Outcomes Measurement Information System Physical Function.

(Table 2). PROMIS PI did not demonstrate statistically significant correlations between worse baseline scores and achieving postoperative PASS and the normative mean.

**Patients Achieving PROMIS PI  $\leq 50$** 

		PROMIS $\leq 50$ Achieved after Surgery	
		Met normative mean	Did not meet normative mean
PROMIS PI Preoperative Score	Better health 38.7 – 54.4	69%	31%
	54.6 – 59.0	48%	52%
	59.1 – 62.6	70%	30%
	62.9 – 66.6	64%	36%
	Worse health 66.9 – 83.8	74%	26%

**Figure 3.** Proportion of patients attaining the normative means ( $\leq 50$ ) for PROMIS PI shown as a function of preoperative PROMIS PI quintile. Differences between these groups were not statistically significant. PROMIS PI, Patient-Reported Outcomes Measurement Information System Pain Interference.

**Table 2.** Association between preoperative diagnosis and outcome.

Outcome	OR	95% CI	P
Overall			
NDI	0.96	0.94, 0.98	<0.001
PROMIS PI	1.04	0.99, 1.09	0.144
PROMIS PF	1.1	1.04, 1.16	<0.001
Radiculopathy			
NDI	0.96	0.92, 0.99	0.022
PROMIS PI	1.03	0.95, 1.12	0.491
PROMIS PF	1.14	1.01, 1.29	0.033
Myelopathy			
NDI	0.98	0.94, 1.02	0.35
PROMIS PI	1.06	0.96, 1.18	0.262
PROMIS PF	1.03	0.95, 1.11	0.515

Abbreviations: NDI, Neck Disability Index; PASS, patient acceptable symptom state; PF, Physical Function; PI, Pain Interference; PROMIS, Patient-Reported Outcomes Measurement Information System.

## DISCUSSION

While the clinical benefit of surgical intervention for refractory pain or myelopathy associated with degenerative cervical spine disorders has been clearly demonstrated, a significant proportion of patients still experience suboptimal outcomes following surgery. Improving our understanding of the extent to which preoperative factors impact postoperative outcomes is a critical step for surgeons in indicating patients for surgery and counseling them about the expected prognosis. In the current study, we assessed the impact of preoperative NDI and PROMIS PF and PI scores on the probability of attaining postoperative acceptable levels of function, using PASS for NDI and the normative mean for PROMIS. We found that worse preoperative function was associated with a lower probability of returning to “normal” postoperatively. When assessed as a function of preoperative diagnosis, we demonstrated that while this relationship holds true for patients with primarily radicular symptoms, there is no significant relationship between baseline preoperative NDI or PROMIS score and the probability of returning to “normal” for patients with primarily myelopathic symptoms.

While there are numerous studies that have assessed outcomes following cervical spine surgery, there are a limited number of studies utilizing preoperative symptoms or disability level to predict the degree of postoperative function, disability, or likelihood of obtaining a clinically significant improvement in symptoms. One example of a preoperative symptom that has been extensively studied with respect to its impact on postoperative outcomes is neck vs radicular arm pain in patients with cervical spine pathology.<sup>17,18,31</sup> Though this topic is well studied, the results are fairly mixed. While the majority of studies have identified neck disability as a predictor of poor outcomes following

ACDF,<sup>32–34</sup> others have found the opposite, namely that higher levels of preoperative neck pain and disability are positive predictors of treatment success.<sup>35,36</sup> These mixed results may be partially explained by the fact that various studies use different methods of characterizing surgical benefit, such as improvement from baseline and overall postoperative PROMs. Though the etiology of these differences is not completely clear, the results of the current study appear to corroborate the findings of the former studies described here, in which a lower degree of preoperative disability is associated with a better postoperative outcome and a higher probability of achieving a “normal” level of function.

Interestingly, while we found a relationship between better preoperative PROMs and the likelihood of achieving a clinically meaningful postoperative outcome for patients with radiculopathy, this relationship did not hold true for patients with predominantly myelopathic symptoms. While the explanation for this outcome is not completely understood, it may be due to differences in the natural history and treatment goals for these 2 different conditions. Because of the presumed progressive and stepwise nature of myelopathic symptoms, surgical treatment for myelopathy is largely focused on preventing the worsening of this condition, and postoperative improvements have been thought to be less predictable. The mixed results seen in this population are likely reflective of the less predictable response to surgery seen in patients with myelopathy. While several studies showed a correlation between greater preoperative disability and worse postoperative outcomes,<sup>13–15</sup> others reported the opposite.<sup>16,25</sup> For example, Fehlings et al found that patients with myelopathy achieved MCID for Nurick, NDI, SF-36, and mJOA.<sup>16</sup> Additionally, Goh et al found that significant portions of patients undergoing surgical treatment for myelopathy attain MCID using the Japanese Orthopaedic Association score.<sup>37</sup>

One important point to address is that patients with no preoperative disability (NDI 0–4) had a nonsignificant trend toward a lower chance of achieving PASS than those with mild preoperative disability (NDI 5–14). While we do not recommend the use of NDI as a screening tool for surgical candidates, this finding may be important for preoperative counseling. In patients with no preoperative disability, there is a slightly lower chance of achieving PASS. However, this likely varies on a patient-to-patient basis. Because disability is a subjective experience, patients with severe symptoms that justify surgical intervention may not have a significant disability if they are highly

resilient and vice versa. Also, in specific instances like myelopathy, symptoms may be minimal despite severe cord compression. These patients may elect to undergo surgical intervention despite no disability in order to prevent future deterioration.

This study had many limitations. First, while our sample size is comparable to prior studies on this topic, it limits a robust assessment of subgroups due to limited power. Second, different patients included in this study were treated using either the anterior or posterior approach, which was not controlled for and may have had subtle impacts on the results, as these patients may have different recovery trajectories at our final endpoint. Third, the follow-up time of 6 months limits the assessment of any long-term differences in patient outcomes. Fourth, although the outcome measures used in this study are well-validated, they were not devised specifically for the assessment of myelopathy. Therefore, they may not have been as sensitive in detecting improvements in myelopathic symptoms, which may explain why no significant results were identified for patients with myelopathy. It is possible that different results would be found with the use of the Japanese Orthopedic Association myelopathy assessment. Future studies should focus on patient-specific assessments of satisfaction to determine individualized thresholds for PASS based on preoperative symptoms and preoperative symptom severity.

## CONCLUSIONS

To our knowledge, this is the first report assessing the impact of preoperative PROMs on the ability for patients to return to “normal” functional levels after cervical spine surgery. Here, we show that preoperative NDI and PROMIS PF are predictive of postoperative benefit, with patients at higher levels of preoperative function being more likely to attain this normative threshold postoperatively. This held true for the overall cohort, as well as for patients with the underlying diagnosis of radiculopathy, but not myelopathy. These results provide invaluable insights for patients and surgeons alike in better understanding the prognosis for patients undergoing operative management for cervical degenerative disease, especially patients with primarily radicular symptoms.

## REFERENCES

1. Saifi C, Fein AW, Cazzulino A, et al. Trends in resource utilization and rate of cervical disc arthroplasty and anterior cervical discectomy and fusion throughout the United States from 2006 to 2013. *Spine J*. 2018;18(6):1022–1029. doi:10.1016/j.spinee.2017.10.072

2. Ebersold MJ, Pare MC, Quast LM. Surgical treatment for cervical spondylitic myelopathy. *J Neurosurg*. 1995;82(5):745–751. doi:10.3171/jns.1995.82.5.0745
3. Bolesta MJ, Rechline GR, Chrin AM. Three- and four-level anterior cervical discectomy and fusion with plate fixation: a prospective study. *Spine*. 2000;25(16):2040–2044. doi:10.1097/00007632-200008150-00007
4. Vaccaro AR, Balderston RA. Anterior plate instrumentation for disorders of the subaxial cervical spine. *Clin Orthop Relat Res*. 1997;1997(335):112–121.
5. Zaveri GR, Ford M. Cervical spondylosis: the role of anterior instrumentation after decompression and fusion. *J Spinal Disord*. 2001;14(1):10–16. doi:10.1097/00002517-200102000-00003
6. Yue WM, Brodner W, Highland TR. Long-term results after anterior cervical discectomy and fusion with allograft and plating: a 5- to 11-year radiologic and clinical follow-up study. *Spine*. 1976;30(19):2138–2144. doi:10.1097/01.brs.0000180479.63092.17
7. Kaiser MG, Haid RW, Subach BR, Barnes B, Rodts GE. Anterior cervical plating enhances arthrodesis after discectomy and fusion with cortical allograft. *Neurosurgery*. 2002;50(2):229–236. doi:10.1097/00006123-200202000-00001
8. Samartzis D, Shen FH, Lyon C, Phillips M, Goldberg EJ, An HS. Does rigid instrumentation increase the fusion rate in one-level anterior cervical discectomy and fusion? *Spine J*. 2004;4(6):636–643. doi:10.1016/j.spinee.2004.04.010
9. Wang JC, McDonough PW, Kanim LEA, Endow KK, Delamarter RB. Increased fusion rates with cervical plating for three-level anterior cervical discectomy and fusion. *Spine*. 2001;26(6):643–646. doi:10.1097/00007632-200103150-00015
10. Wang H, Zhang D, Ma L, Shen Y, Ding W. Factors predicting patient dissatisfaction 2 years after discectomy for lumbar disc herniation in a Chinese older cohort: a prospective study of 843 cases at a single institution. *Medicine*. 2015;94(40). doi:10.1097/MD.0000000000001584
11. Asher AL, Devin CJ, Kerezoudis P, et al. Predictors of patient satisfaction following 1- or 2-level anterior cervical discectomy and fusion: insights from the quality outcomes database. *J Neurosurg Spine*. 2019;31(6):835–843. doi:10.3171/2019.6.SPINE19426
12. Badhiwala JH, Khan O, Wegner A, et al. A partial least squares analysis of functional status, disability, and quality of life after surgical decompression for degenerative cervical myelopathy. *Sci Rep*. 2020;10(1). doi:10.1038/s41598-020-72595-2
13. Karpova A, Arun R, Davis AM, et al. Predictors of surgical outcome in cervical spondylotic myelopathy. *Spine*. 2013;38(5):392–400. doi:10.1097/BRS.0b013e3182715bc3
14. Tetreault LA, Kopjar B, Vaccaro A, et al. A clinical prediction model to determine outcomes in patients with cervical spondylotic myelopathy undergoing surgical treatment: data from the prospective, multi-center aospine North America study. *J Bone Joint Surg Am*. 2013;95(18):1659–1666. doi:10.2106/JBJS.L.01323
15. Setzer M, Vrionis FD, Hermann EJ, Seifert V, Marquardt G. Effect of apolipoprotein E genotype on the outcome after anterior cervical decompression and fusion in patients with cervical spondylotic myelopathy. *J Neurosurg Spine*. 2009;11(6):659–666. doi:10.3171/2009.7.SPINE08667
16. Fehlings MG, Wilson JR, Kopjar B, et al. Efficacy and safety of surgical decompression in patients with cervical spondylotic myelopathy: results of the aospine North America prospective multi-center study. *J Bone Joint Surg Am*. 2013;95(18):1651–1658. doi:10.2106/JBJS.L.00589
17. DePalma AF, Rothman RH, Lewinnek GE, Canale ST. Anterior interbody fusion for severe cervical disc degeneration. *Surg Gynecol Obstet*. 1972;134(5):755–758.
18. Williams JL, Allen MB, Harkness JW. Late results of cervical discectomy and interbody fusion: some factors influencing the results. *J Bone Joint Surg Am*. 1968;50(2):277–286. doi:10.2106/00004623-196850020-00006
19. Yukawa Y, Kato F, Yoshihara H, Yanase M, Ito K. MR T2 image classification in cervical compression myelopathy: predictor of surgical outcomes. *Spine*. 2007;32(15):1675–1678. doi:10.1097/BRS.0b013e318074d62e
20. Nakashima H, Kanemura T, Kanbara S, et al. What are the important predictors of postoperative functional recovery in patients with cervical OPLL? Results of a multivariate analysis. *Glob Spine J*. 2019;9(3):315–320. doi:10.1177/2192568218794665
21. Mastronardi L, Elsawaf A, Roberto R, et al. Prognostic relevance of the postoperative evolution of intramedullary spinal cord changes in signal intensity on magnetic resonance imaging after anterior decompression for cervical spondylotic myelopathy. *J Neurosurg Spine*. 2007;7(6):615–622. doi:10.3171/SPI-07/12/615
22. Pumberger M, Froemel D, Aichmair A, et al. Clinical predictors of surgical outcome in cervical spondylotic myelopathy: an analysis of 248 patients. *Bone Joint J*. 2013;95-B(7):966–971. doi:10.1302/0301-620X.95B7.31363
23. Zhang JT, Wang LF, Wang S, Li J, Shen Y. Risk factors for poor outcome of surgery for cervical spondylotic myelopathy. *Spinal Cord*. 2016;54(12):1127–1131. doi:10.1038/sc.2016.64
24. Sun LQ, Li M, Li YM. Predictors for surgical outcome of laminoplasty for cervical spondylotic myelopathy. *World Neurosurg*. 2016;94:89–96. doi:10.1016/j.wneu.2016.06.092
25. Goh GS, Yue W-M, Guo C-M, Tan S-B, Chen JL. Defining threshold values on the neck disability index corresponding to a patient acceptable symptom state in patients undergoing elective surgery for degenerative disorders of the cervical spine. *Spine J*. 2020;20(8):1316–1326. doi:10.1016/j.spinee.2020.05.004
26. Vernon H, Mior S. The neck disability index: a study of reliability and validity. *J Manipul Physiol Ther*. 1991;14(7):409–415.
27. Cheung WY, Arvinte D, Wong YW, Luk KDK, Cheung KMC. Neurological recovery after surgical decompression in patients with cervical spondylotic myelopathy - a prospective study. *Int Orthop*. 2008;32(2):273–278. doi:10.1007/s00264-006-0315-4
28. Ayling OGS, Ailon T, McIntosh G, et al. Clinical outcomes research in spine surgery: what are appropriate follow-up times? *J Neurosurg Spine*. 2019;30(3):397–404. doi:10.3171/2018.8.SPINE18715
29. Shahi P, Subramanian T, Tuma O. Temporal trends of improvement after minimally invasive transforaminal lumbar interbody fusion. *Spine*. 2014. doi:10.1097/BRS.0000000000005024
30. Calek AK, Hochreiter B, Buckland AJ. Reassessing the minimum 2-year follow-up standard after lumbar decompression surgery: a 2-month follow-up seems to be an acceptable minimum. *Spine J*. 2024;24(7):1244–1252. doi:10.1016/j.spinee.2024.03.018
31. Passias PG, Hasan S, Radcliff K, et al. Arm pain versus neck pain: a novel ratio as a predictor of post-operative clinical outcomes in cervical radiculopathy patients. *Int J Spine Surg*. 2018;12(5):629–637. doi:10.14444/5078
32. Peolsson A, Peolsson M. Predictive factors for long-term outcome of anterior cervical decompression and fusion: a multivariate data analysis. *Eur Spine J*. 2008;17(3):406–414. doi:10.1007/s00586-007-0560-2



33. Peolsson A, Hedlund R, Vavruch L, Oberg B. Predictive factors for the outcome of anterior cervical decompression and fusion. *Eur Spine J*. 2003;12(3):274–280. doi:10.1007/s00586-003-0530-2
34. Peolsson A, Vavruch L, Oberg B. Predictive factors for arm pain, neck pain, neck specific disability and health after anterior cervical decompression and fusion. *Acta Neurochir*. 2006;148(2):167–173. doi:10.1007/s00701-005-0660-x
35. Hermansen A, Hedlund R, Vavruch L, Peolsson A. Positive predictive factors and subgroup analysis of clinically relevant improvement after anterior cervical decompression and fusion for cervical disc disease: a 10- to 13-year follow-up of a prospective randomized study: clinical article. *J Neurosurg Spine*. 2013;19(4):403–411. doi:10.3171/2013.7.SPINE12843
36. Anderson PA, Subach BR, Riew KD. Predictors of outcome after anterior cervical discectomy and fusion. *Spine*. 2009;34(2):161–166. doi:10.1097/BRS.0b013e31819286ea
37. Goh GS-H, Liow MHL, Ling ZM, et al. Severity of pre-operative myelopathy symptoms affects patient-reported outcomes, satisfaction, and return to work after anterior cervical discectomy and fusion for degenerative cervical myelopathy. *Spine*. 2020;45(10):649–656. doi:10.1097/BRS.0000000000003354

**Funding:** Virginie Lafage received a grant (paid to her institution) from CSRS for the present manuscript. Sravisht Iyer reports support from Innovasis.

**Declaration of Conflicting Interests:** The authors report no conflicts of interest in this work.

**Disclosures:** Sravisht Iyer reports consulting fees from Stryker; payment/honoraria from Globus Medical; participation on a Data Safety Monitoring or advisory board for Healthgrades; and other financial/nonfinancial interests in HS2 LLC. Todd J. Albert reports grants from JP Medical Publishers, Theime Medical Publishers, Springer, and Elsevier; royalties/licenses from Biomet (now Zimmer Biomet) and DePuy Synthes Spine; consulting fees from NuVasive; and other financial/nonfinancial interests in Back Story LLC, Innovative Surgical Designs, Inc, InVivo Therapeutics,

Spinicity, CytoDyn Inc., Paradigm Spine LLC, HS2 LLC, Stathspey Crown, Parvizi Surgical Innovation LLC, Surg.IO LLC, Care Equity, Augmedics, Morphogenesis, Precision Orthopedics, Pulse Equity, and Physician Recommended Nutraceuticals. Darren R. Lebl reports royalties from NuVasive and Stryker; consulting fees from DePuy Synthes and Stryker; participation on an advisory board for Remedy Logic; and ownership interest in HS2 LLC, ISPH II LLC, Remedy Logic, Vestia Ventures MiRus Investment LLC, Viseon, and Woven Orthopedic Technologies. Virginie Lafage reports a grant from the International Spine Study Group (paid to her institution); royalties/licenses from NuVasive; consulting fees from Globus Medical and Alphatec; payment/honoraria for lectures, presentation, speakers bureaus, manuscript writing, or education events from J&J, Stryker, and Implanet; and leadership roles for the International Spine Study Group and Scoliosis Research Society. Han Jo Kim reports grants/contracts from ISSGF (paid to institution); royalties/licenses from Zimmer Biomet, K2M/Stryker, and Acuity Surgical; consulting fees from NuVasive; participation on a Data Safety Monitoring or advisory board for Vivex Biology and Aspen Medical; and other financial/nonfinancial interests in AO Spine. The remaining authors have nothing to disclose.

**Corresponding Author:** Han Jo Kim, Department of Orthopaedic Surgery, Hospital for Special Surgery, 535 E 70th St, New York, NY, USA; KimH@HSS.edu

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2024 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.